

Additional Online Appendix for “Jackknife Estimation of a Cluster-Sample IV Regression Model with Many Weak Instruments”*

John C. Chao, Norman R. Swanson, and Tiemen Woutersen

First Draft: September 5, 2020 *This Version:* August, 2022

Abstract

This online appendix is comprised of four sections. Section 1 contains proofs for the key lemmas used to show the main theorems of the paper “Jackknife Estimation of a Cluster-Sample IV Regression with Many Weak Instruments.” More specifically, in section 1, we provide proofs for Lemmas S2-1 to S2-18 which are stated without proof in the Supplemental Appendix of the aforementioned paper. In section 2, we provide a proof of Lemma 1 of the main paper. Section 3 of this online appendix provides proofs of some additional lemmas which give further supporting arguments for the proofs presented in section 1. Finally, section 4 gives some additional Monte Carlo results which complement those reported in the main paper.

Section 1: Proof of Lemmas S2-1 to S2-18

In this section of the online appendix, we provide proof of Lemmas S2-1 to S2-18, which have been stated without proof in the Appendix S2 of the Supplemental Appendix to our paper.

Lemma S2-1: Let $A = P^\perp - M^{(Z,Q)}D_{\hat{\vartheta}}M^{(Z,Q)}$. Then, under Assumptions 2-6, the following statements hold as $K_{2,n}$, $n \rightarrow \infty$.

(a)

$$\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 = O_{a.s.}(K_{2,n}).$$

*Corresponding author: John C. Chao, Department of Economics, 7343 Preinkert Drive, University of Maryland, chao@econ.umd.edu. Norman R. Swanson, Department of Economics, 9500 Hamilton Street, Rutgers University, nswanson@econ.rutgers.edu. Tiemen Woutersen, Department of Economics, 1130 E Helen Street, University of Arizona, woutersen@arizona.edu. The authors owe special thanks to Jerry Hausman and Whitney Newey for many discussions on the topic of this paper over a number of years. In addition, thanks are owed to all of the participants of the 2019 MIT Conference Honoring Whitney Newey for comments on and advice given about the topic of this paper. Finally, the authors wish to thank Miriam Arden for excellent research assistance. Chao thanks the University of Maryland for research support, and Woutersen’s work was supported by an Eller College of Management Research Grant.

(b)

$$\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 = O_{a.s.} \left(\frac{K_{2,n}^3}{n^2} \right).$$

(c)

$$\sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s) \\ (k,v) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 = O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right)$$

(d)

$$\max_{1 \leq (i,t) \leq m_n} \left(\sum_{(j,s)=1}^{m_n} A_{(i,t),(j,s)}^2 \right) = O_{a.s.} \left(\frac{K_{2,n}}{n} \right)$$

(e)

$$\begin{aligned} \sum_{i_1,i_2=1}^n \sum_{\substack{j=1 \\ j \neq i_1,i_2}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{t_2=1}^{T_{i_2}} \sum_{s_1,s_2=1}^{T_j} A_{(i_1,t_1),(j,s_1)}^2 A_{(i_2,t_2),(j,s_2)}^2 &= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right), \\ \sum_{i=1}^n \sum_{j_1=1}^n \sum_{\substack{j_2=1 \\ j_1 \neq j_2}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_{j_1}} \sum_{s_1,s_2=1}^{T_{j_2}} A_{(i,t_1),(j_1,s_1)}^2 A_{(i,t_2),(j_2,s_2)}^2 &= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right). \end{aligned}$$

(f)

$$\sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_i} A_{(i,t),(i,s)}^2 = O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right).$$

(g)

$$\sum_{i=1}^n \sum_{\substack{s_1,t_1=1 \\ s_1 \neq t_1}}^{T_i} \sum_{\substack{s_2,t_2=1 \\ s_2 \neq t_2}}^{T_i} A_{(i,t_1),(i,s_1)}^2 A_{(i,t_2),(i,s_2)}^2 = O_{a.s.} \left(\frac{K_{2,n}^4}{n^3} \right).$$

Proof of Lemma S2-1:

To show part (a), note first that, by Lemma OA-1 given in section 3 of this online appendix, we have

$$tr \left\{ D_{\hat{\vartheta}}^2 \right\} = O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right),$$

where $D_{\hat{\vartheta}} = diag \left(\hat{\vartheta}_1, \dots, \hat{\vartheta}_{m_n} \right)$. Now, by straightforward calculations and by making use of the inequality

$$\left| \sum_{i=1}^G a_i \right|^r \leq G^{r-1} \sum_{i=1}^G |a_i|^r, \quad (1)$$

we get

$$\begin{aligned}
\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 &= \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} \left(P_{(i,t),(j,s)}^\perp - e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \\
&\leq 2 \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} \left(P_{(i,t),(j,s)}^\perp \right)^2 + 2 \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \\
&\leq 2 \left[K_{2,n} + \sum_{(i,t)=1}^{m_n} e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \right] \\
&\leq 2 \left[K_{2,n} + \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}}^2 M^{(Z,Q)} \right\} \right] = 2 \left[K_{2,n} + \text{tr} \left\{ D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} \right\} \right] \\
&\leq 2 \left[K_{2,n} + \text{tr} \left\{ D_{\widehat{\vartheta}}^2 \right\} \right] = O_{a.s.}(K_{2,n}) + O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) = O_{a.s.}(K_{2,n}).
\end{aligned}$$

Next, to show part (b), note that, by applying the inequality in expression (1) as well as the

CS inequality and using the fact that $\lambda_{\max}(M^{(Z,Q)}) = 1$, we obtain

$$\begin{aligned}
& \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 \\
= & \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} \left(P_{(i,t),(j,s)}^\perp - e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^4 \\
\leq & 2^3 \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} \left\{ \left(P_{(i,t),(j,s)}^\perp \right)^4 + \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^4 \right\} \\
\leq & 8 \sum_{(i,t),(j,s)=1}^{m_n} \left(P_{(i,t),(j,s)}^\perp \right)^4 \\
& + 8 \sum_{(i,t),(j,s)=1}^{m_n} \left\{ \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}}^2 M^{(Z,Q)} e_{(i,t)} \right) \left(e'_{(j,s)} M^{(Z,Q)} e_{(j,s)} \right) \right. \\
& \quad \left. \times e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} e'_{(j,s)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \right\} \\
\leq & 8 \sum_{(i,t),(j,s)=1}^{m_n} \left(P_{(i,t),(j,s)}^\perp \right)^4 + 8 \max_{1 \leq (i,t) \leq m_n} \left| \widehat{\vartheta}_{(i,t)} \right|^2 \sum_{(i,t)=1}^{m_n} e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \\
\leq & 8 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} \left(P_{(i,t),(j,s)}^\perp \right)^2 \\
& + 8 \left(\max_{1 \leq (i,t) \leq m_n} \left| \widehat{\vartheta}_{(i,t)} \right|^2 \right) \operatorname{tr} \left\{ D_{\widehat{\vartheta}} M^{(Z,Q)} \sum_{(i,t)=1}^{m_n} e_{(i,t)} e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} \right\} \\
\leq & 8 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 K_{2,n} + 8 \left(\max_{1 \leq (i,t) \leq m_n} \left| \widehat{\vartheta}_{(i,t)} \right|^2 \right) \operatorname{tr} \left\{ D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} \right\} \\
\leq & 8 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 K_{2,n} + 8 \left(\max_{1 \leq (i,t) \leq m_n} \left| \widehat{\vartheta}_{(i,t)} \right|^2 \right) \operatorname{tr} \left\{ D_{\widehat{\vartheta}}^2 \right\} \\
= & O_{a.s.} \left(\frac{K_{2,n}^3}{n^2} \right) + O_{a.s.} \left(\frac{K_{2,n}^4}{n^3} \right) \quad (\text{by parts (a) and (b) of Lemma OA-1}) \\
= & O_{a.s.} \left(\frac{K_{2,n}^3}{n^2} \right).
\end{aligned}$$

To show part (c), note that

$$\begin{aligned}
& \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 \\
= & \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s) \\ (k,v) \neq (j,s)}} \left(P_{(i,t),(j,s)}^\perp - e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \left(P_{(j,s),(k,v)}^\perp - e'_{(j,s)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(k,v)} \right)^2 \\
\leq & \sum_{(i,t),(j,s)=1}^{m_n} \left(P_{(i,t),(j,s)}^\perp - e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \\
& \times \sum_{(k,v)=1}^{m_n} \left(P_{(j,s),(k,v)}^\perp - e'_{(j,s)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(k,v)} \right)^2 \\
\leq & 4 \sum_{(i,t),(j,s)=1}^{m_n} \left[\left(P_{(i,t),(j,s)}^\perp \right)^2 + \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \right] \\
& \times \sum_{(k,v)=1}^{m_n} \left[\left(P_{(j,s),(k,v)}^\perp \right)^2 + \left(e'_{(j,s)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(k,v)} \right)^2 \right] \\
& \quad (\text{by the inequality in expression (1)}) \\
= & 4 \sum_{(i,t),(j,s)=1}^{m_n} \left\{ \left[\left(P_{(i,t),(j,s)}^\perp \right)^2 + \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \right] \right. \\
& \quad \left. \times \left[P_{(j,s),(j,s)}^\perp + e'_{(j,s)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right] \right\} \\
\leq & 4 \left(\max_{1 \leq (j,s) \leq m_n} P_{(j,s),(j,s)}^\perp \right) \sum_{(i,t),(j,s)=1}^{m_n} \left[\left(P_{(i,t),(j,s)}^\perp \right)^2 + \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \right] \\
& + 4 \sum_{(i,t),(j,s)=1}^{m_n} \left[\left(P_{(i,t),(j,s)}^\perp \right)^2 + \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \right] e'_{(j,s)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \\
\leq & 4 \left(\max_{1 \leq (j,s) \leq m_n} P_{(j,s),(j,s)}^\perp \right) \sum_{(i,t)=1}^{m_n} \left[P_{(i,t),(i,t)}^\perp + e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \right] \\
& + 4 \left(\max_{1 \leq (j,s) \leq m_n} |\widehat{\vartheta}_{(j,s)}|^2 \right) \sum_{(i,t)=1}^{m_n} \left[P_{(i,t),(i,t)}^\perp + e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \right] \\
\leq & 4 \left[\left(\max_{(j,s) \in \Lambda_2} P_{(j,s),(j,s)}^\perp \right) + \left(\max_{1 \leq (j,s) \leq m_n} |\widehat{\vartheta}_{(j,s)}|^2 \right) \right] \left[K_{2,n} + \text{tr} \left\{ D_{\widehat{\vartheta}}^2 \right\} \right] \\
= & \left[O_{a.s.} \left(\frac{K_{2,n}}{n} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right) \right] \left[O_{a.s.} (K_{2,n}) + O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) \right] \quad (\text{using Lemma OA-1}) \\
= & O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right).
\end{aligned}$$

Finally, to show part (d), note that

$$\begin{aligned}
\sum_{(j,s)=1}^{m_n} A_{(i,t),(j,s)}^2 &= \sum_{(j,s)=1}^{m_n} \left(P_{(i,t),(j,s)}^\perp - e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right)^2 \\
&= \sum_{(j,s)=1}^{m_n} \left(P_{(i,t),(j,s)}^\perp \right)^2 - 2e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \sum_{(j,s)=1}^{m_n} e_{(j,s)} e'_{(j,s)} P^\perp e_{(i,t)} \\
&\quad + e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \sum_{(j,s)=1}^{m_n} e_{(j,s)} e'_{(j,s)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \\
&= P_{(i,t),(i,t)}^\perp + e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \\
&\leq P_{(i,t),(i,t)}^\perp + e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}}^2 M^{(Z,Q)} e_{(i,t)} \\
&\leq P_{(i,t),(i,t)}^\perp + e'_{(i,t)} M^{(Z,Q)} e_{(i,t)} \left(\max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \right) \\
&\leq P_{(i,t),(i,t)}^\perp + \left(\max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \right) \text{ (since } e'_{(i,t)} M^{(Z,Q)} e_{(i,t)} \leq 1)
\end{aligned}$$

It follows that

$$\begin{aligned}
\max_{1 \leq (i,t) \leq m_n} \left(\sum_{(j,s)=1}^{m_n} A_{(i,t),(j,s)}^2 \right) &\leq \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right) + \left(\max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \right) \\
&= O_{a.s.} \left(\frac{K_{2,n}}{n} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right) \\
&= O_{a.s.} \left(\frac{K_{2,n}}{n} \right).
\end{aligned}$$

To show part (e), we apply the inequality (1), Assumption 6, and parts (a) and (b) of Lemma

OA-1 to obtain

$$\begin{aligned}
& \sum_{i_1, i_2=1}^n \sum_{\substack{j=1 \\ j \neq i_1, i_2}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{t_2=1}^{T_{i_2}} \sum_{s_1, s_2=1}^{T_j} A_{(i_1, t_1), (j, s_1)}^2 A_{(i_2, t_2), (j, s_2)}^2 \\
& \leq \sum_{i_1=1}^n \sum_{j=1}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_j} \left(P_{(i_1, t_1), (j, s_1)}^\perp - e'_{(i_1, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_1)} \right)^2 \\
& \quad \times \sum_{i_2=1}^n \sum_{t_2=1}^{T_{i_2}} \sum_{s_2=1}^{T_j} \left(P_{(i_2, t_2), (j, s_2)}^\perp - e'_{(i_2, t_2)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_2)} \right)^2 \\
& \leq 4 \sum_{i_1=1}^n \sum_{j=1}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_j} \left[\left(P_{(i_1, t_1), (j, s_1)}^\perp \right)^2 + \left(e'_{(i_1, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_1)} \right)^2 \right] \\
& \quad \times \sum_{s_2=1}^{T_j} \sum_{i_2=1}^{n} \sum_{t_2=1}^{T_{i_2}} \left[\left(P_{(i_2, t_2), (j, s_2)}^\perp \right)^2 + \left(e'_{(i_2, t_2)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_2)} \right)^2 \right] \\
& = 4 \sum_{i_1=1}^n \sum_{j=1}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_j} \left[\left(P_{(i_1, t_1), (j, s_1)}^\perp \right)^2 + \left(e'_{(i_1, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_1)} \right)^2 \right] \\
& \quad \times \sum_{s_2=1}^{T_j} \left[P_{(j, s_2), (j, s_2)}^\perp + e'_{(j, s_2)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_2)} \right] \\
& \leq 4\bar{T} \left(\max_{1 \leq (i, t) \leq m_n} P_{(i, t), (i, t)}^\perp \right) \sum_{i_1=1}^n \sum_{j=1}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_j} \left[\left(P_{(i_1, t_1), (j, s_1)}^\perp \right)^2 + \left(e'_{(i_1, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_1)} \right)^2 \right] \\
& \quad + 4\bar{T} \left(\max_{1 \leq (i, t) \leq m_n} |\widehat{\vartheta}_{(i, t)}|^2 \right) \sum_{i_1=1}^n \sum_{j=1}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_j} \left[\left(P_{(i_1, t_1), (j, s_1)}^\perp \right)^2 + \left(e'_{(i_1, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(j, s_1)} \right)^2 \right] \\
& \leq 4\bar{T} \left[\left(\max_{1 \leq (i, t) \leq m_n} P_{(i, t), (i, t)}^\perp \right) + \left(\max_{1 \leq (i, t) \leq m_n} |\widehat{\vartheta}_{(i, t)}|^2 \right) \right] \left[K_{2,n} + \text{tr} \left\{ D_{\widehat{\vartheta}}^2 \right\} \right] \\
& = \left[O_{a.s.} \left(\frac{K_{2,n}}{n} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right) \right] \left[O_{a.s.} (K_{2,n}) + O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) \right] \text{ (using Lemma OA-1)} \\
& = O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right)
\end{aligned} \tag{2}$$

Next, note that, since A is a symmetric matrix, we have $A_{(i, t_1), (j_1, s_1)}^2 = A_{(j_1, s_1), (i, t_1)}^2$ and $A_{(i, t_2), (j_2, s_2)}^2 = A_{(j_2, s_2), (i, t_2)}^2$. It follows, by switching the indices $(i, t_1) \rightarrow (j, s_1)$, $(i, t_2) \rightarrow (j, s_2)$, $(j_1, s_1) \rightarrow (i_1, t_1)$,

and $(j_2, s_2) \rightarrow (i_2, t_2)$, and applying the result given in expression (2) above, we have

$$\begin{aligned}
& \sum_{i=1}^n \sum_{\substack{j_1=1 \\ j_1 \neq i}}^n \sum_{\substack{j_2=1 \\ j_2 \neq i}}^n \sum_{t_1, t_2=1}^{T_i} \sum_{s_1=1}^{T_{j_1}} \sum_{s_2=1}^{T_{j_2}} A_{(i, t_1), (j_1, s_1)}^2 A_{(i, t_2), (j_2, s_2)}^2 \\
&= \sum_{i=1}^n \sum_{\substack{j_1=1 \\ j_1 \neq i}}^n \sum_{\substack{j_2=1 \\ j_2 \neq i}}^n \sum_{t_1, t_2=1}^{T_i} \sum_{s_1=1}^{T_{j_1}} \sum_{s_2=1}^{T_{j_2}} A_{(j_1, s_1), (i, t_1)}^2 A_{(j_2, s_2), (i, t_2)}^2 \\
&= \sum_{j=1}^n \sum_{\substack{i_1=1 \\ i_1 \neq j}}^n \sum_{\substack{i_2=1 \\ i_2 \neq j}}^n \sum_{s_1, s_2=1}^{T_j} \sum_{t_1=1}^{T_{i_1}} \sum_{t_2=1}^{T_{i_2}} A_{(i_1, t_1), (j, s_1)}^2 A_{(i_2, t_2), (j, s_2)}^2 \\
&= \sum_{i_1, i_2=1}^n \sum_{\substack{j=1 \\ j \neq i_1, i_2}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{t_2=1}^{T_{i_2}} \sum_{s_1, s_2=1}^{T_j} A_{(i_1, t_1), (j, s_1)}^2 A_{(i_2, t_2), (j, s_2)}^2 \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right).
\end{aligned}$$

Turning our attention to part (f), note that, by applying the inequality (1) and making use of

Assumption 6(ii), we have

$$\begin{aligned}
& \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_i} A_{(i,t),(i,s)}^2 \\
&= \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_i} \left(P_{(i,t),(i,s)}^\perp - e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,s)} \right)^2 \\
&= \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_i} \left\{ \left(P_{(i,t),(i,s)}^\perp \right)^2 - 2P_{(i,t),(i,s)}^\perp e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,s)} \right. \\
&\quad \left. + \left[e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,s)} \right]^2 \right\} \\
&\leq 2 \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_i} \left\{ \left(P_{(i,t),(i,s)}^\perp \right)^2 + \left[e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} e_{(i,s)} \right]^2 \right\} \\
&\leq 2 \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_i} \left\{ P_{(i,t),(i,t)}^\perp P_{(i,s),(i,s)}^\perp + \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}}^2 M^{(Z,Q)} e_{(i,t)} \right) \left(e'_{(i,s)} M^{(Z,Q)} e_{(i,s)} \right) \right\} \\
&= 2 \sum_{i=1}^n \sum_{t=1}^{T_i} P_{(i,t),(i,t)}^\perp \sum_{s=1}^{T_i} P_{(i,s),(i,s)}^\perp \\
&\quad + 2 \sum_{i=1}^n \sum_{t=1}^{T_i} \left(e'_{(i,t)} M^{(Z,Q)} D_{\widehat{\vartheta}}^2 M^{(Z,Q)} e_{(i,t)} \right) \sum_{s=1}^{T_i} e'_{(i,s)} M^{(Z,Q)} e_{(i,s)} \\
&\leq 2\bar{T} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right) \sum_{(i,t)=1}^{m_n} P_{(i,t),(i,t)}^\perp + 2\bar{T} \left(\max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \right) \sum_{(i,t)=1}^{m_n} \left(e'_{(i,t)} M^{(Z,Q)} e_{(i,t)} \right) \\
&\leq 2\bar{T} \left[\left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right) K_{2,n} + \left(\max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \right) (m_n - K_n - n) \right] \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) \quad (\text{by part (b) of Lemma OA-1}) \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right).
\end{aligned}$$

To show part (g), note that, by making use of Assumption 6(ii), we have

$$\begin{aligned}
& \sum_{i=1}^n \sum_{\substack{s_1, t_1=1 \\ s_1 \neq t_1}}^{T_i} \sum_{\substack{s_2, t_2=1 \\ s_2 \neq t_2}}^{T_i} A_{(i, t_1), (i, s_1)}^2 A_{(i, t_2), (i, s_2)}^2 \\
&= \sum_{i=1}^n \sum_{\substack{s_1, t_1=1 \\ s_1 \neq t_1}}^{T_i} \sum_{\substack{s_2, t_2=1 \\ s_2 \neq t_2}}^{T_i} \left\{ \left(P_{(i, t_1), (i, s_1)}^\perp - e'_{(i, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(i, s_1)} \right)^2 \right. \\
&\quad \left. \times \left(P_{(i, t_2), (i, s_2)}^\perp - e'_{(i, t_2)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(i, s_2)} \right)^2 \right\} \\
&\leq 4 \sum_{i=1}^n \sum_{\substack{s_1, t_1=1 \\ s_1 \neq t_1}}^{T_i} \left[\left(P_{(i, t_1), (i, s_1)}^\perp \right)^2 + \left(e'_{(i, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(i, s_1)} \right)^2 \right] \\
&\quad \times \sum_{\substack{s_2, t_2=1 \\ s_2 \neq t_2}}^{T_i} \left[\left(P_{(i, t_2), (i, s_2)}^\perp \right)^2 + \left(e'_{(i, t_2)} M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} e_{(i, s_2)} \right)^2 \right] \quad (\text{using inequality (1)}) \\
&\leq 4 \sum_{i=1}^n \sum_{t_1=1}^{T_i} \left[\left(P_{(i, t_1), (i, t_1)}^\perp \right) \sum_{s_1=1}^{T_i} \left(P_{(i, s_1), (i, s_1)}^\perp \right) + \left(e'_{(i, t_1)} M^{(Z, Q)} D_{\widehat{\vartheta}}^2 M^{(Z, Q)} e_{(i, t_1)} \right) \sum_{s_1=1}^{T_i} \left(e'_{(i, s_1)} M^{(Z, Q)} e_{(i, s_1)} \right) \right] \\
&\quad \times \sum_{\substack{s_2, t_2=1 \\ s_2 \neq t_2}}^{T_i} \left[\left(P_{(i, t_2), (i, t_2)}^\perp \right) \left(P_{(i, s_1), (i, s_1)}^\perp \right) + \left(e'_{(i, t_2)} M^{(Z, Q)} D_{\widehat{\vartheta}}^2 M^{(Z, Q)} e_{(i, t_2)} \right) \left(e'_{(i, s_1)} M^{(Z, Q)} e_{(i, s_1)} \right) \right] \\
&\leq 4\bar{T}^3 \left[\left(\max_{1 \leq (i, t) \leq m_n} P_{(i, t), (i, t)}^\perp \right)^2 + \left(\max_{1 \leq (i, t) \leq m_n} |\widehat{\vartheta}_{(i, t)}|^2 \right) \right] \\
&\quad \times \left[\left(\max_{1 \leq (i, t) \leq m_n} P_{(i, t), (i, t)}^\perp \right) \sum_{(i, t_1)=1}^{m_n} P_{(i, t_1), (i, t_1)}^\perp + \left(\max_{1 \leq (i, t) \leq m_n} |\widehat{\vartheta}_{(i, t)}|^2 \right) \sum_{(i, t_1)=1}^{m_n} e'_{(i, t_1)} M^{(Z, Q)} e_{(i, t_1)} \right] \\
&\leq 4\bar{T}^3 \left[\left(\max_{1 \leq (i, t) \leq m_n} P_{(i, t), (i, t)}^\perp \right)^2 + \left(\max_{1 \leq (i, t) \leq m_n} |\widehat{\vartheta}_{(i, t)}|^2 \right) \right] \\
&\quad \times \left[\left(\max_{1 \leq (i, t) \leq m_n} P_{(i, t), (i, t)}^\perp \right) K_{2,n} + \left(\max_{1 \leq (i, t) \leq m_n} |\widehat{\vartheta}_{(i, t)}|^2 \right) (m_n - K_n - n) \right] \\
&= \left[O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right) \right] \left[O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) \right] \quad (\text{using Lemma OA-1(b)}) \\
&= O_{a.s.} \left(\frac{K_{2,n}^4}{n^3} \right). \quad \square
\end{aligned}$$

Lemma S2-2: Let Assumptions 1-6 be satisfied. Then, the following statements are true:
(a) $D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1} = O_p \left(n (\mu_n^{\min})^{-2} \right)$; (b) $D_\mu^{-1} X' A X D_\mu^{-1} = H_n + o_p(1)$, where $H_n = \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n = O_p(1)$.

Proof of Lemma S2-2:

To show part (a), note first that $D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1}$ $\leq 2 [\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n + D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1}]$, where we take $A \leq B$ for two square matrices A and B to mean that $A - B$ is negative semi-definite, or, alternatively, $B - A$ is positive semidefinite. Now, for $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we can apply the CS inequality and Assumption 3(iii) to obtain $|a' \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon b / n| \leq \sqrt{a' \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon a / n} \sqrt{b' \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon b / n} \leq \sqrt{a' \Upsilon' Z'_2 Z_2 \Upsilon a / n} \sqrt{b' \Upsilon' Z'_2 Z_2 \Upsilon b / n} = O_{a.s.}(1)$. Since the above argument holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we further deduce that $\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n = O_p(1)$.

Next, note that, by the CS inequality, $|a' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} b| \leq \sqrt{a' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} a} \sqrt{b' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} b}$. Now, by Assumptions 2(i), 3(ii), and 6(ii),

$$\begin{aligned} E \left[a' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} a | \mathcal{F}_n^Z \right] &\leq a' D_\mu^{-1} E \left[U' U | \mathcal{F}_n^Z \right] D_\mu^{-1} a \\ &\leq \frac{\bar{T} \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right) n}{(\mu_n^{\min})^2} = O_{a.s.} \left(\frac{n}{(\mu_n^{\min})^2} \right), \end{aligned}$$

where $\bar{T} = \max_{1 \leq (i,t) \leq m_n} T_i$. Hence, by Theorem 16.1 of Billingsley (1995), there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large $E \left[\left((\mu_n^{\min})^2 / n \right) a' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} a \right] = E_Z \left(\left((\mu_n^{\min})^2 / n \right) E \left[\frac{a' U' M^{(Z_1, Q)} U a}{n} | \mathcal{F}_n^Z \right] \right) \leq \bar{C}$. It follows from the Markov's inequality that $a' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} a = O_p \left(n (\mu_n^{\min})^{-2} \right)$. In the same way, we also have $b' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} b = O_p \left(n / (\mu_n^{\min})^2 \right)$, so that $|a' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} b| \leq \sqrt{a' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} a} \sqrt{b' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} b} = O_p \left(n (\mu_n^{\min})^{-2} \right)$. Since this result holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we further deduce that $D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} = O_p \left(n (\mu_n^{\min})^{-2} \right)$. Putting these results together, it follows that $D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1} = O_p(1) + O_p \left(n (\mu_n^{\min})^{-2} \right) = O_p \left(n (\mu_n^{\min})^{-2} \right)$, as required to show part (a).

To show part (b), note that, by applying parts (a)-(c) of Lemma OA-12, we obtain

$$\begin{aligned}
& D_\mu^{-1} X' A X D_\mu^{-1} \\
&= D_\mu^{-1} \left(\frac{D_\kappa \Theta' Z'_1}{\sqrt{n}} + \frac{D_\mu \Upsilon' Z'_2}{\sqrt{n}} + \Xi' Q' + U' \right) A \left(\frac{Z_1 \Theta D_\kappa}{\sqrt{n}} + \frac{Z_2 \Upsilon D_\mu}{\sqrt{n}} + Q \Xi + U \right) D_\mu^{-1} \\
&= D_\mu^{-1} \left(\frac{D_\mu \Upsilon' Z'_2}{\sqrt{n}} + U \right)' A \left(\frac{Z_2 \Upsilon D_\mu}{\sqrt{n}} + U \right) D_\mu^{-1} \quad (\text{since } AZ_1 = 0 \text{ and } AQ = 0) \\
&= \frac{D_\mu^{-1} D_\mu \Upsilon' Z'_2 A Z_2 \Upsilon D_\mu D_\mu^{-1}}{n} + D_\mu^{-1} U' A U D_\mu^{-1} + \frac{D_\mu^{-1} D_\mu \Upsilon' Z'_2 A U D_\mu^{-1}}{\sqrt{n}} \\
&\quad + \frac{D_\mu^{-1} U' A Z_2 \Upsilon D_\mu D_\mu^{-1}}{\sqrt{n}} \\
&= \frac{\Upsilon' Z'_2 A Z_2 \Upsilon}{n} + D_\mu^{-1} U' A U D_\mu^{-1} + \frac{\Upsilon' Z'_2 A U D_\mu^{-1}}{\sqrt{n}} + \frac{D_\mu^{-1} U' A Z_2 \Upsilon}{\sqrt{n}} \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon}{n} + o_p(1).
\end{aligned}$$

Moreover, $\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n = O_p(1)$, as shown in the proof of part (a) above. This shows part (b). \square

Lemma S2-3: Let $\underline{U} = U - \varepsilon \rho'$ and $\underline{U}_{(i,t)} = U_{(i,t)} - \rho \varepsilon_{(i,t)}$ and let $VC(X|\mathcal{F}_n^Z)$ denote the conditional covariance matrix of the random vector X given \mathcal{F}_n^Z . Under Assumptions 1-2, 5-6, and 8; there exists positive constants $0 < \underline{C} \leq \bar{C} < \infty$ such that the following statements are true.

(a) $\lambda_{\max}[VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z)] \leq \bar{C}$ a.s. and $\lambda_{\min}[VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z)] \geq \underline{C}$ a.s. for all n sufficiently large.

(b) $VC(\underline{U}' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z) \geq \underline{C} I_d > \underset{d \times d}{0}$ a.s., for all n sufficiently large.

(c) $\lambda_{\max}(VC[\underline{U}' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z]) \leq \bar{C}$ a.s., $\lambda_{\max}(VC[\underline{U}' A \varepsilon / \sqrt{K_{2,n}}]) \leq \bar{C}$, $\lambda_{\max}(VC[U' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z]) \leq \bar{C}$ a.s., and $\lambda_{\max}(VC[U' A \varepsilon / \sqrt{K_{2,n}}]) \leq \bar{C}$, for all n sufficiently large.

(d) For any $a \in \mathbb{R}^d$ with $\|a\|_2 = 1$ and for all n sufficiently large, $\lambda_{\min}(\Sigma_n) \geq \underline{C} > 0$ a.s. and $a' \Sigma_n^{-1} a \leq \bar{C} < \infty$ a.s., where $\Sigma_n = VC(\mathcal{Y}_n | \mathcal{F}_n^Z) = \Sigma_{1,n} + \Sigma_{2,n}$, as defined in section 4 of the main paper, and where $\mathcal{Y}_n = \Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} + D_\mu^{-1} \underline{U}' A \varepsilon$.

Proof of Lemma S2-3:

For part (a), note that, by Assumptions 1, 2, and 3(iii); there exists a pair of constants $0 < \underline{C} \leq \bar{C} < \infty$ such that, for any $b \in \mathbb{R}^d$ such that $\|b\| = 1$ and for all n sufficiently large, $b' VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z) b = b' \Upsilon' Z'_2 M^{(Z_1, Q)} E[\varepsilon \varepsilon' | \mathcal{F}_n^Z] M^{(Z_1, Q)} Z_2 \Upsilon b / n \leq \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \lambda_{\max} \bar{C}$ a.s. and $b' VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z) b \geq \left(\min_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) b' \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon b / n \geq \underline{C}$ a.s. Since the above bounds hold for any $b \in \mathbb{R}^d$ such that $\|b\|_2 = 1$, it follows that, almost surely, $\lambda_{\max}[VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z)] = \max_{\|b\|=1} b' VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z) b \leq \bar{C} < \infty$ and $\lambda_{\min}[VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z)] = \min_{\|b\|=1} b' VC(\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z) b \geq \underline{C} > 0$ for all n sufficiently large, which establishes the required result.

To show part (b), let $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$, and we define $\underline{U}_{(i,t)} = U_{(i,t)} - \rho \varepsilon_{(i,t)}$,

$\underline{u}_{a,(i,t)} = a' \underline{U}_{(i,t)}, \sigma_{(i,t)}^2 = E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z], \tilde{\omega}_{a,(i,t)}^2 = E[\underline{u}_{a,(i,t)}^2 | \mathcal{F}_n^Z], \tilde{\psi}_{a,(i,t)} = E[\varepsilon_{(i,t)} \underline{u}_{a,(i,t)} | \mathcal{F}_n^Z]$, and $\varrho_{a,(i,t)} = \tilde{\psi}_{a,(i,t)} / (\sigma_{(i,t)} \tilde{\omega}_{a,(i,t)})$; for $(i, t) = 1, \dots, m_n$, where we have suppressed the dependence of $\sigma_{(i,t)}^2, \tilde{\omega}_{a,(i,t)}^2, \tilde{\psi}_{a,(i,t)}$, and $\varrho_{a,(i,t)}$ on $\mathcal{F}_n^Z = \sigma(Z)$ for notational convenience. Note also that we can write $a' V C (\underline{U}' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z) a = K_{2,n}^{-1} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 E\left[\left(\varepsilon_{(j,s)} \underline{u}_{a,(i,t)} + \varepsilon_{(i,t)} \underline{u}_{a,(j,s)}\right)^2 | \mathcal{F}_n^Z\right]$, since, by construction, $A_{(i,t),(i,t)} = 0$ for $(i, t) = 1, \dots, m_n$. Moreover, define $\delta_{a,(i,t),(j,s)} = (\sigma_{(j,s)} \tilde{\omega}_{a,(i,t)} \quad \sigma_{(i,t)} \tilde{\omega}_{a,(j,s)})'$ and

$$\Delta_{(i,t),(j,s)}^a = \begin{pmatrix} 1 & \varrho_{a,(i,t)} \varrho_{a,(j,s)} \\ \varrho_{a,(i,t)} \varrho_{a,(j,s)} & 1 \end{pmatrix}$$

and note that, given that $(j, s) < (i, t)$, we have $E\left[\left(\varepsilon_{(j,s)} \underline{u}_{a,(i,t)} + \varepsilon_{(i,t)} \underline{u}_{a,(j,s)}\right)^2 | \mathcal{F}_n^Z\right] = \delta'_{a,(i,t),(j,s)} \Delta_{(i,t),(j,s)}^a \delta_{a,(i,t),(j,s)}$. Now, by the quadratic formula, the smallest eigenvalue of $\Delta_{(i,t),(j,s)}^a$ is given by $\lambda_{\min}(\Delta_{(i,t),(j,s)}^a) = 1 - |\varrho_{a,(i,t)}| |\varrho_{a,(j,s)}|$. In addition, write

$$\begin{aligned} \tilde{\Omega}_{(i,t)} &= \begin{pmatrix} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] & E[\varepsilon_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z] \\ E[\underline{U}_{(i,t)} \varepsilon_{(i,t)} | \mathcal{F}_n^Z] & E[\underline{U}_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z] \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ -\rho & I_d \end{pmatrix} \begin{pmatrix} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] & E[\varepsilon_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z] \\ E[\underline{U}_{(i,t)} \varepsilon_{(i,t)} | \mathcal{F}_n^Z] & E[\underline{U}_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z] \end{pmatrix} \begin{pmatrix} 1 & -\rho' \\ 0 & I_d \end{pmatrix} \\ &= L_\rho \Omega_{(i,t)} L'_\rho, \text{ where } L_\rho = \begin{pmatrix} 1 & 0 \\ -\rho & I_d \end{pmatrix}. \end{aligned}$$

Note that L_ρ is nonsingular, so that $L_\rho L'_\rho$ is positive definite. Hence, by Assumption 2 part (ii) and by the fact that $L_\rho L'_\rho$ is a fixed, finite-dimensional positive definite matrix, there exists some constant $C_1 > 1$ such that

$$\min_{1 \leq (i,t) \leq m_n} \lambda_{\min}(\tilde{\Omega}_{(i,t)}) \geq \min_{1 \leq (i,t) \leq m_n} \lambda_{\min}(\Omega_{(i,t)}) \lambda_{\min}(L_\rho L'_\rho) \geq 1/C_1 > 0 \text{ a.s.n.} \quad (3)$$

Next, let

$$D_a = \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix}, D_{SD,(i,t)} = \begin{pmatrix} \sigma_{(i,t)} & 0 \\ 0 & \tilde{\omega}_{a,(i,t)} \end{pmatrix}, \text{ and } D_{\varrho,(i,t)} = \begin{pmatrix} 1 & \varrho_{a,(i,t)} \\ \varrho_{a,(i,t)} & 1 \end{pmatrix}$$

and note that

$$\begin{aligned} D_a' \tilde{\Omega}_{(i,t)} D_a &= \begin{pmatrix} 1 & 0 \\ 0 & a' \end{pmatrix} \begin{pmatrix} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] & E[\varepsilon_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z] \\ E[\underline{U}_{(i,t)} \varepsilon_{(i,t)} | \mathcal{F}_n^Z] & E[\underline{U}_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z] \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix} \\ &= \begin{pmatrix} \sigma_{(i,t)} & 0 \\ 0 & \tilde{\omega}_{a,(i,t)} \end{pmatrix} \begin{pmatrix} 1 & \varrho_{a,(i,t)} \\ \varrho_{a,(i,t)} & 1 \end{pmatrix} \begin{pmatrix} \sigma_{(i,t)} & 0 \\ 0 & \tilde{\omega}_{a,(i,t)} \end{pmatrix} = D_{SD,(i,t)} D_{\varrho,(i,t)} D_{SD,(i,t)}, \end{aligned}$$

Now, as can be seen from expression (3) above, an implication of Assumption 2(ii) is that

$$\min_{1 \leq (i,t) \leq m_n} \sigma_{(i,t)}^2 = e'_{1,d+1} \tilde{\Omega}_{(i,t)} e_{1,d+1} \geq 1/C_1 > 0 \text{ a.s.n.} \quad (4)$$

$$\min_{1 \leq (i,t) \leq m_n} \tilde{\omega}_{a,(i,t)}^2 = \underline{a}' \tilde{\Omega}_{(i,t)} \underline{a} \geq 1/C_1 > 0 \text{ a.s.n.} \quad (5)$$

where $e_{1,d+1} = (1 \ 0')'$ and $\underline{a} = (0 \ a')$, from which we deduce that $D_{SD,(i,t)}$ is invertible almost surely for each $(i,t) \in \{1, \dots, m_n\}$ and for all n sufficiently large. The invertibility of $D_{SD,(i,t)}$ then allows us to write $D_{\varrho,(i,t)} = D_{SD,(i,t)}^{-1} D_a' \tilde{\Omega}_{(i,t)} D_a D_{SD,(i,t)}^{-1}$. On the other hand, Assumption 2(i) implies that there exists some constant $C_2 > 1$ such that

$$\min_{1 \leq (i,t) \leq m_n} \lambda_{\min}(D_{SD,(i,t)}^{-1}) = \frac{1}{\max_{1 \leq (i,t) \leq m_n} \lambda_{\max}(D_{SD,(i,t)})} \geq \frac{1}{C_2} > 0 \text{ a.s.} \quad (6)$$

It follows from the fact that $\lambda_{\min}(D_a' D_a) = \lambda_{\min}(I_2) = 1$ and from making use of Assumptions 2(i) and (ii) and the lower bounds given in (3) and (6) that

$$\begin{aligned} \min_{1 \leq (i,t) \leq m_n} \lambda_{\min}(D_{\varrho,(i,t)}) &\geq \min_{1 \leq (i,t) \leq m_n} \lambda_{\min}(\tilde{\Omega}_{(i,t)}) \lambda_{\min}(D_a' D_a) \lambda_{\min}(D_{SD,(i,t)}^{-2}) \\ &\geq \frac{\min_{1 \leq (i,t) \leq m_n} \lambda_{\min}(\tilde{\Omega}_{(i,t)})}{\max_{1 \leq (i,t) \leq m_n} (\lambda_{\max}(D_{SD,(i,t)}))^2} \geq \frac{1}{C^3} > 0 \text{ a.s.n.}, \end{aligned}$$

where $C = \max\{C_1, C_2\}$. Moreover, by solving the characteristic equation of $D_{\varrho,(i,t)}$, we see that the smallest eigenvalue of $D_{\varrho,(i,t)}$ is given by $\lambda_{\min}(D_{\varrho,(i,t)}) = 1 - |\varrho_{a,(i,t)}|$, so that $\min_{1 \leq (i,t) \leq m_n} \lambda_{\min}(D_{\varrho,(i,t)}) = 1 - \max_{1 \leq (i,t) \leq m_n} |\varrho_{a,(i,t)}| \geq 1/C^3 > 0$ a.s.n., from which we further deduce that $\max_{1 \leq (i,t) \leq m_n} |\varrho_{a,(i,t)}| \leq 1 - (1/C^3) < 1$ a.s.n. Applying this upper bound along with the lower bounds given by (4) and (5) as well as the fact that $\lambda_{\min}(\Delta_{(i,t),(j,s)}^a) = 1 - |\varrho_{a,(i,t)}| |\varrho_{a,(j,s)}|$, as derived earlier, we have

$$\begin{aligned} E \left[\left(\varepsilon_{(j,s)} \underline{u}_{a,(i,t)} + \varepsilon_{(i,t)} \underline{u}_{a,(j,s)} \right)^2 | \mathcal{F}_n^Z \right] &= \delta'_{a,(i,t),(j,s)} \Delta_{(i,t),(j,s)}^a \delta_{a,(i,t),(j,s)} \\ &\geq \left[1 - |\varrho_{a,(i,t)}| |\varrho_{a,(j,s)}| \right] \left[\sigma_{(j,s)}^2 \tilde{\omega}_{a,(i,t)}^2 + \sigma_{(i,t)}^2 \tilde{\omega}_{a,(j,s)}^2 \right] \\ &\geq \left(\frac{2}{C^3} - \frac{1}{C^6} \right) \left[\frac{1}{C^2} + \frac{1}{C^2} \right] \geq \frac{2}{C^5} > 0 \text{ a.s.n.} \end{aligned}$$

Summing over $1 \leq (j,s) < (i,t) \leq m_n$, we obtain

$$\begin{aligned} &K_{2,n}^{-1} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 E \left[\left(\varepsilon_{(j,s)} \underline{u}_{a,(i,t)} + \varepsilon_{(i,t)} \underline{u}_{a,(j,s)} \right)^2 | \mathcal{F}_n^Z \right] \\ &\geq (2/C^5) K_{2,n}^{-1} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 = (1/C^5) K_{2,n}^{-1} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} A_{(i,t),(j,s)}^2, \text{ where the} \\ &\text{last equality follows from the symmetry of } A \text{ and by the fact that } A_{(i,t),(i,t)} = 0 \text{ for } (i,t) = 1, \dots, m_n. \end{aligned}$$

Furthermore, by straightforward calculation, we obtain

$$\begin{aligned} \frac{1}{K_{2,n}} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} A_{(i,t),(j,s)}^2 &= \frac{1}{K_{2,n}} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} \left[e'_{(i,t)} P^\perp e_{(j,s)} - e'_{(i,t)} M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} e_{(j,s)} \right]^2 \\ &= 1 + \frac{1}{K_{2,n}} \sum_{(i,t)=1}^{m_n} e'_{(i,t)} M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \geq 1 \end{aligned}$$

Putting everything together, we have

$$\begin{aligned} K_{2,n}^{-1} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 E \left[\left(\varepsilon_{(j,s)} \underline{u}_{a,(i,t)} + \varepsilon_{(i,t)} \underline{u}_{a,(j,s)} \right)^2 | \mathcal{F}_n^Z \right] &\geq \\ (1/C^5) K_{2,n}^{-1} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} A_{(i,t),(j,s)}^2 &\geq 1/C^5 \geq \underline{C} > 0, \text{ by choosing } \underline{C} \text{ such that } 0 < \underline{C} \leq \\ 1/C^5. \text{ Since the above argument holds for any } a \in \mathbb{R}^d \text{ such that } \|a\| = 1, \text{ it further follows that } &VC(\underline{U}' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z) \geq \underline{C} I_d > 0 \text{ a.s., as required.} \end{aligned}$$

To show part (c), note first that, given Assumption 2(i), there exists a positive constant C such that

$$\begin{aligned} &\max_{1 \leq (j,s) \leq m_n} \lambda_{\max} \left(E \left[\underline{U}_{(j,s)} \underline{U}'_{(j,s)} | \mathcal{F}_n^Z \right] \right) \\ &\leq \max_{1 \leq (j,s) \leq m_n} \text{tr} \left\{ E \left[\underline{U}_{(j,s)} \underline{U}'_{(j,s)} | \mathcal{F}_n^Z \right] \right\} \\ &\leq \max_{1 \leq (j,s) \leq m_n} \left\{ E \left[\|U_{(j,s)}\|_2^2 | \mathcal{F}_n^Z \right] + 2E \left[\left| U'_{(j,s)} \rho \varepsilon_{(j,s)} \right| | \mathcal{F}_n^Z \right] + \rho' \rho E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right] \right\} \\ &\leq \max_{1 \leq (j,s) \leq m_n} \left\{ E \left[\|U_{(j,s)}\|_2^2 | \mathcal{F}_n^Z \right] + 2 \|\rho\|_2 \sqrt{E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right]} \sqrt{E \left[\|U_{(j,s)}\|_2^2 | \mathcal{F}_n^Z \right]} \right. \\ &\quad \left. + \|\rho\|_2^2 E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right] \right\} \\ &\leq 2 \left\{ \left(\max_{1 \leq (j,s) \leq m_n} E \left[\|U_{(j,s)}\|_2^2 | \mathcal{F}_n^Z \right] \right) + \|\rho\|_2^2 \left(\max_{1 \leq (j,s) \leq m_n} E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right] \right) \right\} \\ &\leq C < \infty \text{ a.s.} \end{aligned} \tag{7}$$

where the third inequality above follows from applying the CS inequality while the fourth inequality stems in part from applying the inequality $|XY| \leq (1/2) X^2 + (1/2) Y^2$. Now, for any $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$; we obtain by applying the triangle and CS inequalities, expression (7), as well as

part (a) of Lemma S2-1 and Assumptions 2(i) and 8

$$\begin{aligned}
& a' V C \left(\underline{U}' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z \right) a \\
& \leq \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left(E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] a' E \left[\underline{U}_{(j,s)} \underline{U}'_{(j,s)} | \mathcal{F}_n^Z \right] a \right. \\
& \quad \left. + \sqrt{a' E \left[\underline{U}_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z \right] a} \sqrt{E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right]} \sqrt{a' E \left[\underline{U}_{(j,s)} \underline{U}'_{(j,s)} | \mathcal{F}_n^Z \right] a} \sqrt{E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right]} \right) \\
& \leq \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \lambda_{\max} \left(E \left[\underline{U}_{(j,s)} \underline{U}'_{(j,s)} | \mathcal{F}_n^Z \right] \right) \right. \\
& \quad \left. + \sqrt{\lambda_{\max} \left(E \left[\underline{U}_{(j,s)} \underline{U}'_{(j,s)} | \mathcal{F}_n^Z \right] \right)} \sqrt{E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right]} \sqrt{\lambda_{\max} \left(E \left[\underline{U}_{(j,s)} \underline{U}'_{(j,s)} | \mathcal{F}_n^Z \right] \right)} \sqrt{E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right]} \right\} \\
& = O_{a.s.}(1).
\end{aligned}$$

From this, we deduce that $\lambda_{\max} [V C (\underline{U}' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z)] = \max_{\|a\|=1} a' V C (\underline{U}' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z) a \leq \bar{C}$ a.s.n. Moreover, by applying the law of iterated expectations and part (i) of Theorem 16.1 of Billingsley (1995) that there exists a constant $\bar{C} > 0$ such that $a' V C (\underline{U}' A \varepsilon / \sqrt{K_{2,n}}) a = E_Z \{ E [(a' \underline{U}' A \varepsilon)^2 / K_{2,n}] \} \leq \bar{C}$, for all $a \in \mathbb{R}^d$ such that $\|a\| = 1$, from which we further deduce the unconditional version of this inequality, i.e., $\lambda_{\max} (V C [\underline{U}' A \varepsilon / \sqrt{K_{2,n}}]) = \max_{\|a\|=1} a' V C (\underline{U}' A \varepsilon / \sqrt{K_{2,n}}) a \leq \bar{C} < \infty$, where $\underline{U} = U - \varepsilon \rho'$ and $\underline{U}_{(i,t)} = U_{(i,t)} - \rho \varepsilon_{(i,t)}$.

Furthermore, since $\underline{U} = U - \varepsilon \rho'$, we see that, by setting $\rho = 0$ in the argument given above, we can also show that there exists a constant \bar{C} such that $\lambda_{\max} (V C [U' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z]) \leq \bar{C} < \infty$ a.s.n. and $\lambda_{\max} (V C [U' A \varepsilon / \sqrt{K_{2,n}}]) \leq \bar{C} < \infty$ for all n sufficiently large.

Finally, to show part (d), note first that, by straightforward calculations, we get $\Sigma_n = V C (\mathcal{Y}_n | \mathcal{F}_n^Z) = V C (\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z) + V C (D_\mu^{-1} \underline{U}' A \varepsilon | \mathcal{F}_n^Z) = \Sigma_{1,n} + \Sigma_{2,n}$. It follows by part (a) of this lemma that there exists a positive constant \underline{C} such that $\lambda_{\min} (\Sigma_n) \geq \lambda_{\min} [V C (\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z)] + \lambda_{\min} [V C (D_\mu^{-1} \underline{U}' A \varepsilon | \mathcal{F}_n^Z)] \geq \lambda_{\min} [V C (\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z)] \geq \underline{C} > 0$ a.s.n., so that Σ_n is positive definite a.s.n. Moreover, again by part (a) of this lemma, for any $a \in \mathbb{R}^d$ such that $\|a\| = 1$,

$$\begin{aligned}
a' \Sigma_n^{-1} a & \leq \frac{1}{\lambda_{\min} \{ V C (\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z) + V C (D_\mu^{-1} \underline{U}' A \varepsilon | \mathcal{F}_n^Z) \}} \\
& \leq \frac{1}{\lambda_{\min} [V C (\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} | \mathcal{F}_n^Z)]} \leq \frac{1}{\underline{C}} \leq \bar{C} < \infty \text{ a.s.n.}
\end{aligned}$$

where \bar{C} can be taken to be any finite, positive constant such that $\bar{C} \geq 1/\underline{C}$. \square

Lemma S2-4: Under Assumptions 1-6, $D_\mu^{-1} X' A \varepsilon = \Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} + D_\mu^{-1} U' A \varepsilon = O_p (\max \{ 1, \sqrt{K_{2,n}} / (\mu_n^{\min}) \})$
Proof of Lemma S2-4:

To proceed, since

$$\begin{aligned} A &= P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \\ &= P^{(Z,Q)} - P^{(Z_1,Q)} - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \\ &= M^{(Z_1,Q)} - \left[M^{(Z,Q)} + M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right], \end{aligned}$$

we can write

$$\begin{aligned} &D_\mu^{-1} X' A \varepsilon \\ &= D_\mu^{-1} \left(\frac{D_\kappa}{\sqrt{n}} \Theta' Z'_1 + \frac{D_\mu}{\sqrt{n}} \Upsilon' Z'_2 + \Xi' Q' + U' \right) \left(M^{(Z_1,Q)} - \left[M^{(Z,Q)} + M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right] \right) \varepsilon \\ &= \frac{\Upsilon' Z'_2 A \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' A \varepsilon \end{aligned} \tag{8}$$

Moreover, by straightforward calculations, it can be easily shown that

$$\begin{aligned} \frac{\Upsilon' Z'_2 A \varepsilon}{\sqrt{n}} &= \frac{\Upsilon' Z'_2 M^{(Z_1,Q)} \varepsilon}{\sqrt{n}} - \frac{\Upsilon' Z'_2 M^{(Z,Q)} \varepsilon}{\sqrt{n}} - \frac{\Upsilon' Z'_2 M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \varepsilon}{\sqrt{n}} \\ &= \frac{\Upsilon' Z'_2 M^{(Z_1,Q)} \varepsilon}{\sqrt{n}} \\ &= O_p(1) \end{aligned} \tag{9}$$

Now, let $b \in \mathbb{R}^d$ such that $\|b\|_2 = 1$, and, by straightforward calculations, we obtain

$$\begin{aligned} E \left([b' D_\mu^{-1} U' A \varepsilon]^2 | \mathcal{F}_n^Z \right) &= K_{2,n} b' D_\mu^{-1} E \left(\frac{U' A \varepsilon \varepsilon' A U}{K_{2,n}} | \mathcal{F}_n^Z \right) D_\mu^{-1} b \\ &= K_{2,n} b' D_\mu^{-1} V C \left(U' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z \right) D_\mu^{-1} b \\ &\leq \lambda_{\max} \left[V C \left(U' A \varepsilon / \sqrt{K_{2,n}} | \mathcal{F}_n^Z \right) \right] \frac{K_{2,n}}{(\mu_n^{\min})^2} \\ &\leq \overline{C} \frac{K_{2,n}}{(\mu_n^{\min})^2} \text{ a.s.} \end{aligned}$$

for some positive constant $\overline{C} < \infty$. Hence, for all n sufficiently large

$$E \left(\frac{(\mu_n^{\min})^2}{K_{2,n}} [b' D_\mu^{-1} U' A \varepsilon]^2 \right) = E_Z \left\{ \frac{(\mu_n^{\min})^2}{K_{2,n}} E \left([b' D_\mu^{-1} U' A \varepsilon]^2 | \mathcal{F}_n^Z \right) \right\} \leq \overline{C}.$$

It follows from Markov's inequality, for any $\epsilon > 0$, we can set $C_\epsilon = \sqrt{\overline{C}/\epsilon}$ so that for all n sufficiently

large

$$\begin{aligned}
\Pr \left(\left| \frac{\mu_n^{\min}}{\sqrt{K_{2,n}}} b' D_\mu^{-1} U' A \varepsilon \right| \geq C_\epsilon \right) &= \Pr \left(\left| \frac{\mu_n^{\min}}{\sqrt{K_{2,n}}} b' D_\mu^{-1} U' A \varepsilon \right|^2 \geq C_\epsilon^2 \right) \\
&\leq \frac{1}{C_\epsilon^2} E \left(\frac{(\mu_n^{\min})^2}{K_{2,n}} [b' D_\mu^{-1} U' A \varepsilon]^2 \right) \\
&\leq \frac{\overline{C}}{\overline{C}/\epsilon} = \epsilon
\end{aligned}$$

which shows that

$$b' D_\mu^{-1} U' A \varepsilon = O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} \right).$$

Since the above result holds for all $b \in \mathbb{R}^d$ such that $\|b\|_2 = 1$, we further deduce that

$$D_\mu^{-1} U' A \varepsilon = O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} \right). \quad (10)$$

Expressions (8)-(10) together imply that

$$\begin{aligned}
\frac{D_\mu^{-1} X' A \varepsilon}{\mu_n^{\min}} &= \frac{\Upsilon' Z'_2 A \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' A \varepsilon \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' A \varepsilon \\
&= O_p \left(\max \left\{ 1, \frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} \right\} \right). \quad \square
\end{aligned}$$

Lemma S2-5: Under Assumptions 1-6, $D_\mu^{-1} X' M^{(Z_1, Q)} \varepsilon = O_p(n/\mu_n^{\min})$.

Proof of Lemma S2-5:

To proceed, let $b \in \mathbb{R}^d$ such that $\|b\|_2 = 1$ and let $u_b = U D_\mu^{-1} b$ and $u_{b,(i,t)} = U'_{(i,t)} D_\mu^{-1} b$. Now, write

$$\begin{aligned}
&b' D_\mu^{-1} X' M^{(Z_1, Q)} \varepsilon \\
&= b' D_\mu^{-1} \left(\frac{D_\kappa}{\sqrt{n}} \Theta' Z'_1 + \frac{D_\mu}{\sqrt{n}} \Upsilon' Z'_2 + \Xi' Q' + U' \right) M^{(Z_1, Q)} \varepsilon \\
&= \frac{b' \Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + u'_b M^{(Z_1, Q)} \varepsilon.
\end{aligned}$$

By straightforward calculations and by applying Assumptions 1, 2(i), 3, 4, 5(i), and 5(iii) as well as the CS and Markov's inequalities, we can show that $b' \Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} = O_p(1)$. From the CS inequality, we also obtain $E [|u'_b M^{(Z_1, Q)} \varepsilon| | \mathcal{F}_n^Z]$

$\leq \sqrt{E [u'_b M^{(Z_1, Q)} u_b | \mathcal{F}_n^Z]} \sqrt{E [\varepsilon' M^{(Z_1, Q)} \varepsilon | \mathcal{F}_n^Z]}$. Next, note that, by applying Assumptions 2(i) and

6(ii), we have $E[\varepsilon' M^{(Z_1, Q)} \varepsilon | \mathcal{F}_n^Z] \leq E[\varepsilon' \varepsilon | \mathcal{F}_n^Z] \leq n \bar{T} \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) = O_{a.s.}(n)$. Similarly, by applying Assumptions 2(i), 3(ii), and 6(ii); we obtain $E[u_b' M^{(Z_1, Q)} u_b | \mathcal{F}_n^Z]$
 $= E[b' D_\mu^{-1} U' M^{(Z_1, Q)} U D_\mu^{-1} b | \mathcal{F}_n^Z] \leq \bar{T} \left(\max_{1 \leq (i,t) \leq m_n} E[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z] \right) \left[n / (\mu_n^{\min})^2 \right]$
 $= O_{a.s.} \left(n (\mu_n^{\min})^{-2} \right)$. It follows from these results that $E[|u_b' M^{(Z_1, Q)} \varepsilon| | \mathcal{F}_n^Z] = O_{a.s.} \left(n (\mu_n^{\min})^{-1} \right)$. Hence, by the law of iterated expectations and Theorem 16.1 of Billingsley (1995), there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large $E[((\mu_n^{\min})/n) |u_b' M^{(Z_1, Q)} \varepsilon|]$
 $= E_Z((\mu_n^{\min})/n) E[|u_b' M^{(Z_1, Q)} \varepsilon| | \mathcal{F}_n^Z]) \leq \bar{C}$. By applying Markov's inequality, we further obtain

$$b' D_\mu^{-1} U' M^{(Z_1, Q)} \varepsilon = O_p \left(n (\mu_n^{\min})^{-1} \right). \quad (11)$$

Putting everything together, we have

$$\begin{aligned} b' D_\mu^{-1} X' M^{(Z_1, Q)} \varepsilon &= \frac{b' \Upsilon' Z_2' M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + b' D_\mu^{-1} U' M^{(Z_1, Q)} \varepsilon \\ &= O_p(1) + O_p \left(\frac{n}{(\mu_n^{\min})} \right) = O_p \left(\frac{n}{(\mu_n^{\min})} \right) \end{aligned}$$

Since the above argument holds for all $b \in \mathbb{R}^d$ such that $\|b\|_2 = 1$, we further deduce that $D_\mu^{-1} X' M^{(Z_1, Q)} \varepsilon = O_p \left(n (\mu_n^{\min})^{-1} \right)$. \square

Lemma S2-6: If Assumptions 2 and 8 are satisfied; then, for $1 \leq p \leq 8$ and for all n , there exists a positive constant C such that $\max_{1 \leq (i,t) \leq m_n} E[\|\underline{U}_{(i,t)}\|_2^p | \mathcal{F}_n^Z] \leq C < \infty$ a.s., where $\underline{U}_{(i,t)} = U_{(i,t)} - \rho \varepsilon_{(i,t)}$.

Proof of Lemma S2-6: Note that, for $1 \leq p \leq 8$ and for any $(i,t) \in \{1, \dots, m_n\}$, there exists a positive constant C such that

$$\begin{aligned} E[\|\underline{U}_{(i,t)}\|_2^p | \mathcal{F}_n^Z] &= E[\left(\|U_{(i,t)} - \rho \varepsilon_{(i,t)}\|_2\right)^p | \mathcal{F}_n^Z] \\ &\leq 2^{p-1} \left\{ \left(E[\|U_{(i,t)}\|_2^8 | \mathcal{F}_n^Z] \right)^{p/8} + \|\rho\|_2^p \left(E[|\varepsilon_{(i,t)}|^8 | \mathcal{F}_n^Z] \right)^{p/8} \right\} \\ &\leq C < \infty \text{ a.s.}, \end{aligned}$$

where the first inequality follows by applying the triangle inequality, Loève's c_r inequality, and Liapunov's inequality in sequence and where the second inequality follows from applying Assumption 2(i) and from the fact that $\rho \in \mathcal{S}_\rho$, some compact subset of \mathbb{R}^d as stated in Assumption 8. Since the upper bound above holds for all $(i,t) \in \{1, \dots, m_n\}$, for all $\rho \in \mathcal{S}_\rho$, and for all n , it further follows that $\max_{1 \leq (i,t) \leq m_n} E[\|\underline{U}_{(i,t)}\|_2^p | \mathcal{F}_n^Z] \leq C < \infty$ a.s., as required. \square

Lemma S2-7: Under Assumptions 1-6, the following results hold: (a) $\hat{\ell}_{L,n} = o_p \left([\mu_n^{\min}]^2/n \right)$; (b) $\hat{\ell}_{F,n} = o_p \left([\mu_n^{\min}]^2/n \right)$.

Proof of Lemma S2-7: To proceed, first define

$$\bar{D}_n = \begin{pmatrix} \mu_n^{\min} & 0 \\ 0 & D_\mu \end{pmatrix} \text{ and } L_\delta = \begin{pmatrix} 1 & 0 \\ \delta_0 & I_d \end{pmatrix},$$

and note that

$$L_\delta^{-1} = \begin{pmatrix} 1 & 0 \\ -\delta_0 & I_d \end{pmatrix}$$

Now, for any $\beta \in \mathbb{R}^{d+1}$ such that $\|\beta\|_2 = 1$ and for $\bar{X} = [y \ X]$, we can write $\beta' \bar{X}' A \bar{X} \beta = \beta' L'_\delta \bar{D}_n \left(\bar{D}_n^{-1} L'_\delta^{-1} \bar{X}' A \bar{X} L_\delta^{-1} \bar{D}_n^{-1} \right) \bar{D}_n L_\delta \beta$. Moreover, by direct multiplication,

$$\bar{D}_n^{-1} L'_\delta^{-1} \bar{X}' A \bar{X} L_\delta^{-1} \bar{D}_n^{-1} = \begin{pmatrix} (y - X\delta_0)' A (y - X\delta_0) / (\mu_n^{\min})^2 & (y - X\delta_0)' A X D_\mu^{-1} / (\mu_n^{\min}) \\ D_\mu^{-1} X' A (y - X\delta_0) / (\mu_n^{\min}) & D_\mu^{-1} X' A X D_\mu^{-1} \end{pmatrix},$$

Now, by straightforward but tedious calculations and by applying Assumptions 1, 2(i), 3(ii)-(iii), 4, 5, and 6(i), as well as Lemmas S2-3(c) and S2-1(a); we can show that, under the rate condition $\sqrt{K_2} / (\mu_n^{\min})^2 \rightarrow 0$,

$$\bar{D}_n^{-1} L'_\delta^{-1} \bar{X}' A \bar{X} L_\delta^{-1} \bar{D}_n^{-1} = \begin{pmatrix} 0 & 0 \\ 0 & \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n \end{pmatrix} + o_p(1)^1,$$

where, in light of Assumption 3(iii), $\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n = O_p(1)$ and $\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n$ is positive definite for all n large sufficiently large. It follows that $\bar{D}_n^{-1} L'_\delta^{-1} \bar{X}' A \bar{X} L_\delta^{-1} \bar{D}_n^{-1}$ is positive semidefinite w.p.a.1, so that

$\beta' \bar{X}' A \bar{X} \beta = \beta' L'_\delta \bar{D}_n \left(\bar{D}_n^{-1} L'_\delta^{-1} \bar{X}' A \bar{X} L_\delta^{-1} \bar{D}_n^{-1} \right) \bar{D}_n L_\delta \beta \geq 0$ w.p.a.1 for all $\beta \in \mathbb{R}^{d+1}$ such that $\|\beta\|_2 = 1$. Moreover, by straightforward calculations, we can show that

$$\frac{\beta' \bar{X}' M^{(Z_1, Q)} \bar{X} \beta}{n} = \frac{\beta' L'_{\delta, 2} D_\mu \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon D_\mu L_{\delta, 2} \beta}{n^2} + \beta' L'_\delta E \left[\frac{V' M^Q V}{n} \right] L_\delta \beta + o_p(1),$$

where $V = [\varepsilon \ U]$ and $L_{\delta, 2} = [\delta_0 \ I_d]$ and where $E[V' M^Q V / n]$ is positive definite for all n sufficiently large in light of Assumptions 2(ii) and 6(i). Since $L_\delta \beta \neq 0$ for all $\beta \in \mathbb{R}^{d+1}$ such that $\|\beta\|_2 = 1$, it follows that $\beta' \bar{X}' M^{(Z_1, Q)} \bar{X} \beta / n > 0$ w.p.a.1 for all $\beta \in \mathbb{R}^{d+1}$ such that $\|\beta\|_2 = 1$. Hence, with probability approaching one as $n \rightarrow \infty$,

$$R(\beta) = \frac{\beta' \bar{X}' A \bar{X} \beta}{\beta' \bar{X}' M^{(Z_1, Q)} \bar{X} \beta}$$

is a continuous function of β for all values of β such that $\|\beta\|_2 = 1$. The Weierstrass extreme value theorem then implies that there exists some $\tilde{\beta}$ such that $\tilde{\beta} = \arg \min_{\|\beta\|_2=1} R(\beta)$ w.p.a.1. Next, note that $\hat{\ell}_{L,n}$ is the smallest root of the determinantal equation

$\det \left\{ \bar{X}' A \bar{X} - \ell \bar{X}' M^{(Z_1, Q)} \bar{X} \right\} = 0$; and, thus, $\hat{\ell}_{L,n}$ has the representation

$$\hat{\ell}_{L,n} = R(\tilde{\beta}) = \frac{\tilde{\beta}' \bar{X}' A \bar{X} \tilde{\beta}}{\tilde{\beta}' \bar{X}' M^{(Z_1, Q)} \bar{X} \tilde{\beta}} = \min_{\|\beta\|_2=1} \left(\frac{\beta' \bar{X}' A \bar{X} \beta}{\beta' \bar{X}' M^{(Z_1, Q)} \bar{X} \beta} \right).$$

Now, let $\delta_* = (1 \ -\delta'_0)' / \left\| (1 \ -\delta'_0)' \right\|_2$; and we have, with probability approaching one as $n \rightarrow \infty$,

$$\begin{aligned} 0 &\leq \hat{\ell}_{L,n} = \min_{\|\beta\|_2=1} \left(\frac{\beta' \bar{X}' A \bar{X} \beta}{\beta' \bar{X}' M^{(Z_1, Q)} \bar{X} \beta} \right) \\ &\leq \frac{\delta'_* \bar{X}' A \bar{X} \delta_*}{\delta'_* \bar{X}' M^{(Z_1, Q)} \bar{X} \delta_*} \\ &= \frac{(\mu_n^{\min})^2}{n} \left\{ \frac{(y - X\delta_0)' A (y - X\delta_0) / \sqrt{K_{2,n}}}{(y - X\delta_0)' M^{(Z_1, Q)} (y - X\delta_0) / n} \right\} \frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \\ &= O\left(\frac{[\mu_n^{\min}]^2}{n}\right) O_p(1) O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2}\right) = o_p\left(\frac{[\mu_n^{\min}]^2}{n}\right), \end{aligned}$$

given the rate condition $\sqrt{K_2} / (\mu_n^{\min})^2 \rightarrow 0$. This shows part (a).

For part (b), we use the result in part (a) above and the fact that $m_n/n \sim 1$ by Assumption 5(i) to obtain

$$\hat{\ell}_{F,n} = \frac{\hat{\ell}_{L,n} - (1 - \hat{\ell}_{L,n}) C/m_n}{1 - (1 - \hat{\ell}_{L,n}) C/m_n} = \left[\hat{\ell}_{L,n} + O_p\left(\frac{1}{n}\right) \right] \left[1 + O_p\left(\frac{1}{n}\right) \right] = o_p\left(\frac{[\mu_n^{\min}]^2}{n}\right). \quad \square \quad (12)$$

We introduce some notations before proceeding to the next lemma. Let

$(u_{(1,1),n}, \varepsilon_{(1,1),n}), \dots, (u_{(1,T_1),n}, \varepsilon_{(1,T_1),n}), (u_{(2,1),n}, \varepsilon_{(2,1),n}), \dots, (u_{(2,T_2),n}, \varepsilon_{(2,T_2),n}), \dots, (u_{(n,1),n}, \varepsilon_{(n,1),n}), \dots, (u_{(n,T_n),n}, \varepsilon_{(n,T_n),n})$ denote a triangular array of (bivariate) random vectors. Also, let $\phi_{(i,t),n} = \phi_{(i,t),n}(Z)$, $(i, t) = 1, \dots, m_n$, denote a triangular array of measurable functions. In applications of the lemma given below, we will take $\phi_{(i,t),n}$ to be either a conditional variance or a conditional covariance given $\mathcal{F}_n^Z = \sigma(Z)$, the σ -algebra generated by Z . In addition, we let $\sigma_{(i,t),n}^2 = E[\varepsilon_{(i,t),n}^2 | \mathcal{F}_n^Z]$, $\bar{\omega}_{(i,t),n}^2 = E[u_{(i,t),n}^2 | \mathcal{F}_n^Z]$, and $\bar{\psi}_{(i,t),n} = E[\varepsilon_{(i,t),n} u_{(i,t),n} | \mathcal{F}_n^Z]$. In order to simplify notation, we suppress the dependence of $\phi_{(i,t),n}$, $\sigma_{(i,t),n}^2$, $\bar{\omega}_{(i,t),n}^2$, and $\bar{\psi}_{(i,t),n}$ on Z .

Lemma S2-8: Let A be as defined above. Assume that i) $(u_{(1,1),n}, \varepsilon_{(1,1),n}), \dots, (u_{(1,T_1),n}, \varepsilon_{(1,T_1),n}), (u_{(2,1),n}, \varepsilon_{(2,1),n}), \dots, (u_{(2,T_2),n}, \varepsilon_{(2,T_2),n}), \dots, (u_{(n,1),n}, \varepsilon_{(n,1),n}), \dots, (u_{(n,T_n),n}, \varepsilon_{(n,T_n),n})$ are independent conditional on $\mathcal{F}_n^Z = \sigma(Z)$; ii) there exists a constant C such that, almost surely for all n sufficiently large, $\max_{1 \leq (i,t) \leq m_n} E(u_{(i,t),n}^4 | \mathcal{F}_n^Z) \leq C$, $\max_{1 \leq (i,t) \leq m_n} E(\varepsilon_{(i,t),n}^4 | \mathcal{F}_n^Z) \leq C$, and $\max_{1 \leq (i,t) \leq m_n} |\phi_{(i,t),n}| \leq C$. In addition, define $\bar{\psi}_{(j,s),n} = E[u_{(j,s),n} \varepsilon_{(j,s),n} | \mathcal{F}_n^Z]$ for $(j, s) = 1, \dots, m_n$. Then, under Assumptions 5 and 6, the following statements are true:

- (a) $K_{2,n}^{-1} \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \left\{ u_{(j,s),n} \varepsilon_{(j,s),n} - \bar{\psi}_{(j,s),n} \right\} \xrightarrow{p} 0$;

- (b) $K_{2,n}^{-1} \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \} \xrightarrow{P} 0$;
- (c) $K_{2,n}^{-1} \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \varepsilon_{(j,s),n} \varepsilon_{(k,v),n} \xrightarrow{P} 0$;
- (d) $K_{2,n}^{-1} \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} u_{(j,s),n} u_{(k,v),n} \xrightarrow{P} 0$.

Proof of Lemma S2-8: To show part (a), note that

$$\begin{aligned}
& E \left[\left(\frac{1}{K_{2,n}} \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n}^2 \{ u_{(j,s),n} \varepsilon_{(j,s),n} - \bar{\psi}_{(j,s),n} \} \right)^2 \mid \mathcal{F}_n^Z \right] \\
&= \frac{1}{K_{2,n}^2} \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^4 \phi_{(i,t),n}^2 \left\{ E \left(\varepsilon_{(j,s),n}^2 u_{(j,s),n}^2 \mid \mathcal{F}_n^Z \right) - \bar{\psi}_{(j,s),n}^2 \right\} \\
&\quad + \frac{2}{K_{2,n}^2} \sum_{1 \leq (j,s) < (k,v) < (i,t) \leq m_n} A_{(k,v),(j,s)}^2 A_{(i,t),(j,s)}^2 \phi_{(k,v),n} \phi_{(i,t),n} \left\{ E \left(u_{(j,s),n}^2 \varepsilon_{(j,s),n}^2 \mid \mathcal{F}_n^Z \right) - \bar{\psi}_{(j,s),n}^2 \right\} \\
&\leq \frac{1}{K_{2,n}^2} \sum_{1 \leq (j,s) < (i,t) \leq m_n} \left\{ A_{(i,t),(j,s)}^4 \phi_{(i,t),n}^2 \right. \\
&\quad \times \left[\sqrt{E \left(u_{(j,s),n}^4 \mid \mathcal{F}_n^Z \right) E \left(\varepsilon_{(j,s),n}^4 \mid \mathcal{F}_n^Z \right)} + E \left(u_{(j,s),n}^2 \mid \mathcal{F}_n^Z \right) E \left(\varepsilon_{(j,s),n}^2 \mid \mathcal{F}_n^Z \right) \right] \} \\
&\quad + \frac{2}{K_{2,n}^2} \sum_{1 \leq (j,s) < (k,v) < (i,t) \leq m_n} A_{(k,v),(j,s)}^2 A_{(i,t),(j,s)}^2 |\phi_{(k,v),n}| |\phi_{(i,t),n}| \\
&\quad \times \left\{ \sqrt{E \left(u_{(j,s),n}^4 \mid \mathcal{F}_n^Z \right) E \left(\varepsilon_{(j,s),n}^4 \mid \mathcal{F}_n^Z \right)} + E \left(u_{(j,s),n}^2 \mid \mathcal{F}_n^Z \right) E \left(\varepsilon_{(j,s),n}^2 \mid \mathcal{F}_n^Z \right) \right\} \\
&\leq C \left\{ \frac{1}{K_{2,n}^2} \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^4 + \frac{2}{K_{2,n}^2} \sum_{1 \leq (j,s) < (k,v) < (i,t) \leq m_n} A_{(k,v),(j,s)}^2 A_{(i,t),(j,s)}^2 \right\} \\
&\leq C \left\{ \frac{1}{K_{2,n}^2} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 + \frac{2}{K_{2,n}^2} \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}}^{m_n} A_{(k,v),(j,s)}^2 A_{(i,t),(j,s)}^2 \right\} \\
&= o_{a.s.}(1)
\end{aligned}$$

where the first inequality is the result of applying T and a conditional version of CS, the second inequality follows by hypothesis, and the convergence to zero almost surely follows from parts (b) and (c) of Lemma S2-1 and the symmetry of A . It follows by the conditional version of the Markov's inequality that for any $\epsilon > 0$

$$\Pr \left(\left| \frac{1}{K_{2,n}} \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \{ \varepsilon_{(j,s),n} u_{(j,s),n} - \bar{\psi}_{(j,s),n} \} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \rightarrow 0 \text{ a.s.}$$

Note further that

$$\sup_n E \left[\left| \Pr \left(\left| \sum_{1 \leq (j,s) < (i,t) \leq m_n} \frac{A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \{ \varepsilon_{(j,s),n} u_{(j,s),n} - \bar{\psi}_{(j,s),n} \}}{K_{2,n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \right|^2 \right] < \infty$$

Hence, by a version of the dominated convergence theorem, as given by Theorem 25.12 of Billingsley (1986), we have, as $n \rightarrow \infty$,

$$\begin{aligned} & \Pr \left(\left| \frac{1}{K_{2,n}} \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \left\{ \varepsilon_{(j,s),n} u_{(j,s),n} - \bar{\psi}_{(j,s),n} \right\} \right| \geq \epsilon \right) \\ &= E \left[\Pr \left(\left| \frac{1}{K_{2,n}} \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \left\{ \varepsilon_{(j,s),n} u_{(j,s),n} - \bar{\psi}_{(j,s),n} \right\} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \right] \rightarrow 0, \end{aligned}$$

as required for part (a).

To show part (b), first let L be the lower triangular matrix such that $L_{(i,t),(j,s)} = A_{(i,t),(j,s)} \mathbb{I}\{(i,t) > (j,s)\}$, and define $D_{\bar{\psi}} = \text{diag}(\bar{\psi}_{(1,1),n}, \dots, \bar{\psi}_{(n,T_n),n}) = \text{diag}(\bar{\psi}_1, \dots, \bar{\psi}_{m_n})$, $D_\phi = \text{diag}(\phi_{(1,1)}, \dots, \phi_{(n,T_n)}) = \text{diag}(\phi_1, \dots, \phi_{m_n})$, $u = (u_{(1,1)}, \dots, u_{(n,T_n)})' = (u_1, \dots, u_{m_n})'$, and $\varepsilon = (\varepsilon_{(1,1)}, \dots, \varepsilon_{(n,T_n)})' = (\varepsilon_1, \dots, \varepsilon_{m_n})'$. It then follows by direct multiplication that

$$\begin{aligned} & \varepsilon' L' D_\phi L u - \text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \\ &= \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \left\{ u_{(j,s),n} \varepsilon_{(j,s),n} - \bar{\psi}_{(j,s),n} \right\} \\ &\quad + \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \left\{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \right\} \end{aligned}$$

so that by making use of Loève's c_r inequality, we have that

$$\begin{aligned} & E \left[\left(\sum_{\substack{1 \leq (k,v) < (j,s) \\ < (i,t) \leq m_n}} \frac{A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \left\{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \right\}}{K_{2,n}^2} \right)^2 \mid \mathcal{F}_n^Z \right] \\ &\leq 2 \frac{1}{K_{2,n}^2} E \left[\left(u' L' D_\phi L \varepsilon - \text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right)^2 \mid \mathcal{F}_n^Z \right] \\ &\quad + 2 \frac{1}{K_{2,n}^2} E \left[\left(\sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \left\{ u_{(j,s),n} \varepsilon_{(j,s),n} - \bar{\psi}_{(j,s),n} \right\} \right)^2 \mid \mathcal{F}_n^Z \right] \quad (13) \end{aligned}$$

From the proof of part (a), we already have

$$\frac{1}{K_{2,n}^2} E \left[\left(\sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)}^2 \phi_{(i,t),n} \left\{ u_{(j,s),n} \varepsilon_{(j,s),n} - \bar{\psi}_{(j,s),n} \right\} \right)^2 \mid \mathcal{F}_n^Z \right] = o_{a.s.}(1).$$

To show that

$$\frac{1}{K_{2,n}^2} E \left[\left(u' L' D_\phi L \varepsilon - \text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right)^2 \mid \mathcal{F}_n^Z \right] = o_{a.s.}(1),$$

note first that

$$\begin{aligned}
& \frac{1}{K_{2,n}^2} E \left[\left(u' L' D_\phi L \varepsilon - \text{tr} \left\{ L' D_\phi L D D_{\bar{\psi}} \right\} \right)^2 \mid \mathcal{F}_n^Z \right] \\
= & \frac{1}{K_{2,n}^2} E \left[(u' L' D_\phi L \varepsilon)^2 \mid \mathcal{F}_n^Z \right] - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 \\
= & \frac{1}{K_{2,n}^2} E \left[u' L' D_\phi L \varepsilon \otimes u' L' D_\phi L \varepsilon \mid \mathcal{F}_n^Z \right] - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 \\
= & \frac{1}{K_{2,n}^2} E \left[\text{tr} \left\{ (u' \otimes u') (L' D_\phi L \otimes L' D_\phi L) (\varepsilon \otimes \varepsilon) \right\} \mid \mathcal{F}_n^Z \right] \\
& - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 \\
= & \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) E \left[\varepsilon u' \otimes \varepsilon u' \mid \mathcal{F}_n^Z \right] \right\} - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2. \quad (14)
\end{aligned}$$

Next, by straightforward calculations, we obtain

$$\begin{aligned}
& E [\varepsilon u' \otimes \varepsilon u' | \mathcal{F}_n^Z] \\
&= \left(\begin{array}{ccc} \sigma_{(1,1),n}^2 \bar{\omega}_{(1,1),n}^2 e_{(1,1)} e'_{(1,1)} & \cdots & \sigma_{(1,1),n}^2 \bar{\omega}_{(n,T_n),n}^2 e_{(1,1)} e'_{(n,T_n)} \\ \vdots & & \vdots \\ \sigma_{(1,T_1),n}^2 \bar{\omega}_{(1,1),n}^2 e_{(1,T_1)} e'_{(1,1)} & \cdots & \sigma_{(1,T_1),n}^2 \bar{\omega}_{(n,T_n),n}^2 e_{(1,T_1)} e'_{(n,T_n)} \\ \vdots & & \vdots \\ \sigma_{(n,1),n}^2 \bar{\omega}_{(1,1),n}^2 e_{(n,1)} e'_{(1,1)} & \cdots & \sigma_{(n,1),n}^2 \bar{\omega}_{(n,T_n),n}^2 e_{(n,1)} e'_{(n,T_n)} \\ \vdots & & \vdots \\ \sigma_{(n,T_n),n}^2 \bar{\omega}_{(1,1),n}^2 e_{(n,T_n)} e'_{(1,1)} & \cdots & \sigma_{(n,T_n),n}^2 \bar{\omega}_{(n,T_n),n}^2 e_{(n,T_n)} e'_{(n,T_n)} \end{array} \right) \\
&+ \left(\begin{array}{ccc} \bar{\psi}_{(1,1),n}^2 e_{(1,1)} e'_{(1,1)} & \cdots & \bar{\psi}_{(1,1),n} \bar{\psi}_{(n,T_n),n} e_{(n,T_n)} e'_{(1,1)} \\ \vdots & & \vdots \\ \bar{\psi}_{(1,T_1),n} \bar{\psi}_{(1,1),n} e_{(1,1)} e'_{(1,T_1)} & \cdots & \bar{\psi}_{(1,T_1),n} \bar{\psi}_{(n,T_n),n} e_{(n,T_n)} e'_{(1,T_1)} \\ \vdots & & \vdots \\ \bar{\psi}_{(n,1),n} \bar{\psi}_{(1,1),n} e_{(1,1)} e'_{(n,1)} & \cdots & \bar{\psi}_{(n,1),n} \bar{\psi}_{(n,T_n),n} e_{(n,T_n)} e'_{(n,1)} \\ \vdots & & \vdots \\ \bar{\psi}_{(n,T_n),n} \bar{\psi}_{(1,1),n} e_{(1,1)} e'_{(n,T_n)} & \cdots & \bar{\psi}_{(n,T_n),n} \bar{\psi}_{(n,T_n),n} e_{(n,T_n)} e'_{(n,T_n)} \end{array} \right) \\
&+ \left(\begin{array}{ccccc} \bar{\kappa}_{(1,1),n} e_{(1,1)} e'_{(1,1)} & 0 & \cdots & \cdots & 0 \\ m_n \times m_n & & & & m_n \times m_n \\ 0 & \ddots & \ddots & & \vdots \\ m_n \times m_n & & & & \vdots \\ \vdots & \ddots & \bar{\kappa}_{(1,T_1),n} e_{(1,T_1)} e'_{(1,T_1)} & \ddots & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ m_n \times m_n & 0 & \cdots & 0 & m_n \times m_n \\ \bar{\psi}_{(1,1),n} \otimes D_{\bar{\psi}} & m_n \times m_n & \cdots & \cdots & 0 \\ m_n \times m_n & \ddots & \ddots & & \vdots \\ \vdots & & \ddots & \ddots & \vdots \\ m_n \times m_n & 0 & \cdots & 0 & m_n \times m_n \\ \bar{\psi}_{(1,T_1),n} \otimes D_{\bar{\psi}} & m_n \times m_n & \cdots & \cdots & 0 \\ \vdots & \ddots & \bar{\psi}_{(1,T_1),n} \otimes D_{\bar{\psi}} & \ddots & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ m_n \times m_n & 0 & \cdots & 0 & m_n \times m_n \\ \bar{\psi}_{(n,T_n),n} \otimes D_{\bar{\psi}} & m_n \times m_n & \cdots & \cdots & 0 \end{array} \right) \\
&= (D_\sigma \otimes I_{m_n}) \text{vec}(I_{m_n}) \text{vec}(I_{m_n}) (D_{\bar{\omega}} \otimes I_{m_n}) \\
&\quad + (D_\psi \otimes I_{m_n}) \underline{K}_{m_n m_n} (D_\psi \otimes I_{m_n}) + \underline{E}' D_{\bar{\kappa}} \underline{E} + \left(D_{\bar{\psi}} \otimes D_{\bar{\psi}} \right), \tag{15}
\end{aligned}$$

where $\underline{K}_{m_n m_n}$ is an $m_n^2 \times m_n^2$ commutation matrix such that for any $m_n \times m_n$ matrix A , $\underline{K}_{m_n m_n} \text{vec}(A) = \text{vec}(A')$. Also, here, $D_{\bar{\psi}} = \text{diag}(\bar{\psi}_1, \dots, \bar{\psi}_{m_n})$, $D_\sigma = \text{diag}(\sigma_1^2, \dots, \sigma_{m_n}^2)$, $D_{\bar{\omega}} = \text{diag}(\bar{\omega}_1^2, \dots, \bar{\omega}_{m_n}^2)$, $D_{\bar{\kappa}} = \text{diag}(\bar{\kappa}_1, \dots, \bar{\kappa}_{m_n})$ with $\bar{\kappa}_{(i,t),n} = E[\varepsilon_{(i,t),n}^2 u_{(i,t),n}^2 | \mathcal{F}_n^Z] - \sigma_{(i,t),n}^2 \bar{\omega}_{(i,t),n}^2 - 2\bar{\psi}_{(i,t),n}^2$ for $(i, t) =$

$1, \dots, m_n$. Also, let

$\underline{E} = (e_{1,m_n} \otimes e_{1,m_n}, \dots, e_{m_n,m_n} \otimes e_{m_n,m_n})'$; and e_{i,m_n} is the i^{th} column of an $m_n \times m_n$ identity matrix . It follows from (14) and (15) that

$$\begin{aligned}
& \frac{1}{K_{2,n}^2} E \left[\left(u' L' D_\phi L \varepsilon - \text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right)^2 \mid \mathcal{F}_n^Z \right] \\
&= \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) E [\varepsilon u' \otimes \varepsilon u' | \mathcal{F}_n^Z] \right\} - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 \\
&= \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) (D_\sigma \otimes I_{m_n}) \text{vec}(I_{m_n}) \text{vec}(I_{m_n})' (D_{\bar{\omega}} \otimes I_{m_n}) \right\} \\
&\quad + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) (D_\psi \otimes I_{m_n}) \underline{K}_{m_n m_n} \left(D_{\bar{\psi}} \otimes I_{m_n} \right) \right\} \\
&\quad + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) \underline{E}' D_{\bar{\omega}} \underline{E} \right\} + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) \left(D_{\bar{\psi}} \otimes D_{\bar{\psi}} \right) \right\} \\
&\quad - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 \\
&= \frac{1}{K_{2,n}^2} \text{vec}(I_{m_n})' (D_{\bar{\omega}} L' D_\phi L D_\sigma \otimes L' D_\phi L) \text{vec}(I_{m_n}) \\
&\quad + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (D_\psi L' D_\phi L D_{\bar{\psi}} \otimes L' D_\phi L) \underline{K}_{m_n m_n} \right\} + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) \underline{E}' D_{\bar{\omega}} \underline{E} \right\} \\
&\quad + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L D_{\bar{\psi}} \otimes L' D_\phi L D_{\bar{\psi}}) \right\} - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 \\
&= \frac{1}{K_{2,n}^2} \text{tr} \left\{ L' D_\phi L D_{\bar{\omega}} L' D_\phi L D_\sigma \right\} + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (D_\psi L' D_\phi L D_{\bar{\psi}} \otimes L' D_\phi L) \underline{K}_{m_n m_n} \right\} \\
&\quad + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) \underline{E}' D_{\bar{\omega}} \underline{E} \right\} + \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 - \frac{1}{K_{2,n}^2} \left[\text{tr} \left\{ L' D_\phi L D_{\bar{\psi}} \right\} \right]^2 \\
&= \frac{1}{K_{2,n}^2} \text{tr} \left\{ L' D_\phi L D_{\bar{\omega}} L' D_\phi L D_\sigma \right\} + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (D_{\bar{\psi}} L' D_\phi L D_{\bar{\psi}} \otimes L' D_\phi L) \underline{K}_{m_n m_n} \right\} \\
&\quad + \frac{1}{K_{2,n}^2} \text{tr} \left\{ (L' D_\phi L \otimes L' D_\phi L) \underline{E}' D_{\bar{\omega}} \underline{E} \right\} \tag{16}
\end{aligned}$$

Focusing first on the first term of (16), we get

$$\begin{aligned}
& \frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' D_\phi L D_{\bar{\omega}} L' D_\phi L D_\sigma \} \\
& \leq \sqrt{\frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' D_\phi L D_{\bar{\omega}}^2 L' D_\phi L \}} \sqrt{\frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' D_\phi L D_\sigma^2 L' D_\phi L \}} \\
& \leq \sqrt{\max_{1 \leq (i,t) \leq m_n} \bar{\omega}_{(i,t),n}^4} \sqrt{\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^4} \frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' D_\phi L L' D_\phi L \} \\
& \leq \sqrt{\max_{1 \leq (i,t) \leq m_n} \bar{\omega}_{(i,t),n}^4} \sqrt{\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^4} \sqrt{\frac{1}{K_{2,n}^2} \operatorname{tr} \{ D_\phi L L' D_\phi^2 L L' D_\phi \}} \sqrt{\frac{1}{K_{2,n}^2} \operatorname{tr} \{ L L' L L' \}} \\
& \leq \sqrt{\max_{1 \leq (i,t) \leq m_n} \bar{\omega}_{(i,t),n}^4} \sqrt{\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^4} \sqrt{\max_{1 \leq (i,t) \leq m_n} \phi_{(i,t),n}^2} \\
& \quad \times \sqrt{\frac{1}{K_{2,n}^2} \operatorname{tr} \{ L L' D_\phi^2 L L' \}} \sqrt{\frac{1}{K_{2,n}^2} \operatorname{tr} \{ L L' L L' \}} \\
& \leq \sqrt{\max_{1 \leq (i,t) \leq m_n} \bar{\omega}_{(i,t),n}^4} \sqrt{\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^4} \left(\max_{1 \leq (i,t) \leq m_n} \phi_{(i,t),n}^2 \right) \frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' L L' L \} \\
& \leq \sqrt{\max_{1 \leq (i,t) \leq m_n} E(u_{(i,t),n}^4 | \mathcal{F}_n^Z)} \sqrt{\max_{1 \leq (i,t) \leq m_n} E(\varepsilon_{(i,t),n}^4 | \mathcal{F}_n^Z)} \left(\max_{1 \leq (i,t) \leq m_n} \phi_{(i,t),n}^2 \right) \frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' L L' L \} \\
& \leq C \frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' L L' L \} = \frac{C}{K_{2,n}^2} \|L L'\|_F^2 \quad a.s., \tag{17}
\end{aligned}$$

where the first and third inequalities follow from CS and where the sixth inequality follows from the conditional version of the Jensen's inequality and the last inequality follows in light of the assumptions of this lemma. In addition, let G be an $m_n \times m_n$ matrix and $D = \operatorname{diag}(d_1, \dots, d_{m_n})$ such that $d_{(i,t)} \geq 0$ for all $(i,t) \in \{1, \dots, m_n\}$, and note that the second and fourth inequalities in (17) above follows from the inequality

$$\operatorname{tr} \{ G' D G \} \leq \left\{ \max_{(i,t)} (d_{(i,t)}) \right\} \operatorname{tr} (G' G). \tag{18}$$

Turning our attention now to the second term of (16), we see that, using the identity $\operatorname{tr} \{(A \otimes B) \underline{K}_{m_n m_n}\} = \operatorname{tr} \{AB\}$ for $m_n \times m_n$ matrices A and B ,

$$\begin{aligned}
& \frac{1}{K_{2,n}^2} \operatorname{tr} \left\{ \left(D_{\bar{\psi}} L' D_\phi L D_{\bar{\psi}} \otimes L' D_\phi L \right) \underline{K}_{m_n m_n} \right\} \\
& = \frac{1}{K_{2,n}^2} \operatorname{tr} \left\{ D_{\bar{\psi}} L' D_\phi L D_{\bar{\psi}} L' D_\phi L \right\} = \frac{1}{K_{2,n}^2} \operatorname{tr} \left\{ L' D_\phi L D_{\bar{\psi}} L' D_\phi L D_{\bar{\psi}} \right\}.
\end{aligned}$$

It follows by calculations similar to that used to obtain (17), we obtain

$$\begin{aligned}
& \frac{1}{K_{2,n}^2} \operatorname{tr} \left\{ \left(D_{\bar{\psi}} L' D_{\phi} L D_{\bar{\psi}} \otimes L' D_{\phi} L \right) \underline{K}_{m_n m_n} \right\} \\
& \leq \left(\max_{1 \leq (i,t) \leq m_n} \bar{\psi}_{(i,t),n}^2 \right) \left(\max_{1 \leq (i,t) \leq m_n} \phi_{(i,t),n}^2 \right) \frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' L L' L \} \\
& \leq C \frac{1}{K_{2,n}^2} \operatorname{tr} \{ L' L L' L \} = \frac{C}{K_{2,n}^2} \| L L' \|_F^2 \quad a.s.
\end{aligned} \tag{19}$$

Finally, to analyze the third term of (16), we note that

$$\begin{aligned}
& \frac{1}{K_{2,n}^2} \left| \operatorname{tr} \{ (L' D_{\phi} L \otimes L' D_{\phi} L) \underline{E}' D_{\bar{\pi}} \underline{E} \} \right| \\
& \leq \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} |\bar{\pi}_{(i,t),n}| \operatorname{tr} \left\{ e'_{(i,t)} L' D_{\phi} L e_{(i,t)} L' D_{\phi} L e_{(i,t)} e'_{(i,t)} \right\} \\
& = \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} |\bar{\pi}_{(i,t),n}| \left(e'_{(i,t)} L' D_{\phi} L e_{(i,t)} \right)^2 \\
& \leq \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} |\bar{\pi}_{(i,t),n}| \left(e'_{(i,t)} L' D_{\phi}^2 L e_{(i,t)} \right) \left(e'_{(i,t)} L' L e_{(i,t)} \right) \\
& \leq \left(\max_{1 \leq (i,t) \leq m_n} \phi_{(i,t),n}^2 \right) \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} |\bar{\pi}_{(i,t),n}| \left(e'_{(i,t)} L' L e_{(i,t)} \right)^2 \\
& \leq \left\{ \sqrt{E \left[\varepsilon_{(i,t),n}^4 | \mathcal{F}_n^Z \right]} \sqrt{E \left[u_{(i,t),n}^4 | \mathcal{F}_n^Z \right]} + \left(\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^2 \right) \left(\max_{1 \leq (i,t) \leq m_n} \bar{\omega}_{(i,t),n}^2 \right) \right. \\
& \quad \left. + 2 \left(\max_{1 \leq (i,t) \leq m_n} \bar{\psi}_{(i,t),n}^2 \right) \right\} \left(\max_{1 \leq (i,t) \leq m_n} \phi_{(i,t),n}^2 \right) \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(e'_{(i,t)} L' L e_{(i,t)} \right)^2 \\
& \leq C \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(e'_{(i,t)} L' L e_{(i,t)} \right)^2 \leq C \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(e'_{(i,t)} A' A e_{(i,t)} \right)^2 \\
& = C \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(e'_{(i,t)} \left[P^{\perp} - M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} \right]^2 e_{(i,t)} \right)^2 \\
& = C \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(e'_{(i,t)} P^{\perp} e_{(i,t)} + e'_{(i,t)} M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \right)^2
\end{aligned}$$

where the first inequality above follows from T, the second inequality follows from CS, the third inequality makes use of (18), the fourth inequality uses CS and T, and the fifth inequality stems from our assumption about the (almost sure) boundedness of the conditional moments. Applying

the inequality $\left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r$ for $r \geq 1$, we further obtain

$$\begin{aligned}
& \frac{1}{K_{2,n}^2} \left| \text{tr} \left\{ (L'D_\phi L \otimes L'D_\phi L) \underline{E}' D_{\bar{\vartheta}} \underline{E} \right\} \right| \\
& \leq 2C \left[\frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(P_{(i,t),(i,t)}^\perp \right)^2 + \frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(e'_{(i,t)} M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} D_{\hat{\vartheta}} M^{(Z,Q)} e_{(i,t)} \right)^2 \right] \\
& \leq 2C \left[\frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} \left(P_{(i,t),(i,t)}^\perp \right)^2 + \left(\max_{1 \leq (i,t) \leq m_n} |\hat{\vartheta}_{(i,t)}|^2 \right) \frac{1}{K_{2,n}^2} \text{tr} \left\{ D_{\hat{\vartheta}} M^{(Z,Q)} D_{\hat{\vartheta}} \right\} \right] \\
& \leq 2C \left[1 + \left(\frac{1}{C_*} \right)^2 \right] \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right) \left[\frac{1}{K_{2,n}^2} \sum_{(i,t)=1}^{m_n} P_{(i,t),(i,t)}^\perp + \frac{1}{K_{2,n}^2} \text{tr} \left\{ D_{\hat{\vartheta}}^2 \right\} \right] \\
& = O_{a.s.} \left(\frac{1}{n} \right)
\end{aligned} \tag{20}$$

where we have used Assumption 5(iv) and part (a) of Lemma OA-1 in arriving at the last line above.

In light of (16), it follows from (17), (19), (20), and Lemma OA-11 (given in section 3 below) that

$$\frac{1}{K_{2,n}^2} E \left[\left(u'L'D_\phi L \varepsilon - \text{tr} \left\{ L'D_\phi L D_{\bar{\psi}} \right\} \right)^2 \mid \mathcal{F}_n^Z \right] \leq 2C (1/K_{2,n}^2) \|LL'\|_F^2 + C (1/K_{2,n}) \leq C/K_{2,n} \text{ a.s.}$$

It follows from (13) that

$$\begin{aligned}
& E \left[\left(\sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} \frac{A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \}}{K_{2,n}^2} \right)^2 \mid \mathcal{F}_n^Z \right] \\
& = o_{a.s.}(1)
\end{aligned}$$

Moreover, by the conditional version of the Markov's inequality, we deduce for any $\epsilon > 0$

$$\begin{aligned}
& \Pr \left(\left| \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} \frac{A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \}}{K_{2,n}^2} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \\
& \rightarrow 0 \text{ a.s.}
\end{aligned}$$

Since

$$\begin{aligned}
& \sup_n E \left[\left| \Pr \left(\sum_{\substack{1 \leq (k,v) < (j,s) \\ < (i,t) \leq m_n}} \frac{A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \}}{K_{2,n}^2} \geq \epsilon \mid \mathcal{F}_n^Z \right) \right|^2 \right] \\
& < \infty,
\end{aligned}$$

it further follows by a version of the dominated convergence theorem, as given by Theorem 25.12 of Billingsley (1986), that as $n \rightarrow \infty$

$$\begin{aligned} & \Pr \left(\left| \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} \frac{A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \}}{K_{2,n}} \right| \geq \epsilon \right) \\ &= E \left[\Pr \left(\left| \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} \frac{A_{(i,t),(j,s)} A_{(i,t),(k,v)} \phi_{(i,t),n} \{ u_{(j,s),n} \varepsilon_{(k,v),n} + \varepsilon_{(j,s),n} u_{(k,v),n} \}}{K_{2,n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \right] \\ &\rightarrow 0, \end{aligned}$$

as required for part (b).

It is easily seen that parts (c) and (d) can be proved in essentially the same way as part (b); hence, to avoid redundancy, we do not provide detailed arguments here. \square

Lemma S2-9: Let

$$\hat{\Delta}(\delta_0) = -\frac{(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)}{2} \frac{\partial}{\partial \delta} \left\{ \frac{(y - X\delta)' A (y - X\delta)}{(y - X\delta)' M^{(Z_1,Q)} (y - X\delta)} \right\} \Big|_{\delta=\delta_0}.$$

If Assumptions 1-6 and 8 are satisfied; then, $D_\mu^{-1} \hat{\Delta}(\delta_0) = \Upsilon' Z'_2 M^{(Z_1,Q)} \varepsilon / \sqrt{n} + D_\mu^{-1} \underline{U}' A \varepsilon + o_p(1)$, where $\underline{U} = U - \varepsilon \rho'$ and where $\rho = \lim_{n \rightarrow \infty} E[U' M^Q \varepsilon] / E[\varepsilon' M^Q \varepsilon]$.

Proof of Lemma S2-9: To proceed, note first that we can write

$$\begin{aligned} & \hat{\Delta}(\delta_0) \\ &= -\frac{(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)}{2} \frac{\partial}{\partial \delta} \left\{ \frac{(y - X\delta)' A (y - X\delta)}{(y - X\delta)' M^{(Z_1,Q)} (y - X\delta)} \right\} \Big|_{\delta=\delta_0} \\ &= -\frac{(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)}{2} \\ &\quad \times \left\{ \frac{-2X'A(y - X\delta_0)}{(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)} + \frac{(y - X\delta_0)' A (y - X\delta_0) 2X'M^{(Z_1,Q)} (y - X\delta_0)}{[(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)]^2} \right\} \\ &= X'A(y - X\delta_0) - \frac{(y - X\delta_0)' A (y - X\delta_0)}{(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)} X'M^{(Z_1,Q)} (y - X\delta_0) \\ &= X'A(y - X\delta_0) - \hat{\ell}(\delta_0) X'M^{(Z_1,Q)} (y - X\delta_0) \end{aligned}$$

where $A = P^\perp - M^{(Z,Q)} D_{\hat{\theta}} M^{(Z,Q)}$. It follows from making use of the fact that $\rho_n = E[U' M^{(Z_1,Q)} \varepsilon] / E[\varepsilon' M^{(Z_1,Q)} \varepsilon]$ and from applying Lemma OA-5 and part (c) of Lemma

OA-4 that

$$\begin{aligned}
& D_\mu^{-1} \widehat{\Delta}(\delta_0) \\
&= D_\mu^{-1} \left[X' A(y - X\delta_0) - \widehat{\ell}(\delta_0) X' M^{(Z_1, Q)}(y - X\delta_0) \right] \\
&= D_\mu^{-1} X' A(y - X\delta_0) - \widehat{\ell}(\delta_0) D_\mu^{-1} X' M^{(Z_1, Q)}(y - X\delta_0) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' A \varepsilon - \frac{\varepsilon' A \varepsilon}{\varepsilon' M^Q \varepsilon} \left[1 + O_p \left(\frac{K_{1,n}}{n} \right) \right] \left[\frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' M^{(Z_1, Q)} \varepsilon \right] \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} (U - \varepsilon \rho')' A \varepsilon - \varepsilon' A \varepsilon D_\mu^{-1} \left[\frac{U' M^{(Z_1, Q)} \varepsilon}{\varepsilon' M^Q \varepsilon} - \rho \right] \left[1 + O_p \left(\frac{K_{1,n}}{n} \right) \right] \\
&\quad - \frac{\sqrt{K_{2,n}} \varepsilon' A \varepsilon / \sqrt{K_{2,n}}}{\varepsilon' M^Q \varepsilon / n} \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + O_p \left(\frac{K_{1,n} \sqrt{K_{2,n}}}{n^2} \right) + O_p \left(\frac{K_{1,n} \sqrt{K_{2,n}}}{(\mu_n^{\min}) n} \right) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} (U - \varepsilon \rho')' A \varepsilon - \varepsilon' A \varepsilon D_\mu^{-1} \left[\frac{U' M^{(Z_1, Q)} \varepsilon}{\varepsilon' M^Q \varepsilon} - \rho \right] \left[1 + O_p \left(\frac{K_{1,n}}{n} \right) \right] \\
&\quad + O_p \left(\frac{\sqrt{K_{2,n}}}{n} \right) + O_p \left(\frac{K_{1,n} \sqrt{K_{2,n}}}{(\mu_n^{\min}) n} \right) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} (U - \varepsilon \rho')' A \varepsilon + O_p \left(\frac{1}{\mu_n^{\min}} \max \left\{ \sqrt{\frac{K_{2,n}}{n}}, \frac{K_{1,n} \sqrt{K_{2,n}}}{n} \right\} \right) \\
&\quad + O_p \left(\frac{\sqrt{K_{2,n}}}{n} \right) + O_p \left(\frac{K_{1,n} \sqrt{K_{2,n}}}{(\mu_n^{\min}) n} \right) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} \underline{U}' A \varepsilon + O_p \left(\max \left\{ \frac{1}{\mu_n^{\min}} \sqrt{\frac{K_{2,n}}{n}}, \frac{K_{1,n} \sqrt{K_{2,n}}}{(\mu_n^{\min}) n} \right\} \right) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} \underline{U}' A \varepsilon + o_p(1). \quad \square
\end{aligned}$$

Lemma S2-10: Let Assumptions 1-6 be satisfied, and let $\bar{\delta}_n$ be any estimator such that, as $n \rightarrow \infty$, $D_\mu(\bar{\delta}_n - \delta_0) / \mu_n^{\min} = o_p(1)$. Then, $-D_\mu^{-1} (\partial \widehat{\Delta}(\bar{\delta}_n) / \partial \delta') D_\mu^{-1} = H_n + o_p(1)$, where $H_n = \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n$ and where

$$\begin{aligned}
\widehat{\Delta}(\delta) &= - \left[(y - X\delta)' M^{(Z_1, Q)} (y - X\delta) / 2 \right] \left[\partial \widehat{Q}_{FELIM}(\delta) / \partial \delta \right] \\
&= X' A(y - X\delta) - \widehat{\ell}(\delta) X' M^{(Z_1, Q)}(y - X\delta),
\end{aligned}$$

with

$$\widehat{\ell}(\delta) = (y - X\delta)' A(y - X\delta) / \left[(y - X\delta)' M^{(Z_1, Q)} (y - X\delta) \right].$$

In addition, we also have

$$D_\mu^{-1} X' \left[A - \hat{\ell}(\bar{\delta}_n) M^{(Z_1, Q)} \right] X D_\mu^{-1} = H_n + o_p(1). \quad (21)$$

Proof of Lemma S2-10:

Taking derivative of $-\hat{\Delta}(\delta)$ with respect to δ , we obtain

$$\begin{aligned} & -\frac{\partial \hat{\Delta}(\delta)}{\partial \delta'} \\ = & X' AX - \frac{(y - X\delta)' A (y - X\delta)}{(y - X\delta)' M^{(Z_1, Q)} (y - X\delta)} X' M^{(Z_1, Q)} X - \frac{2X' M^{(Z_1, Q)} (y - X\delta) (y - X\delta)' AX}{(y - X\delta)' M^{(Z_1, Q)} (y - X\delta)} \\ & + 2X' M^{(Z_1, Q)} (y - X\delta) (y - X\delta)' M^{(Z_1, Q)} X \frac{(y - X\delta)' A (y - X\delta)}{[(y - X\delta)' M^{(Z_1, Q)} (y - X\delta)]^2} \\ = & X' AX - \hat{\ell}(\delta) X' M^{(Z_1, Q)} X \\ & - \frac{2X' M^{(Z_1, Q)} (y - X\delta)}{(y - X\delta)' M^{(Z_1, Q)} (y - X\delta)} \left\{ (y - X\delta)' AX - \hat{\ell}(\delta) (y - X\delta)' M^{(Z_1, Q)} X \right\} \\ = & X' AX - \hat{\ell}(\delta) X' M^{(Z_1, Q)} X - \frac{2X' M^{(Z_1, Q)} (y - X\delta)}{(y - X\delta)' M^{(Z_1, Q)} (y - X\delta)} \hat{\Delta}(\delta)' \end{aligned}$$

so that evaluating $-\partial \hat{\Delta}(\delta) / \partial \delta'$ at $\delta = \bar{\delta}_n$, we have

$$\begin{aligned} -\frac{\partial \hat{\Delta}(\bar{\delta}_n)}{\partial \delta'} &= X' AX - \hat{\ell}(\bar{\delta}_n) X' M^{(Z_1, Q)} X - \frac{2X' M^{(Z_1, Q)} (y - X\bar{\delta}_n)}{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} (y - X\bar{\delta}_n)} \hat{\Delta}(\bar{\delta}_n)' \\ &= X' \left[A - \hat{\ell}(\bar{\delta}_n) M^{(Z_1, Q)} \right] X - \frac{2X' M^{(Z_1, Q)} (y - X\bar{\delta}_n)}{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} (y - X\bar{\delta}_n)} \hat{\Delta}(\bar{\delta}_n)' \end{aligned}$$

Pre-multiplying and post-multiplying the above expression by D_μ^{-1} , we then obtain

$$\begin{aligned} & -D_\mu^{-1} \left(\frac{\partial \hat{\Delta}(\bar{\delta}_n)}{\partial \delta'} \right) D_\mu^{-1} \\ = & D_\mu^{-1} X' \left[A - \hat{\ell}(\bar{\delta}_n) M^{(Z_1, Q)} \right] X D_\mu^{-1} - \frac{2D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\bar{\delta}_n)}{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} (y - X\bar{\delta}_n)} \hat{\Delta}(\bar{\delta}_n)' D_\mu^{-1} \end{aligned}$$

Now, part (b) of Lemma S2-2 gives

$$D_\mu^{-1} X' A X D_\mu^{-1} = \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon}{n} + o_p(1) = H_n + o_p(1), \quad (22)$$

and, by part (a) of Lemma S2-2 as well as parts (c) and (d) of Lemma OA-6 given section 3 of this

online appendix below, we have

$$\begin{aligned}
& \widehat{\ell}(\bar{\delta}_n) D_{\mu}^{-1} X' M^{(Z_1, Q)} X D_{\mu}^{-1} \\
&= \frac{(y - X\bar{\delta}_n)' A(y - X\bar{\delta}_n) / (\mu_n^{\min})^2}{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} (y - X\bar{\delta}_n) / n} \left((\mu_n^{\min})^2 \frac{D_{\mu}^{-1} X' M^{(Z_1, Q)} X D_{\mu}^{-1}}{n} \right) \\
&= o_p(1) O_p(1) = o_p(1).
\end{aligned} \tag{23}$$

It follows from expressions (22) and (23) that

$$\begin{aligned}
D_{\mu}^{-1} X' \left[A - \widehat{\ell}(\bar{\delta}_n) M^{(Z_1, Q)} \right] X D_{\mu}^{-1} &= D_{\mu}^{-1} X' A X D_{\mu}^{-1} - \widehat{\ell}(\bar{\delta}_n) D_{\mu}^{-1} X' M^{(Z_1, Q)} X D_{\mu}^{-1} \\
&= H_n + o_p(1).
\end{aligned}$$

where $H_n = \Upsilon' Z_2' M^{(Z_1, Q)} Z_2 \Upsilon / n$. Moreover, making use of parts (a)-(d) of Lemma OA-6, we further obtain

$$\begin{aligned}
& \frac{2D_{\mu}^{-1} X' M^{(Z_1, Q)} (y - X\bar{\delta}_n)}{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} (y - X\bar{\delta}_n)} \widehat{\Delta}(\bar{\delta}_n)' D_{\mu}^{-1} \\
&= \frac{2D_{\mu}^{-1} X' M^{(Z_1, Q)} (y - X\bar{\delta}_n) / n}{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} (y - X\bar{\delta}_n) / n} \\
&\quad \times \left[(y - X\bar{\delta}_n)' A X D_{\mu}^{-1} - \frac{(y - X\bar{\delta}_n)' A (y - X\bar{\delta}_n)}{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} (y - X\bar{\delta}_n) / n} \frac{(y - X\bar{\delta}_n)' M^{(Z_1, Q)} X D_{\mu}^{-1}}{n} \right] \\
&= O_p\left(\frac{1}{\mu_n^{\min}}\right) \left[o_p(\mu_n^{\min}) - o_p([\mu_n^{\min}]^2) O_p\left(\frac{1}{\mu_n^{\min}}\right) \right] = o_p(1)
\end{aligned}$$

so that

$$-D_{\mu}^{-1} \left(\frac{\partial \widehat{\Delta}(\bar{\delta}_n)}{\partial \delta'} \right) D_{\mu}^{-1} = \frac{\Upsilon' Z_2' M^{(Z_1, Q)} Z_2 \Upsilon}{n} + o_p(1) = H_n + o_p(1). \quad \square$$

Lemma S2-11: Let $\widehat{\ell}_L = Q(\tilde{\beta}) = \min_{\beta \in \overline{B}} Q(\beta)$, where $Q(\beta)$ is as defined in Assumption 9. Then, $\widehat{\ell}_L$ is also the smallest root of the determinantal equation $\det[\overline{X}' A \overline{X} - \ell \overline{X}' M^{(Z_1, Q)} \overline{X}] = 0$, where $\overline{X} = [y, X]$. Assume in addition that condition (13) in Assumption 9 is satisfied; then, $\widehat{\ell}_L$ has the representation

$$\widehat{\ell}_L = \frac{(y - X\widehat{\delta}_L)' A (y - X\widehat{\delta}_L)}{(y - X\widehat{\delta}_L)' M^{(Z_1, Q)} (y - X\widehat{\delta}_L)}, \tag{24}$$

where $\widehat{\delta}_L$ denotes the FELIM estimator. Moreover, $\overline{X}' A (y - X\widehat{\delta}_L) - \widehat{\ell}_L \overline{X}' M^{(Z_1, Q)} (y - X\widehat{\delta}_L) = 0$. In particular, this implies that $\widehat{\Delta}(\widehat{\delta}_L) = 0$, where

$\widehat{\Delta}(\delta) = -[(y - X\delta)' M^{(Z_1, Q)} (y - X\delta) / 2] (\partial \widehat{Q}_{FELIM}(\delta) / \partial \delta)$, so that $\widehat{\delta}_L$ satisfies the set of (normalized) first-order conditions for minimizing the variance ratio objective function $\widehat{Q}_{FELIM}(\delta) =$

$$(y - X\delta)' A (y - X\delta) / [(y - X\delta)' M^{(Z_1, Q)} (y - X\delta)].$$

Proof of Lemma S2-11:

Note that the first-order condition for minimizing the objective function $Q(\beta)$ can be written as $\partial Q(\tilde{\beta})/\partial \beta = 2\bar{X}'A\bar{X}\tilde{\beta}/(\tilde{\beta}'\bar{X}'M^{(Z_1, Q)}\bar{X}\tilde{\beta}) - \tilde{\beta}'\bar{X}'A\bar{X}\tilde{\beta}(2\bar{X}'M^{(Z_1, Q)}\bar{X}\tilde{\beta})/[\tilde{\beta}'\bar{X}'M^{(Z_1, Q)}\bar{X}\tilde{\beta}]^2 = 0$. Pre-multiplying this first order condition by the factor $\frac{1}{2}\tilde{\beta}'\bar{X}'M^{(Z_1, Q)}\bar{X}\tilde{\beta}$, we then obtain

$$0 = [\bar{X}'A\bar{X} - \hat{\ell}_L\bar{X}'M^{(Z_1, Q)}\bar{X}] \tilde{\beta} \quad (25)$$

where we have set $\hat{\ell}_L = Q(\tilde{\beta}) = \tilde{\beta}'\bar{X}'A\bar{X}\tilde{\beta}/\tilde{\beta}'\bar{X}'M^{(Z_1, Q)}\bar{X}\tilde{\beta}$. It is clear that in order for there to be a nontrivial solution, i.e., $\tilde{\beta} \neq 0$ such that equation (25) is true, $\hat{\ell}_L$ must be a root of the determinantal equation $\det[\bar{X}'A\bar{X} - \ell\bar{X}'M^{(Z_1, Q)}\bar{X}] = 0$. Moreover, since our goal is to minimize the value of the objective function $Q(\beta)$, this implies that we should choose $\hat{\ell}_L$ to be the smallest root of this determinantal equation. Now, define $\tilde{\delta} = -\tilde{\beta}_2/\tilde{\beta}_1$, and rewrite the first-order conditions given by expression (25) as $0 = \bar{X}'A\bar{X}\tilde{\beta} - \hat{\ell}_L\bar{X}'M^{(Z_1, Q)}\bar{X}\tilde{\beta} = \tilde{\beta}_1 \left\{ \bar{X}'A(y - X\tilde{\delta}) - \hat{\ell}_L\bar{X}'M^{(Z_1, Q)}(y - X\tilde{\delta}) \right\}$, so that, given the condition that $|\tilde{\beta}_1| \geq \underline{C} > 0$ a.s.n. for some constant \underline{C} (as stated in Assumption 9), we must have

$$\bar{X}'A(y - X\tilde{\delta}) - \hat{\ell}_L\bar{X}'M^{(Z_1, Q)}(y - X\tilde{\delta}) = 0 \quad (26)$$

Since $\bar{X} = [y, X]$, we can partition (26) into two sets of equations

$$0 = y'A(y - X\tilde{\delta}) - \hat{\ell}_Ly'M^{(Z_1, Q)}(y - X\tilde{\delta}), \quad (27)$$

$$0 = X'A(y - X\tilde{\delta}) - \hat{\ell}_LX'M^{(Z_1, Q)}(y - X\tilde{\delta}). \quad (28)$$

Solving (28) for $\tilde{\delta}$, we obtain $\tilde{\delta} = \left(X' \left[A - \hat{\ell}_L M^{(Z_1, Q)} \right] X \right)^{-1} X' \left[A - \hat{\ell}_L M^{(Z_1, Q)} \right] y = \hat{\delta}_L$, so that the FELIM estimator $\hat{\delta}_L$ is a solution to the second set of equations given by (28). In addition , note that, under condition (13) in Assumption 9, we have

$$\hat{\ell}_L = \frac{\tilde{\beta}'\bar{X}'A\bar{X}\tilde{\beta}}{\tilde{\beta}'\bar{X}'M^{(Z_1, Q)}\bar{X}\tilde{\beta}} = \frac{\tilde{\beta}_1(y - X\tilde{\delta})' A(y - X\tilde{\delta}) \tilde{\beta}_1}{\tilde{\beta}_1(y - X\tilde{\delta}) M^{(Z_1, Q)}(y - X\tilde{\delta}) \tilde{\beta}_1} = \frac{(y - X\hat{\delta}_L)' A(y - X\hat{\delta}_L)}{(y - X\hat{\delta}_L)' M^{(Z_1, Q)}(y - X\hat{\delta}_L)}.$$

which shows (24). Furthermore, note that $\widehat{\delta}_L$ also satisfies equation (27) since

$$\begin{aligned}
& y' A \left(y - X \widehat{\delta}_L \right) - \widehat{\ell}_L y' M^{(Z_1, Q)} \left(y - X \widehat{\delta}_L \right) \\
&= \left(y - X \widehat{\delta}_L \right)' \left[A - \widehat{\ell}_L M^{(Z_1, Q)} \right] \left(y - X \widehat{\delta}_L \right) + \widehat{\delta}'_L X' \left[A - \widehat{\ell}_L M^{(Z_1, Q)} \right] \left(y - X \widehat{\delta}_L \right) \\
&= \left(y - X \widehat{\delta}_L \right)' A \left(y - X \widehat{\delta}_L \right) - \frac{\left(y - X \widehat{\delta}_L \right)' A \left(y - X \widehat{\delta}_L \right) \left(y - X \widehat{\delta}_L \right)' M^{(Z_1, Q)} \left(y - X \widehat{\delta}_L \right)}{\left(y - X \widehat{\delta}_L \right) M^{(Z_1, Q)} \left(y - X \widehat{\delta}_L \right)} \\
&\quad + \widehat{\delta}'_L X' \left[A - \widehat{\ell}_L M^{(Z_1, Q)} \right] y \\
&\quad - \widehat{\delta}'_L X' \left[A - \widehat{\ell}_L M^{(Z_1, Q)} \right] X \left(X' \left[A - \widehat{\ell}_L M^{(Z_1, Q)} \right] X \right)^{-1} X' \left[A - \widehat{\ell}_L M^{(Z_1, Q)} \right] y \\
&= 0
\end{aligned}$$

from which we further deduce that $\widehat{\delta}_L$ is a solution of the complete set of first-order conditions given by (26). Finally, since $\widehat{\Delta}(\widehat{\delta}_L) = X' A \left(y - X \widehat{\delta}_L \right) - \widehat{\ell}_L X' M^{(Z_1, Q)} \left(y - X \widehat{\delta}_L \right)$, the fact that $\widehat{\delta}_L$ is a solution of (28) directly imply that $\widehat{\delta}_L$ satisfies the set of (normalized) first-order conditions for minimizing the variance ratio objective function. \square

Lemma S2-12: If Assumptions 1-6 are satisfied; then,

$$D_\mu^{-1} X' \left[A - \widehat{\ell}_{F,n} M^{(Z_1, Q)} \right] X D_\mu^{-1} = H_n + o_p(1), \text{ where } H_n = \Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon / n,$$

$\widehat{\ell}_{F,n} = \left[\widehat{\ell}_{L,n} - \left(1 - \widehat{\ell}_{L,n} \right) (C/m_n) \right] / \left[1 - \left(1 - \widehat{\ell}_{L,n} \right) (C/m_n) \right]$, and $\widehat{\ell}_{L,n}$ is smallest root of the determinantal equation $\det \left\{ \overline{X}' A \overline{X} - \ell \overline{X}' M^{(Z_1, Q)} \overline{X} \right\} = 0$, with $\overline{X} = [y \ X]$.

Proof of Lemma S2-12: The result follows directly from applying part (b) of Lemma S2-7, parts (a) and (b) of Lemma S2-2, and the Slutsky theorem. \square

Lemma S2-13: If Assumptions 1-6 and 8-9 are satisfied; then,

$$D_\mu^{-1} X' \left[A - \widehat{\ell}_{F,n} M^{(Z_1, Q)} \right] (y - X \delta_0) = \mathcal{Y}_n [1 + o_p(1)], \text{ where } \mathcal{Y}_n = \Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon / \sqrt{n} + D_\mu^{-1} \underline{U}' A \varepsilon \text{ with } \underline{U} = U - \varepsilon \rho' \text{ and } \rho = \lim_{n \rightarrow \infty} E[U' M^Q \varepsilon] / E[\varepsilon' M^Q \varepsilon].$$

Proof of Lemma S2-13:

Note that, by Lemma S2-11 above, $\widehat{\ell}_{L,n}$ has the representation

$$\widehat{\ell}_{L,n} = \frac{\left(y - X \widehat{\delta}_{L,n} \right)' A \left(y - X \widehat{\delta}_{L,n} \right)}{\left(y - X \widehat{\delta}_{L,n} \right)' M^{(Z_1, Q)} \left(y - X \widehat{\delta}_{L,n} \right)}.$$

Next, from expression (12), we have $\widehat{\ell}_{F,n} = \widehat{\ell}_{L,n} + O_p(n^{-1})$. It then follows, by tedious but

straightforward calculations² and by making use of Assumptions 1, 2(i), 3-6, and 8 that

$$\begin{aligned}
& D_\mu^{-1} X' \left[A - \widehat{\ell}_{F,n} M^{(Z_1, Q)} \right] (y - X\delta_0) \\
&= D_\mu^{-1} X' A (y - X\delta_0) - \widehat{\ell}_{L,n} D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\delta_0) + O_p \left(\frac{1}{n} \right) O_p \left(\frac{n}{\mu_n^{\min}} \right) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' A \varepsilon - \frac{\varepsilon' A \varepsilon}{\varepsilon' M^Q \varepsilon} [1 + o_p(1)] \left[\frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' M^{(Z_1, Q)} \varepsilon \right] \\
&\quad + O_p \left(\frac{1}{\mu_n^{\min}} \right) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} (U - \varepsilon \rho')' A \varepsilon [1 + o_p(1)] - \varepsilon' A \varepsilon D_\mu^{-1} \left[\frac{U' M^{(Z_1, Q)} \varepsilon}{\varepsilon' M^Q \varepsilon} - \rho \right] [1 + o_p(1)] \\
&\quad - \frac{\sqrt{K_{2,n}} \varepsilon' A \varepsilon / \sqrt{K_{2,n}}}{\varepsilon' M^Q \varepsilon / n} \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} [1 + o_p(1)] + O_p \left(\frac{1}{\mu_n^{\min}} \right) \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} (U - \varepsilon \rho')' A \varepsilon [1 + o_p(1)] + O_p \left(\frac{1}{\mu_n^{\min}} \max \left\{ \sqrt{\frac{K_{2,n}}{n}}, \frac{K_{1,n} \sqrt{K_{2,n}}}{n} \right\} \right) \\
&\quad + O_p \left(\frac{\sqrt{K_{2,n}}}{n} \right) + O_p \left(\frac{1}{\mu_n^{\min}} \right) \\
&= \left(\frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} \underline{U}' A \varepsilon \right) [1 + o_p(1)], \text{ where } \underline{U} = U - \varepsilon \rho'. \quad \square
\end{aligned}$$

Lemma S2-14: For any $a \in \mathbb{R}^d$ such that $\|a\| = 1$, define $b_{1n} = \Sigma_n^{-1/2} a$, $b_{2n} = \sqrt{K_{2,n}} D_\mu^{-1} \Sigma_n^{-1/2} a$, $\underline{u}_{2,(i,t),n} = b'_{2n} \underline{U}_{(i,t)}$, $= \sqrt{K_{2,n}} a' \Sigma_n^{-1/2} D_\mu^{-1} \underline{U}_{(i,t)}$, $\sigma_{(i,t),n}^2 = E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z]$, $\tilde{\psi}_{(i,t),n} = E[\underline{u}_{2,(i,t),n} \varepsilon_{(i,t)} | \mathcal{F}_n^Z]$, and $\tilde{\omega}_{(i,t)}^2 = E[\underline{u}_{2,(i,t),n}^2 | \mathcal{F}_n^Z]$. If Assumptions 1-2 and 5-6 are satisfied; then, the following statements are true.

(a) $\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i,t)} / \sqrt{n}] (A_{(i,t),(j,s)} / \sqrt{K_{2,n}}) \left\{ \varepsilon_{(j,s)} \tilde{\psi}_{(i,t),n} + \underline{u}_{2,(j,s)} \sigma_{(i,t),n}^2 \right\} = O_p(K_{2,n}^{1/4} / \mu_n^{\min}) = o_p(1)$.

(b) $\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} (A_{(i,t),(j,s)}^2 / K_{2,n}) (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 = O_p(K_{2,n} (\mu_n^{\min})^{-2} n^{-1/2}) = o_p(1)$.

(c) $\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} (A_{(i,t),(j,s)}^2 / K_{2,n}) (\underline{u}_{2,(j,s),n}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 = O_p(K_{2,n} (\mu_n^{\min})^{-2} n^{-1/2}) = o_p(1)$.

Proof of Lemma S2-14:

To show part (a), first let $\mathfrak{W}_n = \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i,t)} / \sqrt{n}] (A_{(i,t),(j,s)} / \sqrt{K_{2,n}}) \times \left\{ \varepsilon_{(j,s)} \tilde{\psi}_{(i,t),n} + \underline{u}_{2,(j,s),n} \sigma_{(i,t),n}^2 \right\}$. By taking expectation and applying the triangle inequality, we obtain $E[\mathfrak{W}_n^2 | \mathcal{F}_n^Z] \leq \mathcal{H}_1 + \mathcal{H}_2 + \mathcal{H}_3 + \mathcal{H}_4$, where $\mathcal{H}_1 = \sum_{(i,t),(k,v)=2}^{m_n} |n^{-1} [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i,t)}]|$

²Further details are available from the authors upon request.

$$\begin{aligned}
& \times [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(k, v)}] \tilde{\psi}_{(i, t), n} \tilde{\psi}_{(k, v), n} \sum_{(j, s)=1}^{\min\{(i, t), (k, v)\}-1} \left(A_{(i, t), (j, s)} A_{(k, v), (j, s)} \sigma_{(j, s), n}^2 / K_{2, n} \right) \Big|, \\
\mathcal{H}_2 &= \sum_{(i, t), (k, v)=2}^{m_n} \left| \left([b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i, t)}] [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(k, v)}] / n \right) \sigma_{(i, t), n}^2 \tilde{\psi}_{(k, v), n} \right. \\
&\quad \times \sum_{(j, s)=1}^{\min\{(i, t), (k, v)\}-1} \left. \left(A_{(i, t), (j, s)} A_{(k, v), (j, s)} \tilde{\psi}_{(j, s), n} / K_{2, n} \right) \right|, \mathcal{H}_3 = \sum_{(i, t), (k, v)=2}^{m_n} \left| n^{-1} [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i, t)}] \right. \\
&\quad \times [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(k, v)}] \sigma_{(k, v), n}^2 \tilde{\psi}_{(i, t), n} \sum_{(j, s)=1}^{\min\{(i, t), (k, v)\}-1} \left. \left(A_{(i, t), (j, s)} A_{(k, v), (j, s)} \tilde{\psi}_{(j, s), n} / K_{2, n} \right) \right|, \\
\mathcal{H}_4 &= \sum_{(i, t), (k, v)=2}^{m_n} \left| \left([b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i, t)}] [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(k, v)}] / n \right) \sigma_{(i, t), n}^2 \sigma_{(k, v), n}^2 \right. \\
&\quad \times \sum_{(j, s)=1}^{\min\{(i, t), (k, v)\}-1} \left. \left(A_{(i, t), (j, s)} A_{(k, v), (j, s)} \tilde{\omega}_{(j, s), n}^2 / K_{2, n} \right) \right|. \text{ Focusing first on } \mathcal{H}_1, \text{ we obtain, by applying the CS inequality,}
\end{aligned}$$

$$\begin{aligned}
& \mathcal{H}_1 \\
& \leq \sqrt{\sum_{(i, t), ((k, v))=2}^{m_n} \frac{b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i, t)} e'_{(i, t)} M^{(Z_1, Q)} Z_2 \Upsilon b_{1n} \tilde{\psi}_{(i, t), n}^2}{n} \frac{b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(k, v)} e'_{(k, v)} M^{(Z_1, Q)} Z_2 \Upsilon b_{1n} \tilde{\psi}_{(k, v), n}^2}{n}} \\
& \quad \times \sqrt{\sum_{(i, t)=2}^{m_n} \sum_{(k, v)=2}^{m_n} \left(\sum_{(j, s)=1}^{\min\{(i, t), (k, v)\}-1} \frac{A_{(i, t), (j, s)} A_{(k, v), (j, s)} \sigma_{(j, s), n}^2}{K_{2, n}} \right)^2}
\end{aligned}$$

Applying the CS inequality, Assumption 2(i), part (d) of Lemma S2-3, and Lemma S2-6, we obtain

$$\begin{aligned}
\max_{1 \leq (i, t) \leq m_n} |\tilde{\psi}_{(i, t), n}| &= \max_{1 \leq (i, t) \leq m_n} \sqrt{K_{2, n}} E \left[\left| \varepsilon_{(i, t)} \underline{U}'_{(i, t)} D_\mu^{-1} \Sigma_n^{-1/2} a \right| \mid \mathcal{F}_n^Z \right] \\
&\leq \frac{\sqrt{K_{2, n}}}{(\mu_n^{\min})} \sqrt{a' \Sigma_n^{-1} a} \sqrt{\max_{1 \leq (i, t) \leq m_n} E \left[\varepsilon_{(i, t)}^2 \mid \mathcal{F}_n^Z \right]} \sqrt{\max_{1 \leq (i, t) \leq m_n} E \left[\left\| \underline{U}_{(i, t)} \right\|_2^2 \mid \mathcal{F}_n^Z \right]} \\
&= O_{a.s.} \left(\frac{\sqrt{K_{2, n}}}{(\mu_n^{\min})} \right)
\end{aligned} \tag{29}$$

Moreover, by direct calculations,

$$\sum_{(i, t)=2}^{m_n} \sum_{(k, v)=2}^{m_n} \left(\sum_{(j, s)=1}^{\min\{(i, t), (k, v)\}-1} A_{(i, t), (j, s)} A_{(k, v), (j, s)} \sigma_{(j, s), n}^2 / K_{2, n} \right)^2 = K_{2, n}^{-2} \operatorname{tr} \{ L D_{\sigma^2} L' L D_{\sigma^2} L' \},$$

where L is the lower triangular matrix such that $L_{(i, t), (j, s)} = A_{(i, t), (j, s)} \mathbb{I}\{(i, t) > (j, s)\}$ and $D_{\sigma^2} = \operatorname{diag}(\sigma_{(1, 1), n}^2, \dots, \sigma_{(n, T_n), n}^2) = \operatorname{diag}(\sigma_{1, n}^2, \dots, \sigma_{m_n, n}^2)$ and where $\sigma_{(i, t), n}^2 = E \left[\varepsilon_{(i, t)}^2 \mid \mathcal{F}_n^Z \right]$ for $(i, t) = 1, \dots, m_n$. In addition, by the result shown in Lemma OA-11 (given in section 3 of this appendix), we have $\|LL'\|_F = O_{a.s.}(\sqrt{K_{2, n}})$. Using these results, we further deduce, by applying the CS

inequality, Assumptions 2 and 3(iii), and part (d) of Lemma S2-3 that

$$\begin{aligned}
\mathcal{H}_1 &\leq \left(\max_{1 \leq (i,t) \leq m_n} |\tilde{\psi}_{(i,t),n}| \right)^2 \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \frac{1}{K_{2,n}} \sqrt{\text{tr} \{ L' L D_{\sigma^2} L' L D_{\sigma^2} \}} \\
&\leq \left(\max_{1 \leq (i,t) \leq m_n} |\tilde{\psi}_{(i,t),n}| \right)^2 \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \frac{1}{K_{2,n}} (\text{tr} \{ L' L D_{\sigma^2}^2 L' L \})^{1/4} (\text{tr} \{ D_{\sigma^2} L' L L' L D_{\sigma^2} \})^{1/4} \\
&\leq \left(\max_{1 \leq (i,t) \leq m_n} |\tilde{\psi}_{(i,t),n}| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^2 \right) \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \frac{1}{K_{2,n}} \|L L'\|_F \\
&= O_{a.s.} \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right). \tag{30}
\end{aligned}$$

Similarly, let $D_{\tilde{\psi}} = \text{diag} (\tilde{\psi}_{(1,1),n}, \dots, \tilde{\psi}_{(n,T_n),n}) = \text{diag} (\tilde{\psi}_{1,n}, \dots, \tilde{\psi}_{m_n,n})$, we can also show

$$\begin{aligned}
\mathcal{H}_2 &\leq \left(\max_{1 \leq (i,t) \leq m_n} |\tilde{\psi}_{(i,t),n}| \right) \left(\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^2 \right) \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \frac{1}{K_{2,n}} \sqrt{\text{tr} \{ L' L D_{\tilde{\psi}} L' L D_{\tilde{\psi}} \}} \\
&\leq \left(\max_{1 \leq (i,t) \leq m_n} |\tilde{\psi}_{(i,t),n}| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^2 \right) \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \frac{1}{K_{2,n}} \|L L'\|_F \\
&= O_{a.s.} \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right), \tag{31}
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{H}_3 &\leq \left(\max_{1 \leq (i,t) \leq m_n} |\tilde{\psi}_{(i,t),n}| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^2 \right) \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \frac{1}{K_{2,n}} \|L L'\|_F \\
&= O_{a.s.} \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right). \tag{32}
\end{aligned}$$

Moreover, let $D_{\tilde{\omega}^2} = \text{diag} (\tilde{\omega}_{(1,1),n}^2, \dots, \tilde{\omega}_{(n,T_n),n}^2) = \text{diag} (\tilde{\omega}_{1,n}^2, \dots, \tilde{\omega}_{m_n,n}^2)$, and note that

$$\begin{aligned}
\mathcal{H}_4 &\leq \left(\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^2 \right)^2 \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \frac{1}{K_{2,n}} \sqrt{\text{tr} \{ L' L D_{\tilde{\omega}^2} L' L D_{\tilde{\omega}^2} \}} \\
&\leq \left(\max_{1 \leq (i,t) \leq m_n} \sigma_{(i,t),n}^2 \right)^2 \left(\frac{b'_{1n} \Upsilon' Z'_2 Z_2 \Upsilon b_{1n}}{n} \right) \left(\max_{1 \leq (i,t) \leq m_n} \tilde{\omega}_{(i,t),n}^2 \right) \frac{\|L L'\|_F}{K_{2,n}} \\
&= O_{a.s.} \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right), \tag{33}
\end{aligned}$$

where the order of magnitude above is calculated by applying Assumptions 2 and 3(iii), part (d)

of Lemma S2-3, and the fact that $\|LL'\|_F = O_{a.s.}(\sqrt{K_{2,n}})$ and by making use of the result

$$\begin{aligned} \max_{1 \leq (i,t) \leq m_n} \tilde{\omega}_{(i,t),n}^2 &= \max_{1 \leq (i,t) \leq m_n} K_{2,n} a' \Sigma_n^{-1/2} D_\mu^{-1} E \left[\underline{U}_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z \right] D_\mu^{-1} \Sigma_n^{-1/2} a \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E \left[\left\| \underline{U}_{(i,t)} \right\|_2^2 | \mathcal{F}_n^Z \right] \right) \frac{K_{2,n} a' \Sigma_n^{-1} a}{(\mu_n^{\min})^2} \\ &= O_{a.s.} \left(\frac{K_{2,n}}{(\mu_n^{\min})^2} \right) \end{aligned} \quad (34)$$

which can be easily deduced from part (d) of Lemma S2-3 and Lemma S2-6. Combining (30)-(33), we obtain $E [\mathfrak{W}_n^2 | \mathcal{F}_n^Z] = O_{a.s.}(\sqrt{K_{2,n}} (\mu_n^{\min})^{-2})$. Hence, by the law of iterated expectations and by Theorem 16.1 of Billingsley (1995), there exists a constant $\bar{C} < \infty$ such that, for all n sufficiently large, $E \left[(\mu_n^{\min})^2 / \sqrt{K_{2,n}} \right] \mathfrak{W}_n^2 = E_Z \left(E \left\{ \left[(\mu_n^{\min})^2 / \sqrt{K_{2,n}} \right] \mathfrak{W}_n^2 | \mathcal{F}_n^Z \right\} \right) \leq \bar{C}$. It follows from the Markov's inequality that

$$\begin{aligned} \mathfrak{W}_n &= \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} [b'_{1n} \Upsilon' Z'_2 M^{(Z_1, Q)} e_{(i,t)} / \sqrt{n}] (A_{(i,t),(j,s)} / \sqrt{K_{2,n}}) \left\{ \varepsilon_{(j,s)} \tilde{\psi}_{(i,t),n} + \underline{u}_{2,(j,s),n} \sigma_{(i,t),n}^2 \right\} \\ &= O_p \left(K_{2,n}^{1/4} / (\mu_n^{\min}) \right) = o_p(1). \end{aligned}$$

For part (b), note that, by Assumption 2, the symmetry of A , part (c) of Lemma S2-1, and expression (34) above; we obtain

$$\begin{aligned} &E \left\{ \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 \right)^2 | \mathcal{F}_n^Z \right\} \\ &= \frac{1}{K_{2,n}^2} \sum_{(i,t)=2}^{m_n} \sum_{(k,v)=2}^{m_n} \sum_{(j,s)=1}^{\min\{(i,t)-1, (k,v)-1\}} A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 \tilde{\omega}_{(i,t),n}^2 \tilde{\omega}_{(k,v),n}^2 E \left[(\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2)^2 | \mathcal{F}_n^Z \right] \\ &\leq \frac{1}{K_{2,n}^2} \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}} A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 \tilde{\omega}_{(i,t),n}^2 \tilde{\omega}_{(k,v),n}^2 \left\{ E \left[\varepsilon_{(j,s)}^4 | \mathcal{F}_n^Z \right] + \sigma_{(j,s),n}^4 \right\} \\ &= O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right) \end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$\begin{aligned} &E \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 \right)^2 \right] \\ &= E_Z \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} E \left\{ \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 \right)^2 | \mathcal{F}_n^Z \right\} \right] \\ &\leq \bar{C} \end{aligned}$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\begin{aligned}
& \Pr \left(\frac{(\mu_n^{\min})^2 \sqrt{n}}{K_{2,n}} \left| \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 \right| \geq \sqrt{\frac{C}{\epsilon}} \right) \\
&= \Pr \left(\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 \right)^2 \geq \frac{C}{\epsilon} \right) \\
&\leq \frac{\epsilon}{C} E \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 \right)^2 \right] \\
&\leq \epsilon
\end{aligned}$$

for all n sufficiently large, which shows that

$$\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\varepsilon_{(j,s)}^2 - \sigma_{(j,s),n}^2) \tilde{\omega}_{(i,t),n}^2 = O_p \left(\frac{K_{2,n}}{(\mu_n^{\min})^2 \sqrt{n}} \right) = o_p(1)$$

For part (c), note that, by Assumption 2, the symmetry of A , part (c) of Lemma S2-1, and expression (34) above; we get

$$\begin{aligned}
& E \left\{ \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\underline{u}_{2,(j,s)}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 \right)^2 | \mathcal{F}_n^Z \right\} \\
&= \frac{1}{K_{2,n}^2} \sum_{(i,t)=2}^{m_n} \sum_{(k,v)=2}^{m_n} \sum_{(j,s)=1}^{\min\{(i,t)-1, (k,v)-1\}} A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 \sigma_{(i,t),n}^2 \sigma_{(k,v),n}^2 E \left[(\underline{u}_{2,(j,s),n}^2 - \tilde{\omega}_{(j,s),n}^2)^2 | \mathcal{F}_n^Z \right] \\
&\leq \frac{1}{K_{2,n}^2} \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 \sigma_{(i,t),n}^2 \sigma_{(k,v),n}^2 \left\{ E \left[\underline{u}_{2,(j,s),n}^4 | \mathcal{F}_n^Z \right] + \tilde{\omega}_{(j,s),n}^4 \right\} \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right)
\end{aligned}$$

where, in obtaining the result above, we have also calculated the almost sure order of magnitude

of $E \left[\underline{u}_{2,(j,s),n}^4 | \mathcal{F}_n^Z \right]$ as follows

$$\begin{aligned}
E \left[\underline{u}_{2,(j,s),n}^4 | \mathcal{F}_n^Z \right] &= E \left[\left(\sqrt{K_{2,n}} a' \Sigma_n^{-1/2} D_\mu^{-1} \underline{U}_{(i,t)} \right)^4 | \mathcal{F}_n^Z \right] \\
&= K_{2,n}^2 E \left[\left(a' \Sigma_n^{-1/2} D_\mu^{-1} \underline{U}_{(i,t)} \underline{U}'_{(i,t)} D_\mu^{-1} \Sigma_n^{-1/2} a \right)^2 | \mathcal{F}_n^Z \right] \\
&\leq E \left[\left\| \underline{U}_{(i,t)} \right\|_2^4 | \mathcal{F}_n^Z \right] [a' \Sigma_n^{-1} a]^2 \frac{K_{2,n}^2}{(\mu_n^{\min})^4} \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4} \right). \tag{35}
\end{aligned}$$

using part (d) of Lemma S2-3 and Lemma S2-6. Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$\begin{aligned}
&E \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\underline{u}_{2,(j,s)}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 \right)^2 \right] \\
&= E_Z \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} E \left\{ \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\underline{u}_{2,(j,s)}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 \right)^2 | \mathcal{F}_n^Z \right\} \right] \\
&\leq \bar{C}
\end{aligned}$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\begin{aligned}
&\Pr \left(\left| \frac{(\mu_n^{\min})^2 \sqrt{n}}{K_{2,n}} \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\underline{u}_{2,(j,s)}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 \right) \right| \geq \sqrt{\frac{\bar{C}}{\epsilon}} \right) \\
&= \Pr \left(\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\underline{u}_{2,(j,s)}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 \right)^2 \geq \frac{\bar{C}}{\epsilon} \right) \\
&\leq \frac{\epsilon}{\bar{C}} E \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\underline{u}_{2,(j,s)}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 \right)^2 \right] \\
&\leq \epsilon
\end{aligned}$$

for all n sufficiently large, which shows that

$$\sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} \frac{A_{(i,t),(j,s)}^2}{K_{2,n}} (\underline{u}_{2,(j,s),n}^2 - \tilde{\omega}_{(j,s),n}^2) \sigma_{(i,t),n}^2 = O_p \left(\frac{K_{2,n}}{(\mu_n^{\min})^2 \sqrt{n}} \right) = o_p(1),$$

as required. \square

Lemma S2-15: Let $\{X_{i,n}, \mathcal{F}_{i,n}, 1 \leq i \leq k_n, n \geq 1\}$ be a square integrable martingale difference

array. Suppose that for all $\epsilon > 0$

$$\sum_{i=1}^{k_n} E \left[X_{i,n}^2 \mathbb{I}\{|X_{i,n}| > \epsilon\} | \mathcal{F}_{i-1,n} \right] \xrightarrow{p} 0 \quad (36)$$

and

$$\sum_{i=1}^{k_n} E \left[X_{i,n}^2 | \mathcal{F}_{i-1,n} \right] \xrightarrow{p} 1. \quad (37)$$

Then, $\sum_{i=1}^{k_n} X_{i,n} \xrightarrow{d} N(0, 1)$.

Proof of Lemma S2-15: The proof of this central limit theorem for square integrable martingale difference array is given in Gänssler and Stute (1977). See also Corollary 3.1 in Hall and Heyde (1980).

Remark: Note that a sufficient condition for condition (36), which we will verify in lieu of (36) in the proof of Theorems 2 and 3 in Appendix S1, is the following

$$\sum_{i=1}^{k_n} E \left[|X_{i,n}|^{2+\delta} \right] \xrightarrow{p} 0, \text{ for some } \delta > 0. \quad (38)$$

Lemma S2-16: Let \tilde{L}_n be a sequence of $l \times d$ nonrandom matrices (with $l \leq d$) such that $\|\tilde{L}_n\|_F^2 \leq \bar{C} < \infty$ for some constant \bar{C} , and let $\Sigma_{2,n} = VC(D_\mu^{-1} \underline{U}' A\varepsilon | \mathcal{F}_n^Z)$ $= D_\mu^{-1} VC(\underline{U}' A\varepsilon | \mathcal{F}_n^Z) D_\mu^{-1}$. Assume that there exists a positive constant \underline{C} such that $\lambda_{\min}((\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}_n' / K_{2,n}) \geq \underline{C} > 0$ a.s.n. Furthermore, let $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$ and let $\underline{u}_{a,(i,t),n} = a' ((\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}_n' / K_{2,n})^{-1/2} \tilde{L}_n D_\mu^{-1} \underline{U}_{(i,t)}$. Let Assumptions 1-2 and 5-6 be satisfied and assume that $(\mu_n^{\min})^2 / K_{2,n} = o(1)$ but $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Under these conditions, the following statements are true:

$$(a) \left[(\mu_n^{\min})^2 / K_{2,n} \right] \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 \left(\underline{u}_{a,(j,s),n}^2 - E \left[\underline{u}_{a,(j,s),n}^2 | \mathcal{F}_n^Z \right] \right) E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \\ = O_p(n^{-1/2}) = o_p(1);$$

$$(b) \left[(\mu_n^{\min})^2 / K_{2,n} \right] \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 \left(\varepsilon_{(j,s)}^2 - E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right] \right) E \left[\underline{u}_{a,(i,t),n}^2 | \mathcal{F}_n^Z \right] \\ = O_p(n^{-1/2}) = o_p(1).$$

Proof of Lemma S2-16:

To proceed, note first that Lemma S2-6 along with the assumptions on

$\lambda_{\min} \left((\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}'_n / K_{2,n} \right)$ and $\left\| \tilde{L}_n \right\|_F^2$ together imply that

$$\begin{aligned}
E \left[\underline{u}_{a,(i,t),n}^2 | \mathcal{F}_n^Z \right] &= a' \left(\frac{(\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}'_n}{K_{2,n}} \right)^{-1/2} \tilde{L}_n D_\mu^{-1} E \left[\underline{U}_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z \right] \\
&\quad D_\mu^{-1} \tilde{L}'_n \left(\frac{(\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}'_n}{K_{2,n}} \right)^{-1/2} a \\
&\leq \frac{1}{(\mu_n^{\min})^2} \frac{\max_{1 \leq (i,t) \leq m_n} E \left[\left\| \underline{U}_{(i,t)} \right\|_2^2 | \mathcal{F}_n^Z \right] \left\| \tilde{L}_n \right\|_F^2}{\lambda_{\min} \left((\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}'_n / K_{2,n} \right)} \\
&= O_{a.s.} \left(\frac{1}{(\mu_n^{\min})^2} \right)
\end{aligned} \tag{39}$$

and that

$$\begin{aligned}
E \left[\underline{u}_{a,(i,t),n}^4 | \mathcal{F}_n^Z \right] &= E \left[\left(a' \left(\frac{(\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}'_n}{K_{2,n}} \right)^{-1/2} \tilde{L}_n D_\mu^{-1} \underline{U}_{(i,t)} \underline{U}'_{(i,t)} D_\mu^{-1} \tilde{L}'_n \right. \right. \\
&\quad \times \left. \left. \left(\frac{(\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}'_n}{K_{2,n}} \right)^{-1/2} a \right)^2 | \mathcal{F}_n^Z \right] \\
&\leq \frac{1}{(\mu_n^{\min})^4} \frac{\max_{1 \leq (i,t) \leq m_n} E \left[\left\| \underline{U}_{(i,t)} \right\|_2^4 | \mathcal{F}_n^Z \right] \left\| \tilde{L}_n \right\|_F^4}{\left[\lambda_{\min} \left((\mu_n^{\min})^2 \tilde{L}_n H_n^{-1} \Sigma_{2,n} H_n^{-1} \tilde{L}'_n / K_{2,n} \right) \right]^2} \\
&= O_{a.s.} \left(\frac{1}{(\mu_n^{\min})^4} \right)
\end{aligned} \tag{40}$$

For part (a), define

$\mathfrak{Z}_n = \left[(\mu_n^{\min})^2 / K_{2,n} \right] \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 \left(\underline{u}_{a,(j,s),n}^2 - E \left[\underline{u}_{a,(j,s),n}^2 | \mathcal{F}_n^Z \right] \right) E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right]$, and note that we can apply Assumption 2(i), part (c) of Lemma S2-1, and the upper bounds given by (39) and (40) above to obtain

$$\begin{aligned}
E \left[\mathfrak{Z}_n^2 | \mathcal{F}_n^Z \right] &\leq \frac{(\mu_n^{\min})^4}{K_{2,n}^2} \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}}^{m_n} \left(A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(k,v)}^2 | \mathcal{F}_n^Z \right] \right. \\
&\quad \times \left. \left\{ E \left[\underline{u}_{a,(j,s),n}^4 | \mathcal{F}_n^Z \right] + \left(E \left[\underline{u}_{a,(j,s),n}^2 | \mathcal{F}_n^Z \right] \right)^2 \right\} \right) \\
&\leq O_{a.s.} \left(\frac{(\mu_n^{\min})^4}{K_{2,n}^2} \right) O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) O_{a.s.} (1) O_{a.s.} (1) O_{a.s.} \left(\frac{1}{(\mu_n^{\min})^4} \right) = O_{a.s.} \left(\frac{1}{n} \right)
\end{aligned}$$

Hence, the law of iterated expectations and Theorem 16.1 of Billingsley (1995) imply that there exists a positive constant $\bar{C} < \infty$ such that, for all n sufficiently large, $E(n\mathfrak{Z}_n^2) = E_Z(nE[\mathfrak{Z}_n^2|\mathcal{F}_n^Z]) \leq \bar{C}$. Application of the Markov's inequality then implies that $\mathfrak{Z}_n = O_p(n^{-1/2}) = o_p(1)$, which shows part (a).

For part (b), we can apply Assumption 2, part (c) of Lemma S2-1, and the result given in expression (39) above to obtain

$$\begin{aligned}
& E \left\{ \left(\frac{(\mu_n^{\min})^2}{K_{2,n}} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 (\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2|\mathcal{F}_n^Z]) E[\underline{u}_{a,(i,t),n}^2|\mathcal{F}_n^Z] \right)^2 |\mathcal{F}_n^Z \right\} \\
&= \frac{(\mu_n^{\min})^4}{K_{2,n}^2} \sum_{(i,t)=2}^{m_n} \sum_{(k,v)=2}^{m_n} \sum_{(j,s)=1}^{\min\{(i,t)-1, (k,v)-1\}} \left(A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 \right. \\
&\quad \times E[\underline{u}_{a,(i,t),n}^2|\mathcal{F}_n^Z] E[\underline{u}_{a,(k,v),n}^2|\mathcal{F}_n^Z] E\left[\left(\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2|\mathcal{F}_n^Z]\right)^2|\mathcal{F}_n^Z\right] \Big) \\
&\leq \frac{(\mu_n^{\min})^4}{K_{2,n}^2} \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}}^{m_n} \left(A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 E[\underline{u}_{a,(i,t),n}^2|\mathcal{F}_n^Z] E[\underline{u}_{a,(k,v),n}^2|\mathcal{F}_n^Z] \right. \\
&\quad \times \left. \left\{ E[\varepsilon_{(j,s)}^4|\mathcal{F}_n^Z] + \left(E[\varepsilon_{(j,s)}^2|\mathcal{F}_n^Z]\right)^2 \right\} \right) \\
&= O_{a.s.}\left(\frac{(\mu_n^{\min})^4}{K_{2,n}^2}\right) O_{a.s.}\left(\frac{K_{2,n}^2}{n}\right) O_{a.s.}\left(\frac{1}{(\mu_n^{\min})^2}\right) O_{a.s.}\left(\frac{1}{(\mu_n^{\min})^2}\right) \\
&= O_{a.s.}\left(\frac{1}{n}\right)
\end{aligned}$$

Hence, the law of iterated expectations and Theorem 16.1 of Billingsley (1995) imply that there exists a positive constant $\bar{C} < \infty$ such that for all n sufficiently large

$$\begin{aligned}
& nE \left[\left(\frac{(\mu_n^{\min})^2}{K_{2,n}} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 (\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2|\mathcal{F}_n^Z]) E[\underline{u}_{a,(i,t),n}^2|\mathcal{F}_n^Z] \right)^2 \right] \\
&= E_W \left\{ nE \left[\left(\frac{(\mu_n^{\min})^2}{K_{2,n}} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 (\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2|\mathcal{F}_n^Z]) E[\underline{u}_{a,(i,t),n}^2|\mathcal{F}_n^Z] \right)^2 |\mathcal{F}_n^Z \right] \right\} \\
&\leq \bar{C}
\end{aligned}$$

For any $\epsilon > 0$, set $\bar{C}_\epsilon = \sqrt{\bar{C}/\epsilon}$, and it follows by Markov's inequality that, for all n sufficiently

large,

$$\begin{aligned}
& \Pr \left(\sqrt{n} \left| \frac{(\mu_n^{\min})^2}{K_{2,n}} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 (\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z]) E[\underline{u}_{a,(i,t),n}^2 | \mathcal{F}_n^Z] \right| \geq \overline{C}_\epsilon \right) \\
&= \Pr \left(n \left(\frac{(\mu_n^{\min})^2}{K_{2,n}} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 (\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z]) E[\underline{u}_{a,(i,t),n}^2 | \mathcal{F}_n^Z] \right)^2 \geq \overline{C}_\epsilon^2 \right) \\
&\leq \frac{1}{\overline{C}_\epsilon^2} n E \left[\left(\frac{(\mu_n^{\min})^2}{K_{2,n}} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 (\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z]) E[\underline{u}_{a,(i,t),n}^2 | \mathcal{F}_n^Z] \right)^2 \right] \\
&\leq \frac{\overline{C}}{\overline{C}/\epsilon} = \epsilon
\end{aligned}$$

which shows that

$$\begin{aligned}
& \frac{(\mu_n^{\min})^2}{K_{2,n}} \sum_{(i,t)=2}^{m_n} \sum_{(j,s)=1}^{(i,t)-1} A_{(i,t),(j,s)}^2 (\varepsilon_{(j,s)}^2 - E[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z]) E[\underline{u}_{a,(i,t),n}^2 | \mathcal{F}_n^Z] \\
&= O_p \left(\frac{1}{\sqrt{n}} \right) = o_p(1). \quad \square
\end{aligned}$$

Lemma S2-17 Under Assumptions 1-6, $D_\mu^{-1} X' A D(\varepsilon \circ \varepsilon) A X D_\mu^{-1} = \Sigma_{1,n}$
 $+ \sum_{(i,t),(j,s)=1, (i,t) \neq (j,s)}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 D_\mu^{-1} \Psi_{(j,s)} D_\mu^{-1} + o_p(1)$, where $\Sigma_{1,n} = \Upsilon' Z_2' M^{(Z_1,Q)} D_{\sigma^2} M^{(Z_1,Q)} Z_2 \Upsilon / n$,
 $\sigma_{(i,t)}^2 = E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z]$, $D_{\sigma^2} = \text{diag}(\sigma_{(1,1)}^2, \dots, \sigma_{(n,T_n)}^2)$, and $\Psi_{(j,s)} = E[U_{(j,s)} U'_{(j,s)} | \mathcal{F}_n^Z]$.

Proof of Lemma S2-17: To proceed, using the fact that $A Z_1 \Theta D_\kappa / \sqrt{n} = 0$ and $A Q = 0$, we can write

$$\begin{aligned}
D_\mu^{-1} X' A D(\varepsilon \circ \varepsilon) A X D_\mu^{-1} &= D_\mu^{-1} \left(\frac{D_\kappa}{\sqrt{n}} \Theta' Z_1' + \frac{D_\mu}{\sqrt{n}} \Upsilon' Z_2' + \Xi' Q' + U' \right) A D(\varepsilon \circ \varepsilon) A \\
&\quad \times \left(Z_1 \Theta \frac{D_\kappa}{\sqrt{n}} + Z_2 \Upsilon \frac{D_\mu}{\sqrt{n}} + Q \Xi + U \right) D_\mu^{-1} \\
&= \frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A Z_2 \Upsilon}{n} + D_\mu^{-1} U' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} \\
&\quad + \frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1}}{\sqrt{n}} + \frac{D_\mu^{-1} U' A D(\varepsilon \circ \varepsilon) A Z_2 \Upsilon}{\sqrt{n}}
\end{aligned}$$

Hence, by applying part (b) of Lemma OA-8 as well as (a)-(e) of Lemma OA-9, we obtain

$$\begin{aligned}
& D_\mu^{-1} X' A D(\varepsilon \circ \varepsilon) A X D_\mu^{-1} - \frac{\Upsilon' Z_2' M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} \\
& - \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] D_\mu^{-1} E \left[U_{(j,s)} U'_{(j,s)} | \mathcal{F}_n^Z \right] D_\mu^{-1} \\
= & \frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A Z_2 \Upsilon}{n} - \frac{\Upsilon' Z_2' M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} \\
& + D_\mu^{-1} U' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} - \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] D_\mu^{-1} E \left[U_{(j,s)} U'_{(j,s)} | \mathcal{F}_n^Z \right] D_\mu^{-1} \\
& + \frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1}}{\sqrt{n}} + \frac{D_\mu^{-1} U' A D(\varepsilon \circ \varepsilon) A Z_2 \Upsilon}{\sqrt{n}} \\
= & o_p(1). \quad \square
\end{aligned}$$

Lemma S2-18 Let Assumptions 1-6 and 8 be satisfied, and let $\{\hat{\delta}_n\}$ be any sequence of estimators such that $\|\hat{\delta}_n - \delta_0\|_2 \xrightarrow{p} 0$ as $n \rightarrow \infty$, as long as $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Also, define the following notations: let $\hat{\varepsilon} = M^{(Z,Q)}(y - X\hat{\delta}_n)$, $J = [M^Q \circ M^Q]^{-1}$, $S_1 = X' A D(J[\hat{\varepsilon} \circ \hat{\varepsilon}]) A X$, $S_2 = (\hat{\varepsilon} \circ \hat{\varepsilon})' J(A \circ A) J(\hat{\varepsilon} \ell_d' \circ M^{(Z,Q)} X)$, $S_2 = (\hat{\varepsilon} \circ \hat{\varepsilon})' J(A \circ A) J(\hat{\varepsilon} \ell_d' \circ \underline{U})$ with $\underline{U} = M^{(Z,Q)} X - \hat{\varepsilon} \hat{\rho}'_n$, $S_3 = (\hat{\varepsilon} \circ \hat{\varepsilon})' J(A \circ A) J(\hat{\varepsilon} \circ \hat{\varepsilon})$, $S_4 = (\hat{\varepsilon} \ell_d' \circ M^{(Z,Q)} X)' J(A \circ A) J(\hat{\varepsilon} \ell_d' \circ M^{(Z,Q)} X)$, $\underline{S}_4 = (\hat{\varepsilon} \ell_d' \circ \underline{U})' J(A \circ A) J(\hat{\varepsilon} \ell_d' \circ \underline{U})$, and $\Sigma_{1,n} = \Upsilon' Z_2' M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon / n$. In addition, define $\sigma_{(i,t)}^2 = E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right]$, $D_{\sigma^2} = \text{diag}(\sigma_{(1,1)}^2, \dots, \sigma_{(n,T_n)}^2)$, $\phi_{(i,t)} = E \left[U_{(i,t)} \varepsilon_{(i,t)} | \mathcal{F}_n^Z \right]$, $\Psi_{(i,t)} = E \left[U_{(i,t)} U'_{(i,t)} | \mathcal{F}_n^Z \right]$, $\underline{\phi}_{(i,t)} = E \left[\underline{U}_{(i,t)} \varepsilon_{(i,t)} | \mathcal{F}_n^Z \right]$, and $\underline{\Psi}_{(i,t)} = E \left[\underline{U}_{(i,t)} \underline{U}'_{(i,t)} | \mathcal{F}_n^Z \right]$ where $\underline{U}_{(i,t)} = U_{(i,t)} - \rho \varepsilon_{(i,t)}$ and where for notational convenience we suppress the dependence of $\sigma_{(i,t)}^2$, $\phi_{(i,t)}$, $\Psi_{(i,t)}$, $\underline{\phi}_{(i,t)}$, and $\underline{\Psi}_{(i,t)}$ on $\mathcal{F}_n^Z = \sigma(Z)$. Then, under the above conditions, the following statements are true.

- (a) $D_\mu^{-1} S_1 D_\mu^{-1} = \Sigma_{1,n} + \sum_{(i,t),(j,s)=1,(i,t) \neq (j,s)}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 \sigma_{(j,s)}^2 D_\mu^{-1} \Psi_{(j,s)} D_\mu^{-1}$
 $+ o_p \left(\max \left\{ 1, K_{2,n} (\mu_n^{\min})^{-2} \right\} \right).$
- (b) $S_3 / K_{2,n} - K_{2,n}^{-1} \sum_{(i,t),(j,s)=1,(i,t) \neq (j,s)}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 \sigma_{(j,s)}^2 = o_p(1)$.
- (c) $D_\mu^{-1} S_4 D_\mu^{-1} - \sum_{(i,t),(j,s)=1,(i,t) \neq (j,s)}^{m_n} A_{(i,t),(j,s)}^2 D_\mu^{-1} \phi_{(i,t)} \phi'_{(j,s)} D_\mu^{-1} = o_p \left(K_{2,n} (\mu_n^{\min})^{-2} \right)$.
- (d) $(\mu_n^{\min} / K_{2,n}) S_2 D_\mu^{-1} - (\mu_n^{\min} / K_{2,n}) \sum_{(i,t),(j,s)=1,(i,t) \neq (j,s)}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 \sigma_{(j,s)}^2 \phi'_{(j,s)} D_\mu^{-1} = o_p(1)$.

- (e) $D_\mu^{-1}\hat{\rho}_n = O_p\left(\left(\mu_n^{\min}\right)^{-1}\right)$ and $D_\mu^{-1}(\hat{\rho}_n - \rho) = o_p\left(\left(\mu_n^{\min}\right)^{-1}\right)$, where $\rho = \lim_{n \rightarrow \infty} \rho_n = \lim_{n \rightarrow \infty} (E[U'M^Q\varepsilon]/n)/(E[\varepsilon'M^Q\varepsilon]/n)$.
- (f) $D_\mu^{-1}S_4D_\mu^{-1} - \sum_{(i,t),(j,s)=1,(i,t)\neq(j,s)}^{m_n} A_{(i,t),(j,s)}^2 D_\mu^{-1} \underline{\phi}_{(i,t)} \underline{\phi}'_{(j,s)} D_\mu^{-1} = o_p\left(K_{2,n} (\mu_n^{\min})^{-2}\right)$.
- (g) $(\mu_n^{\min}/K_{2,n}) - (\mu_n^{\min}/K_{2,n}) \sum_{(i,t),(j,s)=1,(i,t)\neq(j,s)}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 \underline{\phi}'_{(j,s)} D_\mu^{-1} = o_p(1)$.

Proof of Lemma S2-18:

To show part (a), note first that, by making use of the decomposition $M^{(Z,Q)} = M^Q - P^{Z^\perp}$, where $P^{Z^\perp} = M^Q Z (Z' M^Q Z)^{-1} Z' M^Q$, we can write

$$\begin{aligned} \hat{\varepsilon} &= M^{(Z,Q)}(y - X\hat{\delta}_n) \\ &= M^{(Z,Q)}(y - X[\delta_0 - \delta_0 + \hat{\delta}_n]) \\ &= M^{(Z,Q)}(y - X\delta_0) - M^{(Z,Q)}X(\hat{\delta}_n - \delta_0) \\ &= M^{(Z,Q)}(Z_1\varphi_n + Q\alpha) + M^{(Z,Q)}\varepsilon - M^{(Z,Q)}X(\hat{\delta}_n - \delta_0) \\ &= M^{(Z,Q)}\varepsilon - M^{(Z,Q)}X(\hat{\delta}_n - \delta_0) \\ &= -M^{(Z,Q)}X(\hat{\delta}_n - \delta_0) + M^Q\varepsilon - P^{Z^\perp}\varepsilon, \end{aligned}$$

from which we obtain

$$\begin{aligned} J[\hat{\varepsilon} \circ \hat{\varepsilon}] &= J\left[M^{(Z,Q)}X(\hat{\delta}_n - \delta_0) \circ M^{(Z,Q)}X(\hat{\delta}_n - \delta_0)\right] + J[M^Q\varepsilon \circ M^Q\varepsilon] \\ &\quad + J[P^{Z^\perp}\varepsilon \circ P^{Z^\perp}\varepsilon] - 2J\left[M^Q\varepsilon \circ M^{(Z,Q)}X(\hat{\delta}_n - \delta_0)\right] \\ &\quad + 2J\left[P^{Z^\perp}\varepsilon \circ M^{(Z,Q)}X(\hat{\delta}_n - \delta_0)\right] - 2J\left[M^Q\varepsilon \circ P^{Z^\perp}\varepsilon\right] \end{aligned} \quad (41)$$

where $J = [M^Q \circ M^Q]^{-1}$. Substituting the right-hand side of (41) into covariance matrix estimator $D_\mu^{-1}X'AD(J[\hat{\varepsilon} \circ \hat{\varepsilon}])AXD_\mu^{-1}$, we get

$$\begin{aligned} &D_\mu^{-1}S_1D_\mu^{-1} - D_\mu^{-1}X'AD(\varepsilon \circ \varepsilon)AXD_\mu^{-1} \\ &= D_\mu^{-1}X'AD(J[\hat{\varepsilon} \circ \hat{\varepsilon}])AXD_\mu^{-1} - D_\mu^{-1}X'AD(\varepsilon \circ \varepsilon)AXD_\mu^{-1} \\ &= \mathcal{T}_{1,n} + \mathcal{T}_{2,n} + \mathcal{T}_{3,n} + \mathcal{T}_{4,n} + \mathcal{T}_{5,n} + \mathcal{T}_{6,n} + \mathcal{T}_{7,n} + \mathcal{T}_{8,n} + \mathcal{T}_{9,n} + \mathcal{T}_{10,n}, \end{aligned}$$

where $\mathcal{T}_{1,n} = D_\mu^{-1}X'AD\left(J\left[M^{(Z,Q)}X(\hat{\delta}_n - \delta_0) \circ M^{(Z,Q)}X(\hat{\delta}_n - \delta_0)\right]\right)AXD_\mu^{-1}$,
 $\mathcal{T}_{2,n} = D_\mu^{-1}X'AD\left(J\left[M^Q\varepsilon \circ M^Q\varepsilon\right]\right)AXD_\mu^{-1} - D_\mu^{-1}X'AD(\varepsilon \circ \varepsilon)AXD_\mu^{-1}$,
 $\mathcal{T}_{3,n} = D_\mu^{-1}X'AD\left(J\left[P^{Z^\perp}\varepsilon \circ P^{Z^\perp}\varepsilon\right]\right)AXD_\mu^{-1}$,
 $\mathcal{T}_{4,n} = -2D_\mu^{-1}X'AD\left(J\left[M^Q\varepsilon \circ M^{(Z,Q)}X(\hat{\delta}_n - \delta_0)\right]\right)AXD_\mu^{-1}$,
 $\mathcal{T}_{5,n} = 2D_\mu^{-1}X'AD\left(J\left[P^{Z^\perp}\varepsilon \circ M^{(Z,Q)}X(\hat{\delta}_n - \delta_0)\right]\right)AXD_\mu^{-1}$, and

$$\mathcal{T}_{6,n} = -2D_\mu^{-1}X'AD\left(J\left[M^Q\varepsilon \circ P^{Z^\perp}\varepsilon\right]\right)AXD_\mu^{-1}.$$

Consider the term $\mathcal{T}_{1,n}$. Let $\mathcal{G}_i = \{(\ell, h) : \ell = i \text{ and } h = 1, \dots, T_i\}$; and note that for any $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we have

$$\begin{aligned} |a'\mathcal{T}_{1,n}b| &= \left| \sum_{(i,t)=1}^{m_n} \sum_{(j,s)\neq(i,t)} \sum_{(k,v)\neq(i,t)} \sum_{(p,q)=1}^{m_n} a'D_\mu^{-1}X'e_{(j,s)}A_{(i,t),(j,s)}J_{(i,t),(p,q)} \right. \\ &\quad \times e'_{(p,q)}M^{(Z,Q)}X\left(\widehat{\delta}_n - \delta_0\right)\left(\widehat{\delta}_n - \delta_0\right)'X'M^{(Z,Q)}e_{(p,q)}A_{(i,t),(k,v)}e'_{(k,v)}XD_\mu^{-1}b \Big| \\ &= \left| \sum_{(i,t)=1}^{m_n} \sum_{(j,s)\neq(i,t)} a'D_\mu^{-1}X'e_{(j,s)}A_{(i,t),(j,s)} \right. \\ &\quad \times \left(\sum_{(p,q)=1}^{m_n} J_{(i,t),(p,q)}\mathbb{I}\{(p,q) \in \mathcal{G}_i\} e'_{(p,q)}M^{(Z,Q)}X\left(\widehat{\delta}_n - \delta_0\right)\left(\widehat{\delta}_n - \delta_0\right)'X'M^{(Z,Q)}e_{(p,q)} \right) \\ &\quad \times \left. \sum_{(k,v)\neq(i,t)} A_{(i,t),(k,v)}e'_{(k,v)}XD_\mu^{-1}b \right| \end{aligned}$$

where $J_{(i,t),(p,q)}$ is the element in the $(i,t)^{th}$ row and the $(p,q)^{th}$ column of the matrix J for $(i,t), (p,q) \in \{1, \dots, m_n\}$, where $\mathbb{I}\{\cdot\}$ denotes an indicator function, where $e_{(j,s)}$ is an $m_n \times 1$ elementary vector whose $(j,s)^{th}$ component is 1 and all other components are 0, and where the second equality above follows from the fact that $J_{(i,t),(p,q)} = 0$ for $p \neq i$ due to the sparsity (or block diagonal nature) of J . Now, let $D^{\Sigma J}(M^{(Z,Q)}XX'M^{(Z,Q)})$ be a diagonal matrix whose $(i,t)^{th}$ diagonal element is given by $\sum_{q=1}^{T_i} |J_{(i,t),(i,q)}| e'_{(i,q)}M^{(Z,Q)}XX'M^{(Z,Q)}e_{(i,q)}$. Applying the triangle inequality and the inequality $|XY| \leq (1/2)X^2 + (1/2)Y^2$, we then obtain

$$\begin{aligned}
|a' \mathcal{T}_{1,n} b| &= \left| \sum_{(i,t)=1}^{m_n} a' D_\mu^{-1} X' A e_{(i,t)} \sum_{q=1}^{T_i} J_{(i,t),(i,q)} e'_{(i,q)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) (\widehat{\delta}_n - \delta_0)' \right. \\
&\quad \times X' M^{(Z,Q)} e_{(i,q)} e'_{(i,t)} A X D_\mu^{-1} b \Big| \\
&\leq \frac{1}{2} \sum_{(i,t)=1}^{m_n} \sum_{q=1}^{T_i} \left\{ |J_{(i,t),(i,q)}| e'_{(i,q)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) (\widehat{\delta}_n - \delta_0)' X' M^{(Z,Q)} e_{(i,q)} \right. \\
&\quad \times a' D_\mu^{-1} X' A e_{(i,t)} e'_{(i,t)} A X D_\mu^{-1} a \Big\} \\
&\quad + \frac{1}{2} \sum_{(i,t)=1}^{m_n} \sum_{q=1}^{T_i} \left\{ |J_{(i,t),(i,q)}| e'_{(i,q)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) (\widehat{\delta}_n - \delta_0)' X' M^{(Z,Q)} e_{(i,q)} \right. \\
&\quad \times b' D_\mu^{-1} X' A e_{(i,t)} e'_{(i,t)} A X D_\mu^{-1} b \Big\} \\
&\leq \frac{1}{2} \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 a' D_\mu^{-1} X' A D^{\Sigma J} (M^{(Z,Q)} X X' M^{(Z,Q)}) A X D_\mu^{-1} a \\
&\quad + \frac{1}{2} \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 b' D_\mu^{-1} X' A D^{\Sigma J} (M^{(Z,Q)} X X' M^{(Z,Q)}) A X D_\mu^{-1} b \tag{42}
\end{aligned}$$

By tedious but straightforward calculations, we can show that

$D_\mu^{-1} X' A D^{\Sigma J} (M^{(Z,Q)} X X' M^{(Z,Q)}) A X D_\mu^{-1} = O_p \left(\max \left\{ 1, K_{2,n} (\mu_n^{\min})^{-2} \right\} \right)$. Hence, given the assumption that $\left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \xrightarrow{p} 0$, it follows from expression (42) that

$|a' \mathcal{T}_{1,n} b| = o_p \left(\max \left\{ 1, K_{2,n} (\mu_n^{\min})^{-2} \right\} \right)$. Since the above argument holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we further deduce that $\mathcal{T}_{1,n} = o_p \left(\max \left\{ 1, K_{2,n} (\mu_n^{\min})^{-2} \right\} \right)$. By following a similar method of proof, we can also show that $\mathcal{T}_{k,n} = o_p \left(\max \left\{ 1, K_{2,n} (\mu_n^{\min})^{-2} \right\} \right)$ for $k = 2, \dots, 6$.

The fact that $\mathcal{T}_{k,n} = o_p \left(\max \left\{ 1, K_{2,n} (\mu_n^{\min})^{-2} \right\} \right)$ for each $k = \{1, \dots, 6\}$ further implies that

$$D_\mu^{-1} S_1 D_\mu^{-1} - D_\mu^{-1} X' A D (\varepsilon \circ \varepsilon) A X D_\mu^{-1} = \sum_{k=1}^6 \mathcal{T}_{k,n} = o_p \left(\max \left\{ 1, \frac{K_{2,n}}{(\mu_n^{\min})^2} \right\} \right) \tag{43}$$

Moreover, by the result of Lemma S2-17,

$$D_\mu^{-1} X' A D (\varepsilon \circ \varepsilon) A X D_\mu^{-1} = \Sigma_{1,n} + \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 D_\mu^{-1} \Psi_{(j,s)} D_\mu^{-1} + o_p(1). \tag{44}$$

Combining (43) and (44), we further obtain

$D_\mu^{-1} S_1 D_\mu^{-1} = \Sigma_{1,n} + \sum_{(i,t),(j,s)=1, (i,t) \neq (j,s)}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 D_\mu^{-1} \Psi_{(j,s)} D_\mu^{-1} + o_p \left(\max \left\{ 1, K_{2,n} (\mu_n^{\min})^{-2} \right\} \right)$, which shows part (a).

To show part (b), write

$$\frac{S_3}{K_{2,n}} - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 \sigma_{(j,s)}^2 = \mathcal{A} + \mathfrak{A},$$

where

$$\begin{aligned} \mathcal{A} &= \frac{S_3}{K_{2,n}} - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2, \\ \mathfrak{A} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left(\varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 - \sigma_{(i,t)}^2 \sigma_{(j,s)}^2 \right) \end{aligned}$$

and where

$$S_3 = (\widehat{\varepsilon} \circ \widehat{\varepsilon})' J(A \circ A) J(\widehat{\varepsilon} \circ \widehat{\varepsilon})$$

To analyze the term \mathcal{A} , note that, by direct calculation, we can obtain the following decomposition

$$\begin{aligned} \mathcal{A} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left[e'_{(i,h)} M^{(Z,Q)} (y - X \widehat{\delta}_n) \right]^2 \right. \\ &\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left[(y - X \widehat{\delta}_n)' M^{(Z,Q)} e_{(j,v)} \right]^2 \left. \right\} - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 \\ &= \sum_{k=1}^3 \mathcal{A}_k + 2 \sum_{\ell=4}^6 \mathcal{A}_\ell \end{aligned} \tag{45}$$

where

$$\begin{aligned}
\mathcal{A}_1 &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} M^{Q_\varepsilon} \right)^2 \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \\
&\quad - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2, \\
\mathcal{A}_2 &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^{Q_\varepsilon} \right) \right. \\
&\quad \times \left. \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^{Q_\varepsilon} \right) \right\}, \\
\mathcal{A}_3 &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \right. \\
&\quad \times \left. \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left[e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \right\}, \\
\mathcal{A}_4 &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \right. \\
&\quad \times \left. \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^{Q_\varepsilon} \right) \right\}, \\
\mathcal{A}_5 &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right)^2 \right. \\
&\quad \times \left. \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \right\}, \text{ and} \\
\mathcal{A}_6 &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right)^2 \right. \\
&\quad \times \left. \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^{Q_\varepsilon} \right) \right\}.
\end{aligned}$$

It then follows from part (d) of Lemma OA-14; part (e) of Lemmas OA-15 and OA-16; as well as

part (c) of Lemmas OA-17, OA-18, and OA-19 that

$$\begin{aligned}\mathcal{A} &= \frac{S_3}{K_{2,n}} - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 \\ &= \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 + 2\mathcal{A}_4 + 2\mathcal{A}_5 + 2\mathcal{A}_6 = o_p(1).\end{aligned}\quad (46)$$

Now, for the term $\mathfrak{A} = K_{2,n}^{-1} \sum_{(i,t),(j,s)=1, (i,t) \neq (j,s)}^{m_n} A_{(i,t),(j,s)}^2 (\varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 - \sigma_{(i,t)}^2 \sigma_{(j,s)}^2)$, it can be shown by straightforward calculation and by using Assumptions 1, 2(i), 5 and 6 and Lemma S2-1(a) that $E[\mathfrak{A}^2 | \mathcal{F}_n^Z] = O_{a.s.}(n^{-1})$. It then follows by application of the law of iterated expectations, Theorem 16.1 of Billingsley (1995), and the Markov's inequality that

$$\mathfrak{A} = O_p\left(\frac{1}{\sqrt{n}}\right). \quad (47)$$

Combining (46) and (47), we get

$$\frac{S_3}{K_{2,n}} - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sigma_{(i,t)}^2 \sigma_{(j,s)}^2 = \mathcal{A} + \mathfrak{A} = o_p(1) + O_p\left(\frac{1}{\sqrt{n}}\right) = o_p(1),$$

which shows part (b).

The proofs for parts (c) and (d) are very similar to that of part (b). Hence, to avoid duplication, we will not provide detailed proofs of these parts here.

Turning our attention to part (e), note first that, we can write

$$\begin{aligned}\widehat{\rho}_n - \rho &= \frac{X' M^{(Z,Q)} (y - X\widehat{\delta}_n) / n}{(y - X\widehat{\delta}_n)' M^{(Z,Q)} (y - X\widehat{\delta}_n) / n} - \rho \\ &= \frac{X' M^{(Z,Q)} (y - X\widehat{\delta}_n) / n - U' M^{(Z_1,Q)} \varepsilon / n + U' M^{(Z_1,Q)} \varepsilon / n}{(y - X\widehat{\delta}_n)' M^{(Z,Q)} (y - X\widehat{\delta}_n) / n - \varepsilon' M^Q \varepsilon / n + \varepsilon' M^Q \varepsilon / n} - \rho\end{aligned}$$

where $\rho = \lim_{n \rightarrow \infty} \rho_n = \lim_{n \rightarrow \infty} (E[U' M^Q \varepsilon] / n) / (E[\varepsilon' M^Q \varepsilon] / n)$. By straightforward asymptotic analysis, we can show that

$$\begin{aligned}\frac{X' M^{(Z,Q)} (y - X\widehat{\delta}_n)}{n} - \frac{U' M^{(Z_1,Q)} \varepsilon}{n} &= o_p(1), \\ \frac{(y - X\widehat{\delta}_n)' M^{(Z,Q)} (y - X\widehat{\delta}_n)}{n} - \frac{\varepsilon' M^{(Z_1,Q)} \varepsilon}{n} &= o_p(1), \\ \frac{\varepsilon' M^{(Z_1,Q)} \varepsilon}{n} - \frac{\varepsilon' M^Q \varepsilon}{n} &= o_p(1),\end{aligned}$$

$$\frac{U'M^{(Z_1,Q)}\varepsilon}{\varepsilon'M^Q\varepsilon} - \rho = O_p\left(\max\left\{\frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n}\right\}\right) = o_p(1).$$

Next, note that, under Assumption 6(i), $T_i \geq 3$ for all i , so that $\frac{T_i-1}{T_i} \geq \frac{2}{3}$ for all i . Hence, by Assumption 2(ii), there exists a positive constant \underline{C} such that $E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \geq \underline{C} > 0$ a.s., so that

$$\begin{aligned} \frac{E[\varepsilon'M^Q\varepsilon]}{n} &= \frac{1}{n} E_Z \left\{ \sum_{i=1}^n \sum_{t=1}^{T_i} \left[\left(\frac{T_i-1}{T_i} \right) \right] E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right\} \\ &\geq \frac{2}{3n} E_Z \left\{ \sum_{i=1}^n \sum_{t=1}^{T_i} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right\} \\ &\geq \frac{2}{3n} E_Z \left[\sum_{i=1}^n \sum_{t=1}^{T_i} \underline{C} \right] = \frac{2}{3} \frac{m_n}{n} \underline{C} \quad \left(\text{since } m_n = \sum_{i=1}^n T_i \right) \\ &\geq \frac{2}{3} \underline{C} > 0 \end{aligned}$$

for all n sufficiently large. It follows from these results that $\hat{\rho}_n - \rho = [U'M^{(Z_1,Q)}\varepsilon / (\varepsilon'M^Q\varepsilon)] - \rho + o_p(1) = o_p(1)$ and $\|\hat{\rho}_n\|_2 \leq \|\hat{\rho}_n - \rho\|_2 + \|\rho\|_2 = O_p(1)$.

Now, let $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$; and note that, by applying the CS inequality, we have

$$|a'D_\mu^{-1}\hat{\rho}_n| \leq \frac{1}{(\mu_n^{\min})} \|\hat{\rho}_n\|_2 = O_p\left(\frac{1}{(\mu_n^{\min})}\right), \quad (48)$$

$$|a'D_\mu^{-1}(\hat{\rho}_n - \rho)| \leq \frac{1}{(\mu_n^{\min})} \|\hat{\rho}_n - \rho\|_2 = o_p\left(\frac{1}{(\mu_n^{\min})}\right), \quad (49)$$

Since the argument above holds for any $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$, we further deduce that $D_\mu^{-1}\hat{\rho}_n = O_p((\mu_n^{\min})^{-1})$ and $D_\mu^{-1}(\hat{\rho}_n - \rho) = o_p((\mu_n^{\min})^{-1})$, which shows part (e).

Part (f) can be shown by applying the results of parts (b), (c), (d), and (e) of this lemma as well as part (a) of Lemma S2-1 and Assumptions 2(i) and 3(ii). Part (g), on the other hand, can be proved by applying the results of parts (b), (d), and (e) of this lemma. \square

Section 2: Proof of Lemma 1 of the Main Paper

Lemma 1: Suppose that Assumptions 5 and 6(i) are satisfied. Then, there exists a positive constant C such that

$$\lambda_{\min}(M^{(Z,Q)} \circ M^{(Z,Q)}) \geq C > 0 \text{ a.s.}$$

for all n sufficiently large.

Proof of Lemma 1:

To show the required result, we shall apply Geršgorin's theorem (see Theorem 6.1.1 on page 344 of Horn and Johnson, 1985). To apply this theorem to the matrix $M^{(Z,Q)} \circ M^{(Z,Q)}$, we note first that this matrix is symmetric and its elements are real-valued, so that all the eigenvalues of this matrix are real. Now, let $e_{(i,t)}$ denote an $m_n \times 1$ elementary vector whose $(i,t)^{th}$ component is

1 and all other components are 0, and define

$$R'_{(i,t)} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) = \sum_{\substack{(j,s)=1 \\ (j,s) \neq (i,t)}}^{m_n} \left| e'_{(i,t)} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) e_{(j,s)} \right| \text{ for } (i,t) \in \{1, \dots, m_n\}.$$

Applying Geršgorin's theorem to the matrix $M^{(Z,Q)} \circ M^{(Z,Q)}$, we see that

$$\begin{aligned} & \lambda_{\min} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) \\ & \in \bigcup_{(i,t)=1}^{m_n} \left\{ z \in \mathbb{R} : \left| z - \left(M_{(i,t),(i,t)}^{(Z,Q)} \right)^2 \right| \leq R'_{(i,t)} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) \right\}, \end{aligned}$$

so that there exists at least one pair (k,v) (where $k = 1, \dots, n$ and $v = 1, \dots, T_k$ given k) such that

$$\begin{aligned} \left| \lambda_{\min} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) - \left(M_{(k,v),(k,v)}^{(Z,Q)} \right)^2 \right| & \leq R'_{(k,v)} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) \\ & = \sum_{\substack{(j,s)=1 \\ (j,s) \neq (k,v)}}^{m_n} \left(M_{(k,v),(j,s)}^{(Z,Q)} \right)^2 \end{aligned}$$

or

$$\begin{aligned} - \sum_{\substack{(j,s)=1 \\ (j,s) \neq (k,v)}}^{m_n} \left(M_{(k,v),(j,s)}^{(Z,Q)} \right)^2 & \leq \lambda_{\min} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) - \left(M_{(k,v),(k,v)}^{(Z,Q)} \right)^2 \\ & \leq \sum_{\substack{(j,s)=1 \\ (j,s) \neq (k,v)}}^{m_n} \left(M_{(k,v),(j,s)}^{(Z,Q)} \right)^2 \end{aligned}$$

Using the first inequality above, we have

$$\begin{aligned} \lambda_{\min} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right) & \geq \left(M_{(k,v),(k,v)}^{(Z,Q)} \right)^2 - \sum_{\substack{(j,s)=1 \\ (j,s) \neq (k,v)}}^{m_n} \left(M_{(k,v),(j,s)}^{(Z,Q)} \right)^2 \\ & = 2 \left(M_{(k,v),(k,v)}^{(Z,Q)} \right)^2 - \sum_{(j,s)=1}^{m_n} \left(M_{(k,v),(j,s)}^{(Z,Q)} \right)^2 \\ & = 2 \left(M_{(k,v),(k,v)}^{(Z,Q)} \right)^2 - M_{(k,v),(k,v)}^{(Z,Q)} \\ & = 2 M_{(k,v),(k,v)}^{(Z,Q)} \left(M_{(k,v),(k,v)}^{(Z,Q)} - \frac{1}{2} \right) \\ & \geq 2 \min_{1 \leq (k,v) \leq m_n} \left[M_{(k,v),(k,v)}^{(Z,Q)} \left(M_{(k,v),(k,v)}^{(Z,Q)} - \frac{1}{2} \right) \right]. \end{aligned}$$

Hence, a sufficient condition that $\lambda_{\min}(M^{(Z,Q)} \circ M^{(Z,Q)})$ is bounded away from zero *a.s.n.* is that, there exists a positive constant \underline{C} such that

$$\min_{1 \leq (k,v) \leq m_n} \left[M_{(k,v),(k,v)}^{(Z,Q)} \left(M_{(k,v),(k,v)}^{(Z,Q)} - \frac{1}{2} \right) \right] \geq \underline{C} > 0 \text{ a.s.n.} \quad (50)$$

Now, consider the function

$$f(x) = x \left(x - \frac{1}{2} \right) \quad \text{for } 0 \leq x \leq 1$$

and note that

$$f'(x) = 2x - \frac{1}{2} > 0 \text{ for } \frac{1}{4} < x \leq 1,$$

from which we deduce that a sufficient condition for the condition given by (50) is

$$\min_{1 \leq (k,v) \leq m_n} M_{(k,v),(k,v)}^{(Z,Q)} \geq \frac{1}{2} + \epsilon_1 \text{ a.s.n.}$$

for some $\epsilon_1 > 0$. In addition, write $M^{(Z,Q)} = M^Q - P^{Z^\perp}$, where $P^{Z^\perp} = M^Q Z (Z' M^Q Z)^{-1} Z' M^Q$ and where the $(k, v)^{\text{th}}$ diagonal element of $M^Q = I_{m_n} - Q(Q'Q)^{-1}Q'$ is given by

$$M_{(k,v),(k,v)}^Q = 1 - \frac{1}{T_k}$$

Under Assumption 5, $P_{(k,v),(k,v)}^{Z^\perp} = O_{a.s.}(K_n/n) = o_{a.s.}(1)$ for every $(k, v) \in \{1, \dots, m_n\}$, so that for any $0 < \epsilon_1 < 1/6$, there exists a positive integer N_{ϵ_1} such that for all $n \geq N_{\epsilon_1}$, $P_{(k,v),(k,v)}^{Z^\perp} < \frac{1}{6} - \epsilon_1$ *a.s.* It follows that, under the assumption that $\min_{1 \leq k \leq n} T_k \geq 3$, we have for all $n \geq N_{\epsilon_2}$

$$\begin{aligned} \min_{1 \leq (k,v) \leq m_n} M_{(k,v),(k,v)}^{(Z,Q)} &= \min_{1 \leq (k,v) \leq m_n} \left(M_{(k,v),(k,v)}^Q - P_{(k,v),(k,v)}^{Z^\perp} \right) \\ &= \min_{1 \leq (k,v) \leq m_n} \left(1 - \frac{1}{T_k} - P_{(k,v),(k,v)}^{Z^\perp} \right) \\ &> \frac{2}{3} - \left(\frac{1}{6} - \epsilon_1 \right) = \frac{1}{2} + \epsilon_1 \text{ a.s.,} \end{aligned}$$

as required. \square

Section 3: Statement and Proof of Additional Lemmas

In this section, we state and prove a number of additional supporting lemmas whose results are used to prove some of the lemmas given in section 1 of this online appendix.

Lemma OA-1: Under Assumptions 5 and 6(i), the following statements are true.

(a)

$$\text{tr} \left\{ D_{\hat{\vartheta}}^2 \right\} = O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right),$$

where $D_{\widehat{\vartheta}} = \text{diag}(\widehat{\vartheta}_1, \dots, \widehat{\vartheta}_{m_n})$.

(b)

$$\max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 = O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right).$$

Proof of Lemma OA-1:

To show part (a), note that, by the result of Lemma 1, there exists a constant $C > 0$ such that

$$\begin{aligned} \text{tr} \left\{ D_{\widehat{\vartheta}}^2 \right\} &= \sum_{(i,t)=1}^{m_n} \widehat{\vartheta}_{(i,t)}^2 \\ &= d'_{P^\perp} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} \sum_{(i,t)=1}^{m_n} e_{(i,t)} e'_{(i,t)} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} d_{P^\perp} \\ &= d'_{P^\perp} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-2} d_{P^\perp} \\ &\leq \frac{1}{[\lambda_{\min}(M^{(Z,Q)} \circ M^{(Z,Q)})]^2} d'_{P^\perp} d_{P^\perp} \\ &\leq \left(\frac{1}{C} \right)^2 d'_{P^\perp} d_{P^\perp} \quad a.s. \\ &= \left(\frac{1}{C} \right)^2 \sum_{(i,t)=1}^{m_n} \left(P_{(i,t),(i,t)}^\perp \right)^2 \\ &\leq \left(\frac{1}{C} \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right) \sum_{(i,t)=1}^{m_n} P_{(i,t),(i,t)}^\perp \\ &= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) \quad (\text{by Assumption 5(iv)}). \end{aligned}$$

where $e_{(i,t)}$ denotes an $m_n \times 1$ elementary vector whose $(i,t)^{th}$ component is 1 and all other components are 0.

Next, we consider part (b). Note first that, as shown in the proof of Lemma 1, under the assumptions that $\min_{1 \leq i \leq n} T_i \geq 3$ and $K_n/n = o(1)$, the projection matrix $M^{(Z,Q)}$ is strictly diagonally dominate *a.s.n.*, so that there exists a positive constant \underline{C} such that

$$\min_{1 \leq (i,t) \leq m_n} \left[M_{(i,t),(i,t)}^{(Z,Q)} \left(M_{(i,t),(i,t)}^{(Z,Q)} - \frac{1}{2} \right) \right] \geq \underline{C} > 0 \quad a.s.n.$$

See, in particular, expression (50) in the proof of Lemma 1. Under these conditions, then, application of Theorem 1 of Varah (1975) to our problem leads to the following inequality

$$\left\| \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} \right\|_\infty \leq \left(\min_{1 \leq (i,t) \leq m_n} \left[\left(M_{(i,t),(i,t)}^{(Z,Q)} \right)^2 - \sum_{\substack{(k,v)=1 \\ (k,v) \neq (i,t)}}^{m_n} \left(M_{(i,t),(k,v)}^{(Z,Q)} \right)^2 \right] \right)^{-1}.$$

Now, making use of this inequality, we have, almost surely for all n sufficiently large,

$$\begin{aligned}
& \max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \\
&= \max_{1 \leq (i,t) \leq m_n} \left| e'_{(i,t)} \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} d_{P^\perp} \right|^2 \\
&\leq \max_{1 \leq (i,t) \leq m_n} \left\| \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} e_{(i,t)} \right\|_1^2 \|d_{P^\perp}\|_\infty^2 \quad (\text{by H\"older's inequality}) \\
&= \left(\max_{1 \leq (i,t) \leq m_n} \left\| \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} e_{(i,t)} \right\|_1 \right)^2 \|d_{P^\perp}\|_\infty^2 \\
&= \left(\max_{1 \leq (i,t) \leq m_n} \sum_{(j,s)=1}^{m_n} \left| \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1}_{(j,s),(i,t)} \right| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 \\
&= \left\| \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} \right\|_1^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 \\
&= \left\| \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1} \right\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 \quad (\text{by symmetry of } \left(M^{(Z,Q)} \circ M^{(Z,Q)} \right)^{-1}) \\
&\leq \left(\min_{1 \leq (i,t) \leq m_n} \left[\left(M_{(i,t),(i,t)}^{(Z,Q)} \right)^2 - \sum_{\substack{(k,v)=1 \\ (k,v) \neq (i,t)}}^{m_n} \left(M_{(i,t),(k,v)}^{(Z,Q)} \right)^2 \right] \right)^{-2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 \\
&\quad (\text{by Theorem 1 of Varah, 1975}) \\
&= \left(2 \min_{1 \leq (i,t) \leq m_n} \left[M_{(i,t),(i,t)}^{(Z,Q)} \left(M_{(i,t),(i,t)}^{(Z,Q)} - \frac{1}{2} \right) \right] \right)^{-2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 \\
&\leq \left(\frac{1}{2\underline{C}} \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right)^2 \quad (\text{by expression (50)}) \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right). \quad \square
\end{aligned}$$

Lemma OA-2:

Under Assumptions 1, 2, 3(iii), and 5(iii), the following statements are true.

(a)

$$\frac{\Upsilon' Z'_2 M^Q Z_1 (Z'_1 M^Q Z_1)^{-1} Z'_1 M^Q \varepsilon}{\sqrt{n}} = O_p(1),$$

(b)

$$\frac{\Upsilon' Z'_2 M^Q \varepsilon}{\sqrt{n}} = O_p(1).$$

(c)

$$\frac{\Upsilon' Z_2' M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} = O_p(1).$$

Proof of Lemma OA-2:

Note first that, under Assumption 5(iii), $Z' M^Q Z$ is positive definite *a.s.n.*, which implies that $Z_1' M^Q Z_1$ is positive definite and, thus, nonsingular *a.s.n.* as well. Now, to show part (a), note that, for any $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$, we obtain, by applying Assumptions 1, 2(i), and 3(iii),

$$\begin{aligned} & E \left[\left(\frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} \right)^2 | \mathcal{F}_n^Z \right] \\ &= \frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q E [\varepsilon \varepsilon' | \mathcal{F}_n^Z] M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q Z_2 \Upsilon a}{n} \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q Z_2 \Upsilon a}{n} \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n} \\ &= O_{a.s.}(1). \end{aligned}$$

Hence, there exists a positive constant $\bar{C} < \infty$ such that for all n sufficiently large

$$\begin{aligned} & E \left(\left[\frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} \right]^2 \right) \\ &= E_Z \left\{ E \left(\left[\frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} \right]^2 | \mathcal{F}_n^Z \right) \right\} \\ &\leq \bar{C} \end{aligned}$$

It follows from Markov's inequality that, for any $\epsilon > 0$, we can set $C_\epsilon = \sqrt{\bar{C}/\epsilon}$ so that, for all n sufficiently large,

$$\begin{aligned} & \Pr \left(\left| \frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} \right| \geq C_\epsilon \right) \\ &= \Pr \left(\left| \frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} \right|^2 \geq C_\epsilon^2 \right) \\ &\leq \frac{E \left(\left[a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon / \sqrt{n} \right]^2 \right)}{C_\epsilon^2} \\ &\leq \frac{\bar{C}}{\bar{C}/\epsilon} = \epsilon \end{aligned}$$

which shows that

$$\frac{a' \Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} = O_p(1)$$

Since the above result holds for all $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$, we further deduce that

$$\frac{\Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} = O_p(1),$$

as required for part (a).

Turning our attention to part (b), we again let $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$; and note that, by Assumptions 1, 2(i), and 3(iii),

$$\begin{aligned} E \left(\left[\frac{a' \Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} \right]^2 | \mathcal{F}_n^Z \right) &= \frac{a' \Upsilon' Z_2' M^Q E [\varepsilon \varepsilon' | \mathcal{F}_n^Z] M^Q Z_2 \Upsilon a}{n} \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \frac{a' \Upsilon' Z_2' M^Q Z_2 \Upsilon a}{n} \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n} \\ &= O_{a.s.}(1) \end{aligned}$$

Hence, there exists a positive constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left(\left[\frac{a' \Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} \right]^2 \right) = E_Z \left\{ E \left(\left[\frac{a' \Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} \right]^2 | \mathcal{F}_n^Z \right) \right\} \leq \bar{C}.$$

It follows from Markov's inequality, for any $\epsilon > 0$, we can set $C_\epsilon = \sqrt{\bar{C}/\epsilon}$ so that for all n sufficiently large

$$\begin{aligned} \Pr \left(\left| \frac{a' \Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} \right| \geq C_\epsilon \right) &= \Pr \left(\left| \frac{a' \Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} \right|^2 \geq C_\epsilon^2 \right) \\ &\leq \frac{E \left([a' \Upsilon' Z_2' M^Q \varepsilon / \sqrt{n}]^2 \right)}{C_\epsilon^2} \\ &\leq \frac{\bar{C}}{\bar{C}/\epsilon} = \epsilon \end{aligned}$$

which shows that

$$\frac{a' \Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} = O_p(1).$$

Since the above result holds for all $a \in \mathbb{R}^d$ such that $\|a\|_2 = 1$, we further deduce that

$$\frac{\Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} = O_p(1).$$

Finally, for part (c), note that it follows immediately from parts (a) and (b) above that

$$\frac{\Upsilon' Z_2' M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} = \frac{\Upsilon' Z_2' M^Q \varepsilon}{\sqrt{n}} - \frac{\Upsilon' Z_2' M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q \varepsilon}{\sqrt{n}} = O_p(1),$$

as required. \square

Lemma OA-3:

If Assumptions 1, 2(i), 4(ii), 5, and 6(i) are satisfied; then,

$$\frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} = O_p(1).$$

Proof of Lemma OA-3:

To proceed, note that, by the symmetry of A and by Assumptions 1, 2(i), and 5; we have

$$\begin{aligned} & E \left(\left[\frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} \right]^2 \mid \mathcal{F}_n^Z \right) \\ &= \frac{1}{K_{2,n}} \sum_{\substack{(i_1, t_1), (j_1, s_1) = 1 \\ (i_1, t_1) \neq (j_1, s_1)}}^{m_n} \sum_{\substack{(i_2, t_2), (j_2, s_2) = 1 \\ (i_2, t_2) \neq (j_2, s_2)}}^{m_n} A_{(i_1, t_1), (j_1, s_1)} A_{(i_2, t_2), (j_2, s_2)} E \left[\varepsilon_{(i_1, t_1)} \varepsilon_{(j_1, s_1)} \varepsilon_{(i_2, t_2)} \varepsilon_{(j_2, s_2)} \mid \mathcal{F}_n^Z \right] \\ &= \frac{2}{K_{2,n}} \sum_{\substack{(i, t), (j, s) = 1 \\ (i, t) \neq (j, s)}}^{m_n} A_{(i, t), (j, s)}^2 E \left[\varepsilon_{(i, t)}^2 \varepsilon_{(j, s)}^2 \mid \mathcal{F}_n^Z \right] \\ &\leq 2 \left(\max_{1 \leq (i, t) \leq m_n} E \left[\varepsilon_{(i, t)}^4 \mid \mathcal{F}_n^Z \right] \right) \frac{1}{K_{2,n}} \sum_{\substack{(i, t), (j, s) = 1 \\ (i, t) \neq (j, s)}}^{m_n} A_{(i, t), (j, s)}^2 \\ &= O_{a.s.}(1) \end{aligned}$$

Hence, there exists a positive constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left(\left[\frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} \right]^2 \right) = E_Z \left\{ E \left(\left[\frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} \right]^2 \mid \mathcal{F}_n^Z \right) \right\} \leq \bar{C}.$$

It follows from Markov's inequality, for any $\epsilon > 0$, we can set $C_\epsilon = \sqrt{\bar{C}/\epsilon}$ so that for all n sufficiently

large

$$\begin{aligned}
\Pr \left(\left| \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} \right| \geq C_\epsilon \right) &= \Pr \left(\left| \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} \right|^2 \geq C_\epsilon^2 \right) \\
&\leq \frac{E \left([\varepsilon' A \varepsilon / \sqrt{K_{2,n}}]^2 \right)}{C_\epsilon^2} \\
&\leq \frac{\bar{C}}{\bar{C}/\epsilon} = \epsilon
\end{aligned}$$

which shows that

$$\frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} = O_p(1). \quad \square$$

Lemma OA-4: Under Assumptions 1, 2, 6, and 8; the following statements are true.

(a)

$$\frac{\varepsilon' M^{Q_\varepsilon}}{n} - E \left[\frac{\varepsilon' M^{Q_\varepsilon}}{n} \right] = O_p \left(\frac{1}{\sqrt{n}} \right),$$

and there exist positive constants $\underline{C} \leq \bar{C}$ such that

$$0 < \underline{C} \leq E \left[\frac{\varepsilon' M^{Q_\varepsilon}}{n} \right] \leq \bar{C} < \infty$$

for all n sufficiently large.

(b)

$$\frac{U' M^{(Z_1, Q)_\varepsilon}}{n} - E \left[\frac{U' M^{(Z_1, Q)_\varepsilon}}{n} \right] = O_p \left(\max \left\{ \frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n} \right\} \right),$$

where

$$E \left[\frac{U' M^{Q_\varepsilon}}{n} \right] = O(1).$$

(c)

$$\frac{U' M^{(Z_1, Q)_\varepsilon}}{\varepsilon' M^{Q_\varepsilon}} - \rho = O_p \left(\max \left\{ \frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n} \right\} \right),$$

where

$$\rho = \lim_{n \rightarrow \infty} \rho_n = \lim_{n \rightarrow \infty} \frac{E [U' M^{Q_\varepsilon}] / n}{E [\varepsilon' M^{Q_\varepsilon}] / n}.$$

Proof of Lemma OA-4: To show part (a), note first that, making use of Assumptions 1, 2(i),

and 6(ii); we have

$$\begin{aligned}
& E \left(\frac{\varepsilon' M^{Q_\varepsilon} - E[\varepsilon' M^{Q_\varepsilon}]}{n} \right)^2 \\
= & \frac{E \left(\sum_{i=1}^n \sum_{t=1}^{T_i} \left(\frac{T_i-1}{T_i} \right) [\varepsilon_{(i,t)}^2 - E\{\varepsilon_{(i,t)}^2\}] - \sum_{i=1}^n \frac{1}{T_i} \sum_{r=1}^{T_i} \sum_{\substack{s=1 \\ s \neq r}} \varepsilon_{(i,r)} \varepsilon_{(i,s)} \right)^2}{n^2} \\
= & \frac{1}{n^2} \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{j=1}^n \sum_{s=1}^{T_j} \left(\frac{T_i-1}{T_i} \right) \left(\frac{T_j-1}{T_j} \right) E \left([\varepsilon_{(i,t)}^2 - E\{\varepsilon_{(i,t)}^2\}] [\varepsilon_{(j,s)}^2 - E\{\varepsilon_{(j,s)}^2\}] \right) \\
& - \frac{2}{n^2} \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{j=1}^n \sum_{r=1}^{T_j} \sum_{\substack{s=1 \\ s \neq r}} \left(\frac{T_i-1}{T_i} \right) \frac{1}{T_j} E \left\{ [\varepsilon_{(i,t)}^2 - E\{\varepsilon_{(i,t)}^2\}] \varepsilon_{(j,r)} \varepsilon_{(j,s)} \right\} \\
& + \frac{1}{n^2} \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{\substack{s=1 \\ s \neq t}} \sum_{j=1}^n \sum_{r=1}^{T_j} \sum_{\substack{h=1 \\ h \neq r}} \left(\frac{1}{T_i} \right) \left(\frac{1}{T_j} \right) E [\varepsilon_{(i,t)} \varepsilon_{(i,s)} \varepsilon_{(j,r)} \varepsilon_{(j,h)}] \\
= & \frac{1}{n^2} \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{j=1}^n \sum_{s=1}^{T_j} \left(\frac{T_i-1}{T_i} \right) \left(\frac{T_j-1}{T_j} \right) E \left([\varepsilon_{(i,t)}^2 - E\{\varepsilon_{(i,t)}^2\}] [\varepsilon_{(j,s)}^2 - E\{\varepsilon_{(j,s)}^2\}] \right) \\
& + \frac{2}{n^2} \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{\substack{s=1 \\ s \neq t}} \left(\frac{1}{T_i} \right)^2 E [\varepsilon_{(i,t)}^2] E [\varepsilon_{(i,s)}^2] \\
= & \frac{1}{n^2} \sum_{i=1}^n \sum_{t=1}^{T_i} \left(\frac{T_i-1}{T_i} \right)^2 E \left(\varepsilon_{(i,t)}^2 - E\{\varepsilon_{(i,t)}^2\} \right)^2 + \frac{2}{n^2} \sum_{i=1}^n \sum_{t=1}^{T_i} \sum_{\substack{s=1 \\ s \neq t}} \left(\frac{1}{T_i} \right)^2 E [\varepsilon_{(i,t)}^2] E [\varepsilon_{(i,s)}^2] \\
\leq & \frac{1}{n^2} n \bar{T} \max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4] + \frac{2}{n^2} n \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2] \right)^2 \\
= & \frac{1}{n} \bar{T} \max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4] + \frac{2}{n} \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2] \right)^2 \\
= & O \left(\frac{1}{n} \right)
\end{aligned}$$

It follows by Markov's inequality that

$$\frac{\varepsilon' M^{Q_\varepsilon} - E[\varepsilon' M^{Q_\varepsilon}]}{n} = O_p \left(\frac{1}{\sqrt{n}} \right). \quad (51)$$

Next, note that, by Assumption 6(i), $T_i \geq 3$ for all i , so that $\frac{T_i-1}{T_i} \geq \frac{2}{3}$ for all i . Hence, by

Assumption 2(ii), there exists a positive constant \underline{C} such that $E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \geq \underline{C} > 0$ a.s., so that

$$\begin{aligned}
\frac{E[\varepsilon' M^Q \varepsilon]}{n} &= \frac{1}{n} E_Z \left\{ \sum_{i=1}^n \sum_{t=1}^{T_i} \left[\left(\frac{T_i - 1}{T_i} \right) \right] E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right\} \\
&\geq \frac{2}{3n} E_Z \left\{ \sum_{i=1}^n \sum_{t=1}^{T_i} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right\} \\
&\geq \frac{2}{3n} E_Z \left[\sum_{i=1}^n \sum_{t=1}^{T_i} \underline{C} \right] \\
&= \frac{2}{3n} E_Z \left[\sum_{(i,t)=1}^{m_n} \underline{C} \right] \\
&= \frac{2}{3} \frac{m_n}{n} \underline{C} \\
&\geq \frac{2}{3} \underline{C} > 0
\end{aligned}$$

for all n sufficiently large. Furthermore, by Assumption 2(i), there also exists a positive constant C ($\geq \underline{C}$) such that

$$\left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \leq C \text{ a.s.},$$

from which it follows that

$$\begin{aligned}
E\left[\frac{\varepsilon' M^Q \varepsilon}{n}\right] &= \frac{1}{n} E_Z \left\{ \sum_{i=1}^n \sum_{t=1}^{T_i} \left(\frac{T_i - 1}{T_i} \right) E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right\} \\
&\leq \frac{1}{n} E_Z \left\{ \sum_{i=1}^n \sum_{t=1}^{T_i} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right\} \\
&\leq \frac{m_n}{n} C = \bar{C} < \infty.
\end{aligned}$$

Next, to show part (b), let $b \in \mathbb{R}^d$ such that $\|b\|_2 = 1$; and since $M^{(Z_1, Q)} = M^Q - P^{Z_1^\perp}$, where $P^{Z_1^\perp} = M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q$, we can write

$$\frac{b' U' M^{(Z_1, Q)} \varepsilon}{n} = \frac{b' U' M^Q \varepsilon}{n} - \frac{b' U' P^{Z_1^\perp} \varepsilon}{n}.$$

Note first that by argument similar to that given in part (a) above, we can show that

$$\frac{b' U' M^Q \varepsilon}{n} - E\left[\frac{b' U' M^Q \varepsilon}{n}\right] = O_p\left(\frac{1}{\sqrt{n}}\right). \quad (52)$$

Next, note that, by application of the CS inequality and Assumptions 1, 2(i), and 5; we have

$$\begin{aligned}
& E \left[\left| \frac{b' U' P^{Z_1^\perp} \varepsilon}{n} \right| \middle| \mathcal{F}_n^Z \right] \\
& \leq \sqrt{E \left[\frac{b' U' P^{Z_1^\perp} U b}{n} \middle| \mathcal{F}_n^Z \right]} \sqrt{E \left[\frac{\varepsilon' P^{Z_1^\perp} \varepsilon}{n} \middle| \mathcal{F}_n^Z \right]} \\
& = \sqrt{\frac{1}{n} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} P_{(i,t),(j,s)}^{Z_1^\perp} b' E \left[U_{(i,t)} U'_{(j,s)} \middle| \mathcal{F}_n^Z \right] b} \sqrt{\frac{1}{n} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} P_{(i,t),(j,s)}^{Z_1^\perp} E \left[\varepsilon_{(i,t)} \varepsilon_{(j,s)} \middle| \mathcal{F}_n^Z \right]} \\
& = \sqrt{\frac{1}{n} \sum_{(i,t)=1}^{m_n} P_{(i,t),(i,t)}^{Z_1^\perp} b' E \left[U_{(i,t)} U'_{(i,t)} \middle| \mathcal{F}_n^Z \right] b} \sqrt{\frac{1}{n} \sum_{(i,t)=1}^{m_n} P_{(i,t),(i,t)}^{Z_1^\perp} E \left[\varepsilon_{(i,t)}^2 \middle| \mathcal{F}_n^Z \right]} \\
& \leq \sqrt{\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 \middle| \mathcal{F}_n^Z \right]} \sqrt{\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 \middle| \mathcal{F}_n^Z \right]} \frac{K_{1,n}}{n} = O_{a.s.} \left(\frac{K_{1,n}}{n} \right)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left[\frac{n}{K_{1,n}} \left| \frac{b' U' P^{Z_1^\perp} \varepsilon}{n} \right| \right] = E_Z \left(E \left[\frac{n}{K_{1,n}} \left| \frac{b' U' P^{Z_1^\perp} \varepsilon}{n} \right| \middle| \mathcal{F}_n^Z \right] \right) \leq \bar{C}.$$

Application of Markov's inequality then implies that for any $\epsilon > 0$,

$$\Pr \left(\frac{n}{K_{1,n}} \left| \frac{b' U' P^{Z_1^\perp} \varepsilon}{n} \right| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{n}{K_{1,n}} \frac{E \left[\left| b' U' P^{Z_1^\perp} \varepsilon / n \right| \right]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\frac{b' U' P^{Z_1^\perp} \varepsilon}{n} = O_p \left(\frac{K_{1,n}}{n} \right). \quad (53)$$

It follows from (52) and (53) that

$$\frac{b' U' M^{(Z_1, Q)} \varepsilon}{n} - E \left[\frac{b' U' M^Q \varepsilon}{n} \right] = O_p \left(\max \left\{ \frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n} \right\} \right)$$

In addition, by the CS inequality and Assumptions 2(i) and 6(ii),

$$\begin{aligned}
\left| \frac{E[b'U'M^{Q_\varepsilon}]}{n} \right| &= \frac{1}{n} \left| \sum_{i=1}^n \sum_{t=1}^{T_i} \left(\frac{T_i - 1}{T_i} \right) E[b'U_{(i,t)}\varepsilon_{(i,t)}] \right| \\
&\leq \frac{1}{n} \sum_{i=1}^n \sum_{t=1}^{T_i} E[|b'U_{(i,t)}\varepsilon_{(i,t)}|] \\
&\leq \frac{1}{n} \sum_{i=1}^n \sum_{t=1}^{T_i} \sqrt{b'E[U_{(i,t)}U'_{(i,t)}]} b \sqrt{E[\varepsilon_{(i,t)}^2]} \\
&\leq \bar{T} \sqrt{\max_{1 \leq (i,t) \leq m_n} E\|U_{(i,t)}\|^2} \sqrt{\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^2]} \\
&= O(1)
\end{aligned}$$

Since the above argument holds for any $b \in \mathbb{R}^d$ such that $\|b\|_2 = 1$, we further deduce that

$$\frac{U'M^{(Z_1,Q)_\varepsilon}}{n} - E\left[\frac{U'M^{Q_\varepsilon}}{n}\right] = O_p\left(\max\left\{\frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n}\right\}\right).$$

and

$$E\left[\frac{U'M^{Q_\varepsilon}}{n}\right] = O(1),$$

as required to show part (b).

Finally, to show part (c), we first write

$$\frac{U'M^{(Z_1,Q)_\varepsilon}}{\varepsilon'M^{Q_\varepsilon}} - \rho = \frac{U'M^{(Z_1,Q)_\varepsilon}}{\varepsilon'M^{Q_\varepsilon}} - \rho_n + \rho_n - \rho$$

By Assumption 8, the sequence $\{\rho_n\}$ has a limit defined to be ρ so that $\rho_n - \rho \rightarrow 0$ as $n \rightarrow \infty$. Next, note that

$$\begin{aligned}
&\frac{U'M^{(Z_1,Q)_\varepsilon}}{\varepsilon'M^{Q_\varepsilon}} - \rho_n \\
&= \frac{U'M^{(Z_1,Q)_\varepsilon}}{\varepsilon'M^{Q_\varepsilon}} - \frac{E[U'M^{Q_\varepsilon}]}{E[\varepsilon'M^{Q_\varepsilon}]} \\
&= \frac{U'M^{(Z_1,Q)_\varepsilon}}{E[\varepsilon'M^{Q_\varepsilon}]} - \frac{E[U'M^{Q_\varepsilon}]}{E[\varepsilon'M^{Q_\varepsilon}]} + \frac{U'M^{(Z_1,Q)_\varepsilon}}{\varepsilon'M^{Q_\varepsilon}} - \frac{U'M^{(Z_1,Q)_\varepsilon}}{E[\varepsilon'M^{Q_\varepsilon}]} \\
&= \frac{U'M^{(Z_1,Q)_\varepsilon} - E[U'M^{Q_\varepsilon}]}{E[\varepsilon'M^{Q_\varepsilon}]} + \frac{U'M^{(Z_1,Q)_\varepsilon} E[\varepsilon'M^{Q_\varepsilon}]}{(\varepsilon'M^{Q_\varepsilon}) E[\varepsilon'M^{Q_\varepsilon}]} - \frac{(\varepsilon'M^{Q_\varepsilon}) U'M^{(Z_1,Q)_\varepsilon}}{(\varepsilon'M^{Q_\varepsilon}) E[\varepsilon'M^{Q_\varepsilon}]} \\
&= \frac{U'M^{(Z_1,Q)_\varepsilon} - E[U'M^{Q_\varepsilon}]}{E[\varepsilon'M^{Q_\varepsilon}]} - \frac{U'M^{(Z_1,Q)_\varepsilon} (\varepsilon'M^{Q_\varepsilon} - E[\varepsilon'M^{Q_\varepsilon}])}{\varepsilon'M^{Q_\varepsilon} E[\varepsilon'M^{Q_\varepsilon}]}
\end{aligned}$$

Now, applying the results of parts (a) and (b) above and the Slutsky's theorem, we obtain

$$\begin{aligned}\frac{U'M^{(Z_1,Q)}\varepsilon - E[U'M^Q\varepsilon]}{E[\varepsilon'M^Q\varepsilon]} &= \frac{(U'M^{(Z_1,Q)}\varepsilon - E[U'M^Q\varepsilon])/n}{E[\varepsilon'M^Q\varepsilon]/n} = O_p\left(\max\left\{\frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n}\right\}\right), \\ \frac{\varepsilon'M^Q\varepsilon - E[\varepsilon'M^Q\varepsilon]}{E[\varepsilon'M^Q\varepsilon]} &= \frac{(\varepsilon'M^Q\varepsilon - E[\varepsilon'M^Q\varepsilon])/n}{E[\varepsilon'M^Q\varepsilon]/n} = O_p\left(\frac{1}{\sqrt{n}}\right),\end{aligned}$$

and

$$\begin{aligned}\frac{U'M^{(Z_1,Q)}\varepsilon}{\varepsilon'M^Q\varepsilon} &= \frac{U'M^{(Z_1,Q)}\varepsilon/n}{\varepsilon'M^Q\varepsilon/n} \\ &= \frac{(U'M^{(Z_1,Q)}\varepsilon - E[U'M^Q\varepsilon])/n + E[U'M^Q\varepsilon]/n}{(\varepsilon'M^Q\varepsilon - E[\varepsilon'M^Q\varepsilon])/n + E[\varepsilon'M^Q\varepsilon]/n} \\ &= \frac{E[U'M^Q\varepsilon]/n + o_p(1)}{E[\varepsilon'M^Q\varepsilon]/n + o_p(1)} = O_p(1).\end{aligned}$$

It follows that

$$\begin{aligned}&\frac{U'M^{(Z_1,Q)}\varepsilon}{\varepsilon'M^Q\varepsilon} - \rho_n \\ &= \frac{U'M^{(Z_1,Q)}\varepsilon}{\varepsilon'M^Q\varepsilon} - \frac{E[U'M^Q\varepsilon]}{E[\varepsilon'M^Q\varepsilon]} \\ &= \frac{(U'M^{(Z_1,Q)}\varepsilon - E[U'M^Q\varepsilon])/n}{E[\varepsilon'M^Q\varepsilon]/n} - \frac{U'M^{(Z_1,Q)}\varepsilon - E[\varepsilon'M^Q\varepsilon]/n}{\varepsilon'M^Q\varepsilon} \\ &= O_p\left(\max\left\{\frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n}\right\}\right) - O_p(1)O_p\left(\frac{1}{\sqrt{n}}\right) \\ &= O_p\left(\max\left\{\frac{1}{\sqrt{n}}, \frac{K_{1,n}}{n}\right\}\right). \square\end{aligned}$$

Lemma OA-5:

Suppose that Assumptions 1-6 are satisfied. Let

$$\hat{\ell}(\delta_0) = \frac{(y - X\delta_0)' A (y - X\delta_0)}{(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)};$$

then,

$$\hat{\ell}(\delta_0) = \frac{\varepsilon' A \varepsilon}{\varepsilon' M^Q \varepsilon} \left[1 + O_p\left(\max\left\{\frac{\tau_n}{\sqrt{K_{2,n} K_{1,n}^{\varrho_g}}}, \frac{K_{1,n}}{n}\right\}\right)\right],$$

Proof of Lemma OA-5:

To proceed, first write

$$\hat{\ell}(\delta_0) = \frac{(y - X\delta_0)' A (y - X\delta_0)}{(y - X\delta_0)' M^{(Z_1,Q)} (y - X\delta_0)} = \frac{\varepsilon' A \varepsilon}{\varepsilon' M^{(Z_1,Q)} \varepsilon}$$

Moreover, note that $\varepsilon' M^{(Z_1, Q)} \varepsilon / n = \varepsilon' M^Q \varepsilon / n - \varepsilon' P^{Z_1^\perp} \varepsilon / n$. By the results of part (a) of Lemma OA-4, we have that $\varepsilon' M^Q \varepsilon / n - E[\varepsilon' M^Q \varepsilon / n] = O_p(n^{-1/2})$, and also that there exist positive constants $\underline{C} \leq \bar{C}$ such that, for all n sufficiently large, $0 < \underline{C} \leq E[\varepsilon' M^Q \varepsilon / n] \leq \bar{C} < \infty$. In addition, by argument similar to that used to obtain expression (53) in the proof of part (b) of Lemma OA-4, we can show that $\varepsilon' P^{Z_1^\perp} \varepsilon / n = O_p(K_{1,n}/n) = o_p(1)$. It follows that

$$\frac{(y - X\delta_0)' M^{(Z_1, Q)} (y - X\delta_0)}{n} = \frac{\varepsilon' M^{(Z_1, Q)} \varepsilon}{n} = \frac{\varepsilon' M^Q \varepsilon}{n} \left[1 + O_p\left(\frac{K_{1,n}}{n}\right) \right], \quad (54)$$

where $\varepsilon' M^Q \varepsilon / n = O_p(1)$ and where $\varepsilon' M^Q \varepsilon / n > 0$ w.p.a.1. Finally, note that

$$\begin{aligned} \hat{\ell}(\delta_0) &= \frac{\varepsilon' A \varepsilon}{n} \left[\frac{\varepsilon' M^{(Z_1, Q)} \varepsilon}{n} \right]^{-1} \\ &= \frac{\varepsilon' A \varepsilon}{n} \left[\frac{\varepsilon' M^Q \varepsilon}{n} \left(1 + O_p\left(\frac{K_{1,n}}{n}\right) \right) \right]^{-1} \\ &= \frac{\varepsilon' A \varepsilon}{\varepsilon' M^Q \varepsilon} \left[1 + O_p\left(\frac{K_{1,n}}{n}\right) \right]^{-1} \\ &= \frac{\varepsilon' A \varepsilon}{\varepsilon' M^Q \varepsilon} \left[1 + O_p\left(\frac{K_{1,n}}{n}\right) \right] \end{aligned}$$

which shows the required result. \square

Lemma OA-6: Suppose that Assumptions 1-6 are satisfied. Let $\hat{\delta}_n$ be any estimator that satisfies the following conditions as $n \rightarrow \infty$

(i) If $K_{2,n}/(\mu_n^{\min})^2 = O(1)$, then

$$D_\mu(\hat{\delta}_n - \delta_0) = O_p(1)$$

(ii) If $(\mu_n^{\min})^2/K_{2,n} = o(1)$ but $\sqrt{K_{2,n}}/(\mu_n^{\min})^2 \rightarrow 0$, then

$$\frac{(\mu_n^{\min}) D_\mu(\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} = O_p(1)$$

Under these conditions, the following statements are true.

(a) Under case (i),

$$D_\mu^{-1} X' A (y - X\hat{\delta}_n) = O_p(1)$$

while, under case (ii),

$$D_\mu^{-1} X' A (y - X\hat{\delta}_n) = O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right).$$

(b) Under both case (i) and case (ii),

$$D_\mu^{-1} X' M^{(Z_1, Q)} \left(y - X \hat{\delta}_n \right) = O_p \left(\frac{n}{(\mu_n^{\min})} \right)$$

(c) Under both case (i) and case (ii),

$$\frac{\left(y - X \hat{\delta}_n \right)' A \left(y - X \hat{\delta}_n \right)}{\sqrt{K_{2,n}}} = \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} + o_p(1),$$

where $\varepsilon' A \varepsilon / \sqrt{K_{2,n}} = O_p(1)$.

(d) Under both case (i) and case (ii),

$$\frac{\left(y - X \hat{\delta}_n \right)' M^{(Z_1, Q)} \left(y - X \hat{\delta}_n \right)}{n} = \frac{\varepsilon' M^Q \varepsilon}{n} [1 + o_p(1)],$$

where $\varepsilon' M^Q \varepsilon / n = O_p(1)$ and where $\varepsilon' M^Q \varepsilon / n > 0$ w.p.a.1.

(e) Under both case (i) and case (ii),

$$\hat{\ell} \left(\hat{\delta}_n \right) = \frac{\varepsilon' A \varepsilon}{\varepsilon' M^Q \varepsilon} + o_p \left(\frac{\sqrt{K_{2,n}}}{n} \right) = O_p \left(\frac{\sqrt{K_{2,n}}}{n} \right),$$

where

$$\hat{\ell} \left(\hat{\delta}_n \right) = \frac{\left(y - X \hat{\delta}_n \right)' A \left(y - X \hat{\delta}_n \right)}{\left(y - X \hat{\delta}_n \right)' M^{(Z_1, Q)} \left(y - X \hat{\delta}_n \right)}.$$

Proof of Lemma OA-6:

To show part (a), note that

$$\begin{aligned} D_\mu^{-1} X' A \left(y - X \hat{\delta}_n \right) &= D_\mu^{-1} X' A (y - X \delta_0) + D_\mu^{-1} X' A X \left(\hat{\delta}_n - \delta_0 \right) \\ &= D_\mu^{-1} X' A (y - X \delta_0) + D_\mu^{-1} X' A X D_\mu^{-1} D_\mu \left(\hat{\delta}_n - \delta_0 \right) \end{aligned} \quad (55)$$

To analyze the first term on the right-hand side of expression (55) above, write

$$D_\mu^{-1} X' A (y - X \delta_0) = \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} + D_\mu^{-1} U' A \varepsilon$$

Now, consider case (i) where $\sqrt{K_{2,n}} / (\mu_n^{\min}) = O(1)$. In this case, we have by part (c) of Lemma OA-2 and expression (??) that

$$\frac{\Upsilon' Z'_2 M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} = O_p(1) \text{ and } D_\mu^{-1} U' A \varepsilon = O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} \right),$$

from which it follows that, in this case, $D_\mu^{-1}X'A(y - X\delta_0) = O_p(1)$. In addition, by part (b) of Lemma S2-2, we have in this case

$$D_\mu^{-1}X'AXD_\mu^{-1}D_\mu(\hat{\delta}_n - \delta_0) = O_p(1),$$

so that, in this case,

$$\begin{aligned} D_\mu^{-1}X'A(y - X\hat{\delta}_n) &= D_\mu^{-1}X'A(y - X\delta_0) + D_\mu^{-1}X'AXD_\mu^{-1}D_\mu(\hat{\delta}_n - \delta_0) \\ &= O_p(1) + O_p(1) = O_p(1) \end{aligned}$$

Next, consider case (ii) where $(\mu_n^{\min})/\sqrt{K_{2,n}} = o(1)$ but $\sqrt{K_{2,n}}/(\mu_n^{\min})^2 \rightarrow 0$. Here, again by part (c) of Lemma OA-2, and expression (??), we have

$$D_\mu^{-1}X'A(y - X\delta_0) = O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right)$$

Furthermore, in this case, we also have, by using part (b) of Lemma S2-2

$$\begin{aligned} &D_\mu^{-1}X'AXD_\mu^{-1}D_\mu(\hat{\delta}_n - \delta_0) \\ &= \frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} D_\mu^{-1}X'AXD_\mu^{-1} \frac{\mu_n^{\min}D_\mu(\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} = O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right) \end{aligned}$$

from which it follows that in case (ii)

$$\begin{aligned} D_\mu^{-1}X'A(y - X\hat{\delta}_n) &= D_\mu^{-1}X'A(y - X\delta_0) + D_\mu^{-1}X'AXD_\mu^{-1}D_\mu(\hat{\delta}_n - \delta_0) \\ &= O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right) + O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right) \\ &= O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right). \end{aligned}$$

Next, consider part (b). In this case, note that

$$\begin{aligned} &D_\mu^{-1}X'M^{(Z_1,Q)}(y - X\hat{\delta}_n) \\ &= D_\mu^{-1}X'M^{(Z_1,Q)}(y - X\delta_0) + D_\mu^{-1}X'M^{(Z_1,Q)}X(\hat{\delta}_n - \delta_0) \\ &= D_\mu^{-1}X'M^{(Z_1,Q)}(y - X\delta_0) + D_\mu^{-1}X'M^{(Z_1,Q)}XD_\mu^{-1}D_\mu(\hat{\delta}_n - \delta_0) \end{aligned} \tag{56}$$

The first term on the right-hand side of the expression (56) above can be written as

$$D_\mu^{-1}X'M^{(Z_1,Q)}(y - X\delta_0) = D_\mu^{-1}X'M^{(Z_1,Q)}\varepsilon = \frac{\Upsilon'Z'_2M^{(Z_1,Q)}\varepsilon}{\sqrt{n}} + D_\mu^{-1}U'M^{(Z_1,Q)}\varepsilon$$

Now, consider case (i) where $\sqrt{K_{2,n}} / (\mu_n^{\min}) = O(1)$. In this case, we have by part (c) of Lemma OA-2 and expression (11) that

$$\frac{\Upsilon' Z_2' M^{(Z_1, Q)} \varepsilon}{\sqrt{n}} = O_p(1) \text{ and } D_\mu^{-1} U' M^{(Z_1, Q)} \varepsilon = O_p\left(\frac{n}{(\mu_n^{\min})}\right),$$

from which it follows that, in this case,

$$D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\delta_0) = O_p\left(\frac{n}{(\mu_n^{\min})}\right).$$

In addition, by part (a) of Lemma S2-2, we have in this case

$$D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1} D_\mu (\hat{\delta}_n - \delta_0) = O_p\left(\frac{n}{(\mu_n^{\min})^2}\right),$$

so that, in this case,

$$\begin{aligned} & D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\hat{\delta}_n) \\ = & D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\delta_0) + D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1} D_\mu (\hat{\delta}_n - \delta_0) \\ = & O_p\left(\frac{n}{(\mu_n^{\min})}\right) + O_p\left(\frac{n}{(\mu_n^{\min})^2}\right) = O_p\left(\frac{n}{(\mu_n^{\min})}\right) \end{aligned}$$

Next, consider case (ii) where $(\mu_n^{\min}) / \sqrt{K_{2,n}} = o(1)$ but $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Here, again by part (c) of Lemma OA-2 and expression (11), we have

$$D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\delta_0) = O_p\left(\frac{n}{(\mu_n^{\min})}\right)$$

Furthermore, using part (a) of Lemma S2-2, we have

$$\begin{aligned} D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1} D_\mu (\hat{\delta}_n - \delta_0) &= \frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1} \frac{(\mu_n^{\min}) D_\mu (\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\ &= O_p\left(\frac{\sqrt{K_{2,n}} n}{(\mu_n^{\min})^3}\right) \end{aligned}$$

from which it follows that in case (ii)

$$\begin{aligned} & D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\hat{\delta}_n) \\ = & D_\mu^{-1} X' M^{(Z_1, Q)} (y - X\delta_0) + D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1} D_\mu (\hat{\delta}_n - \delta_0) \\ = & O_p\left(\frac{n}{(\mu_n^{\min})}\right) + O_p\left(\frac{\sqrt{K_{2,n}} n}{(\mu_n^{\min})^3}\right) = O_p\left(\frac{n}{(\mu_n^{\min})}\right) \end{aligned}$$

Combining the results for cases (i) and (ii), we have

$$D_\mu^{-1} X' M^{(Z_1, Q)} \left(y - X\hat{\delta}_n \right) = O_p \left(\frac{n}{(\mu_n^{\min})} \right)$$

Consider now part (c). Write

$$\begin{aligned} & \frac{(y - X\hat{\delta}_n)' A (y - X\hat{\delta}_n)}{\sqrt{K_{2,n}}} \\ = & \frac{\left[y - X\delta_0 - X(\hat{\delta}_n - \delta_0) \right]' A \left[y - X\delta_0 - X(\hat{\delta}_n - \delta_0) \right]}{\sqrt{K_{2,n}}} \\ = & \frac{(y - X\delta_0)' A (y - X\delta_0)}{\sqrt{K_{2,n}}} - 2 \frac{(y - X\delta_0)' A X (\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\ & + \frac{(\hat{\delta}_n - \delta_0)' X' A X (\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\ = & \frac{(y - X\delta_0)' A (y - X\delta_0)}{\sqrt{K_{2,n}}} - 2 (y - X\delta_0)' A X D_\mu^{-1} \frac{D_\mu (\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\ & + (\hat{\delta}_n - \delta_0)' D_\mu \frac{D_\mu^{-1} X' A X D_\mu^{-1}}{\sqrt{K_{2,n}}} D_\mu (\hat{\delta}_n - \delta_0) \end{aligned}$$

Note first that

$$\begin{aligned} \frac{(y - X\delta_0)' A (y - X\delta_0)}{\sqrt{K_{2,n}}} &= \frac{(Z_1 \gamma \tau_n / \sqrt{n} + Q\alpha + \varepsilon)' A (Z_1 \gamma \tau_n / \sqrt{n} + Q\alpha + \varepsilon)}{\sqrt{K_{2,n}}} \\ &= \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}}. \end{aligned}$$

and

$$\begin{aligned} & (y - X\delta_0)' A X D_\mu^{-1} \\ = & (Z_1 \gamma \tau_n / \sqrt{n} + Q\alpha + \varepsilon)' A (Z_1 \Theta D_\kappa / \sqrt{n} + Z_2 \Upsilon D_\mu / \sqrt{n} + Q\Xi + U) D_\mu^{-1} \\ = & \frac{\varepsilon' M^{(Z_1, Q)} Z_2 \Upsilon}{\sqrt{n}} + \varepsilon' A U D_\mu^{-1} \end{aligned}$$

Now, consider case (i) where $\sqrt{K_{2,n}} / (\mu_n^{\min}) = O(1)$. In this case, we have by part (c) of Lemma OA-2 and expression (??) that

$$\frac{\varepsilon' M^{(Z_1, Q)} Z_2 \Upsilon}{\sqrt{n}} = O_p(1) \text{ and } \varepsilon' A U D_\mu^{-1} = O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} \right),$$

so that $(y - X\delta_0)' A X D_\mu^{-1} = O_p(1)$ and, thus,

$$(y - X\delta_0)' A X D_\mu^{-1} \frac{D_\mu(\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} = O_p\left(\frac{1}{\sqrt{K_{2,n}}}\right) = o_p(1)$$

In addition, by part (b) of Lemma S2-2, we have in this case

$$\begin{aligned} & (\hat{\delta}_n - \delta_0)' D_\mu \frac{D_\mu^{-1} X' A X D_\mu^{-1}}{\sqrt{K_{2,n}}} D_\mu(\hat{\delta}_n - \delta_0) \\ &= \frac{1}{\sqrt{K_{2,n}}} (\hat{\delta}_n - \delta_0)' D_\mu D_\mu^{-1} X' A X D_\mu^{-1} D_\mu(\hat{\delta}_n - \delta_0) \\ &= O_p\left(\frac{1}{\sqrt{K_{2,n}}}\right) = o_p(1), \end{aligned}$$

from which it follows that

$$\begin{aligned} \frac{(y - X\hat{\delta}_n)' A (y - X\hat{\delta}_n)}{\sqrt{K_{2,n}}} &= \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} + O_p\left(\frac{1}{\sqrt{K_{2,n}}}\right) + O_p\left(\frac{1}{\sqrt{K_{2,n}}}\right) \\ &= \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} + o_p(1). \end{aligned}$$

Next, consider case (ii) where $(\mu_n^{\min})/\sqrt{K_{2,n}} = o(1)$ but $\sqrt{K_{2,n}}/(\mu_n^{\min})^2 \rightarrow 0$. Here, again by part (c) of Lemma OA-2, and expression (??), we have

$$(y - X\delta_0)' A X D_\mu^{-1} = O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right)$$

so that

$$\begin{aligned} & (y - X\delta_0)' A X D_\mu^{-1} \frac{D_\mu(\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\ &= \frac{1}{(\mu_n^{\min})} (y - X\delta_0)' A X D_\mu^{-1} \frac{(\mu_n^{\min}) D_\mu(\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\ &= O_p\left(\frac{1}{(\mu_n^{\min})}\right) O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})}\right) O_p(1) = O_p\left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2}\right) = o_p(1). \end{aligned}$$

Furthermore, in this case,

$$\begin{aligned}
& \left(\hat{\delta}_n - \delta_0 \right)' D_\mu \frac{D_\mu^{-1} X' A X D_\mu^{-1}}{\sqrt{K_{2,n}}} D_\mu \left(\hat{\delta}_n - \delta_0 \right) \\
&= \frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \left[\frac{(\mu_n^{\min}) \left(\hat{\delta}_n - \delta_0 \right)' D_\mu}{\sqrt{K_{2,n}}} D_\mu^{-1} X' A X D_\mu^{-1} \frac{(\mu_n^{\min}) D_\mu \left(\hat{\delta}_n - \delta_0 \right)}{\sqrt{K_{2,n}}} \right] \\
&= O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) = o_p(1)
\end{aligned}$$

from which it follows that in case (ii)

$$\begin{aligned}
\frac{\left(y - X \hat{\delta}_n \right)' A \left(y - X \hat{\delta}_n \right)}{\sqrt{K_{2,n}}} &= \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} + O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) + O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) \\
&= \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} + o_p(1).
\end{aligned}$$

Combining the results for cases (i) and (ii), we have

$$\frac{\left(y - X \hat{\delta}_n \right)' A \left(y - X \hat{\delta}_n \right)}{\sqrt{K_{2,n}}} = \frac{\varepsilon' A \varepsilon}{\sqrt{K_{2,n}}} + o_p(1).$$

where $\varepsilon' A \varepsilon / \sqrt{K_{2,n}} = O_p(1)$ by Lemma OA-3.

Turning our attention to part (d), note that

$$\begin{aligned}
& \frac{\left(y - X \hat{\delta}_n \right)' M^{(Z_1, Q)} \left(y - X \hat{\delta}_n \right)}{n} \\
&= \frac{\left[y - X \delta_0 - X \left(\hat{\delta}_n - \delta_0 \right) \right]' M^{(Z_1, Q)} \left[y - X \delta_0 - X \left(\hat{\delta}_n - \delta_0 \right) \right]}{n} \\
&= \frac{(y - X \delta_0)' M^{(Z_1, Q)} (y - X \delta_0)}{n} - 2 \frac{(y - X \delta_0)' M^{(Z_1, Q)} X \left(\hat{\delta}_n - \delta_0 \right)}{n} \\
&\quad + \frac{\left(\hat{\delta}_n - \delta_0 \right)' X' M^{(Z_1, Q)} X \left(\hat{\delta}_n - \delta_0 \right)}{n} \\
&= \frac{(y - X \delta_0)' M^{(Z_1, Q)} (y - X \delta_0)}{n} - 2 \frac{(y - X \delta_0)' M^{(Z_1, Q)} X D_\mu^{-1} D_\mu \left(\hat{\delta}_n - \delta_0 \right)}{n} \\
&\quad + \left(\hat{\delta}_n - \delta_0 \right)' D_\mu \frac{D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1}}{n} D_\mu \left(\hat{\delta}_n - \delta_0 \right).
\end{aligned}$$

Making use of the argument used to derive expression (54) in the proof of Lemma OA-5, we see

that

$$\frac{(y - X\delta_0)' M^{(Z_1, Q)} (y - X\delta_0)}{n} = \frac{\varepsilon' M^{(Z_1, Q)} \varepsilon}{n} = \frac{\varepsilon' M^Q \varepsilon}{n} + O_p \left(\frac{K_{1,n}}{n} \right).$$

Moreover, we can write

$$\begin{aligned} & (y - X\delta_0)' M^{(Z_1, Q)} X D_\mu^{-1} \\ &= (Z_1 \gamma \tau_n / \sqrt{n} + Q\alpha + \varepsilon)' M^{(Z_1, Q)} (Z_1 \Theta D_\kappa / \sqrt{n} + Z_2 \Upsilon D_\mu / \sqrt{n} + Q\Xi + U) D_\mu^{-1} \\ &= \frac{\varepsilon' M^{(Z_1, Q)} Z_2 \Upsilon}{\sqrt{n}} + \varepsilon' M^{(Z_1, Q)} U D_\mu^{-1} \end{aligned}$$

Now, consider case (i) where $\sqrt{K_{2,n}} / (\mu_n^{\min}) = O(1)$. In this case, we have by part (c) of Lemma OA-2 and expression (11) that

$$\frac{\varepsilon' M^{(Z_1, Q)} Z_2 \Upsilon}{\sqrt{n}} = O_p(1) \text{ and } \varepsilon' M^{(Z_1, Q)} U D_\mu^{-1} = O_p \left(\frac{n}{(\mu_n^{\min})} \right),$$

so that $(y - X\delta_0)' M^{(Z_1, Q)} X D_\mu^{-1} = O_p(n / (\mu_n^{\min}))$ and, thus,

$$\begin{aligned} \frac{(y - X\delta_0)' M^{(Z_1, Q)} X D_\mu^{-1}}{n} D_\mu (\hat{\delta}_n - \delta_0) &= O_p \left(\frac{1}{(\mu_n^{\min})} \right) O_p(1) \\ &= O_p \left(\frac{1}{(\mu_n^{\min})} \right) \end{aligned}$$

In addition, by part (a) of Lemma S2-2, we have in this case

$$\begin{aligned} & (\hat{\delta}_n - \delta_0)' D_\mu \frac{D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1}}{n} D_\mu (\hat{\delta}_n - \delta_0) \\ &= O_p(1) O_p \left(\frac{1}{(\mu_n^{\min})^2} \right) O_p(1) = O_p \left(\frac{1}{(\mu_n^{\min})^2} \right) = o_p(1), \end{aligned}$$

from which it follows that

$$\begin{aligned} & \frac{(y - X\hat{\delta}_n)' M^{(Z_1, Q)} (y - X\hat{\delta}_n)}{n} \\ &= \frac{\varepsilon' M^Q \varepsilon}{n} + O_p \left(\frac{K_{1,n}}{n} \right) + O_p \left(\frac{1}{(\mu_n^{\min})} \right) + O_p \left(\frac{1}{(\mu_n^{\min})^2} \right) \\ &= \frac{\varepsilon' M^Q \varepsilon}{n} + o_p(1). \end{aligned}$$

Next, consider case (ii) where $(\mu_n^{\min}) / \sqrt{K_{2,n}} = o(1)$ but $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Here, again by part (c) of Lemma OA-2 and expression (11); we have

$$\frac{(y - X\delta_0)' M^{(Z_1, Q)} X D_\mu^{-1}}{n} = O_p \left(\frac{1}{(\mu_n^{\min})} \right) = o_p(1)$$

so that

$$\begin{aligned}
& \frac{(y - X\delta_0)' M^{(Z_1, Q)} X D_\mu^{-1}}{n} D_\mu (\hat{\delta}_n - \delta_0) \\
&= \frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} \frac{(y - X\delta_0)' M^{(Z_1, Q)} X D_\mu^{-1}}{n} \frac{(\mu_n^{\min}) D_\mu (\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\
&= O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})} \right) O_p \left(\frac{1}{(\mu_n^{\min})} \right) O_p (1) = O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) = o_p (1).
\end{aligned}$$

Furthermore, in this case,

$$\begin{aligned}
& (\hat{\delta}_n - \delta_0)' D_\mu \frac{D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1}}{n} D_\mu (\hat{\delta}_n - \delta_0) \\
&= \frac{K_{2,n}}{(\mu_n^{\min})^2} \frac{(\mu_n^{\min}) (\hat{\delta}_n - \delta_0)' D_\mu D_\mu^{-1} X' M^{(Z_1, Q)} X D_\mu^{-1}}{\sqrt{K_{2,n}}} \frac{(\mu_n^{\min}) D_\mu (\hat{\delta}_n - \delta_0)}{\sqrt{K_{2,n}}} \\
&= O_p \left(\frac{K_{2,n}}{(\mu_n^{\min})^2} \right) O_p (1) O_p \left(\frac{1}{(\mu_n^{\min})^2} \right) O_p (1) = O_p \left(\frac{K_{2,n}}{(\mu_n^{\min})^4} \right) = o_p (1)
\end{aligned}$$

from which it follows that in case (ii)

$$\begin{aligned}
\frac{(y - X\hat{\delta}_n)' M^{(Z_1, Q)} (y - X\hat{\delta}_n)}{n} &= \frac{\varepsilon' M^Q \varepsilon}{n} + O_p \left(\frac{K_{1,n}}{n} \right) + O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) + O_p \left(\frac{K_{2,n}}{(\mu_n^{\min})^4} \right) \\
&= \frac{\varepsilon' M^Q \varepsilon}{n} + o_p (1)
\end{aligned}$$

Combining the results for cases (i) and (ii), we have

$$\frac{(y - X\hat{\delta}_n)' M^{(Z_1, Q)} (y - X\hat{\delta}_n)}{n} = \frac{\varepsilon' M^Q \varepsilon}{n} + o_p (1).$$

where $\varepsilon' M^Q \varepsilon / n = O_p (1)$ and $\varepsilon' M^Q \varepsilon / n > 0$ w.p.a.1. were both shown in the proof of Lemma OA-5.

Finally, to show part (e), we apply the results of parts (c) and (d) above to obtain

$$\begin{aligned}
& \widehat{\ell}(\widehat{\delta}_n) \\
&= \frac{(y - X\widehat{\delta}_n)' A (y - X\widehat{\delta}_n)}{(y - X\widehat{\delta}_n)' M^{(Z_1, Q)} (y - X\widehat{\delta}_n)} \\
&= \frac{\sqrt{K_{2,n}}}{n} \frac{(y - X\widehat{\delta}_n)' A (y - X\widehat{\delta}_n) / \sqrt{K_{2,n}}}{(y - X\widehat{\delta}_n)' M^{(Z_1, Q)} (y - X\widehat{\delta}_n) / n} \\
&= \frac{\sqrt{K_{2,n}}}{n} \frac{\varepsilon' A \varepsilon / \sqrt{K_{2,n}} + o_p(1)}{\varepsilon' M^Q \varepsilon / n + o_p(1)} \\
&= \frac{\sqrt{K_{2,n}}}{n} \left[\frac{\varepsilon' A \varepsilon / \sqrt{K_{2,n}}}{\varepsilon' M^Q \varepsilon / n} + o_p(1) \right] \quad \left(\text{since } \frac{\varepsilon' M^Q \varepsilon}{n} > 0 \text{ w.p.a.1} \right) \\
&= \frac{\varepsilon' A \varepsilon}{\varepsilon' M^Q \varepsilon} + o_p\left(\frac{\sqrt{K_{2,n}}}{n}\right) = O_p\left(\frac{\sqrt{K_{2,n}}}{n}\right)
\end{aligned}$$

given that $\varepsilon' A \varepsilon / \sqrt{K_{2,n}} = O_p(1)$ and $\varepsilon' M^Q \varepsilon / n > 0$ w.p.a.1. \square

Lemma OA-7: (Decoupling Inequality) For natural number $n \geq G$, let $\{X_i\}_{i=1}^n$ be n independent random variables taking on values in a measurable space (S, \mathfrak{S}) , and let $\{X_i^{(k)}\}_{i=1}^n$ $k = 1, \dots, G$ be G independent copies of this sequence. Let B be a separable Banach space and for each $(i_1, \dots, i_G) \in I_n^{g*}$, where

$$I_n^{g*} = \{(i_1, \dots, i_G) : i_j \in \mathbb{N}, 1 \leq i_j \leq n, i_j \neq i_k \text{ if } j \neq k\},$$

let $h_{i_1 \dots i_G} : S^G \mapsto B$ be a measurable functions such that $E(\|h_{i_1 \dots i_G}(X_{i_1}, \dots, X_{i_G})\|) < \infty$. Let $\Phi : [0, \infty) \mapsto [0, \infty)$ be a convex nondecreasing function such that $E\Phi(\|h_{i_1 \dots i_G}(X_{i_1}, \dots, X_{i_G})\|) < \infty$ for all $(i_1, \dots, i_G) \in I_n^G$. Then,

$$E\Phi\left(\left\|\sum_{I_n^G} h_{i_1 \dots i_G}(X_{i_1}, \dots, X_{i_G})\right\|\right) \leq E\Phi\left(C_G \left\|\sum_{I_n^G} h_{i_1 \dots i_G}(X_{i_1}, \dots, X_{i_G})\right\|\right) \quad (57)$$

where $C_G = 2^G [G^G - 1] [(G-1)^{(G-1)} - 1] \times \dots \times 3$.

Lemma OA-7 gives the inequality result stated in the first half of Theorem 3.1.1 of de la Peña and Giné (1999). Theorem 3.1.1 also gives a reverse inequality under some additional symmetry conditions on the kernel $h_{i_1 \dots i_G}$ which we will not give here, since we will not be using the reverse inequality any of the results stated below. Proof of a more general decoupling inequality which contains the inequality given in expression (57) as a special case is provided in de la Peña (1992). See Theorem 2 of de la Peña (1992).

Lemma OA-8:

Let $D(\varepsilon \circ \varepsilon) = \text{diag}(\varepsilon_{(1,1)}^2, \dots, \varepsilon_{(n,T_n)}^2)$. Under Assumptions 2, 5, and 6; the following statements hold.

(a)

$$\frac{\Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon}{n} = O_p \left(\sqrt{\frac{K_{2,n}}{n}} \right),$$

where $D_{\sigma^2} = \text{diag}\left(E\left[\varepsilon_{(1,1)}^2 | \mathcal{F}_n^Z\right], \dots, E\left[\varepsilon_{(n,T_n)}^2 | \mathcal{F}_n^Z\right]\right)$.

(b)

$$\frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A Z_2 \Upsilon}{n} - \frac{\Upsilon' Z_2' M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} = O_p \left(\sqrt{\frac{K_{2,n}}{n}} \right) = o_p(1).$$

Proof of Lemma OA-8:

To show part (a), let $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, and note that we can write

$$\begin{aligned} & \frac{a' \Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon b}{n} - \frac{a' \Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon b}{n} \\ &= \frac{1}{n} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} \sum_{(k,v)=1}^{m_n} P_{(i,t),(j,s)}^\perp P_{(i,t),(k,v)}^\perp a' \Upsilon' Z_{2,(j,s)} Z_{2,(k,v)}' \Upsilon b \left\{ \varepsilon_{(i,t)}^2 - E\left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z\right] \right\} \end{aligned}$$

Now, making use of the CS inequality as well as Assumptions 2(i), 3(iii) and 5; we obtain

$$\begin{aligned} & E \left[\left(\frac{1}{n} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} \sum_{(k,v)=1}^{m_n} P_{(i,t),(j,s)}^\perp P_{(i,t),(k,v)}^\perp a' \Upsilon' Z_{2,(j,s)} Z_{2,(k,v)}' \Upsilon b \left\{ \varepsilon_{(i,t)}^2 - E\left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z\right] \right\} \right)^2 | \mathcal{F}_n^Z \right] \\ &= \frac{1}{n^2} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} \sum_{(k,v)=1}^{m_n} \sum_{(\ell,h)=1}^{m_n} \sum_{(r,c)=1}^{m_n} P_{(i,t),(j,s)}^\perp P_{(i,t),(k,v)}^\perp P_{(i,t),(\ell,h)}^\perp P_{(i,t),(r,c)}^\perp \\ & \quad \times (a' \Upsilon' Z_{2,(j,s)}) (a' \Upsilon' Z_{2,(\ell,h)}) (Z_{2,(k,v)}' \Upsilon b) (Z_{2,(r,c)}' \Upsilon b) E \left\{ (\varepsilon_{(i,t)}^2 - E\left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z\right])^2 | \mathcal{F}_n^Z \right\} \\ &= \frac{1}{n^2} \sum_{(i,t)=1}^{m_n} E \left\{ (\varepsilon_{(i,t)}^2 - E\left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z\right])^2 | \mathcal{F}_n^Z \right\} (e'_{(i,t)} P^\perp Z_2 \Upsilon a)^2 (e'_{(i,t)} P^\perp Z_2 \Upsilon b)^2 \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E\left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z\right] \right) \frac{1}{n^2} \sum_{(i,t)=1}^{m_n} (a' \Upsilon' Z_2' P^\perp e_{(i,t)} e'_{(i,t)} P^\perp Z_2 \Upsilon a) (b' \Upsilon' Z_2' P^\perp e_{(i,t)} e'_{(i,t)} P^\perp Z_2 \Upsilon b) \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E\left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z\right] \right) \frac{1}{n} \left(a' \Upsilon' Z_2' P^\perp \sum_{(i,t)=1}^{m_n} e_{(i,t)} e'_{(i,t)} P^\perp Z_2 \Upsilon a \right) e'_{(i,t)} P^\perp e_{(i,t)} \frac{b' \Upsilon' Z_2' Z_2 \Upsilon b}{n} \\ &\leq \left(\max_{1 \leq (i,t) \leq m_n} E\left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z\right] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \right) \frac{a' \Upsilon' Z_2' P^\perp Z_2 \Upsilon a}{n} \frac{b' \Upsilon' Z_2' Z_2 \Upsilon b}{n} \\ &= O_{a.s.} \left(\frac{K_{2,n}}{n} \right) = o_{a.s.}(1). \end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$\begin{aligned} & E \left[\frac{n}{K_{2,n}} \left(\frac{a' \Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon b}{n} - \frac{a' \Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon b}{n} \right)^2 \right] \\ = & E_Z \left(\frac{n}{K_{2,n}} E \left[\left(\frac{a' \Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon b}{n} - \frac{a' \Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon b}{n} \right)^2 | \mathcal{F}_n^Z \right] \right) \leq \bar{C}. \end{aligned}$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\begin{aligned} & \Pr \left(\sqrt{\frac{n}{K_{2,n}}} \left| \frac{a' \Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon b}{n} - \frac{a' \Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon b}{n} \right| \geq \sqrt{\frac{\bar{C}}{\epsilon}} \right) \\ = & \Pr \left(\frac{n}{K_{2,n}} \left(\frac{a' \Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon b}{n} - \frac{a' \Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon b}{n} \right)^2 \geq \frac{\bar{C}}{\epsilon} \right) \\ \leq & \frac{\epsilon}{\bar{C}} E \left[\frac{n}{K_{2,n}} \left(\frac{a' \Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon b}{n} - \frac{a' \Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon b}{n} \right)^2 \right] \leq \epsilon \end{aligned}$$

for all n sufficiently large, which shows that

$$\frac{a' \Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon b}{n} - \frac{a' \Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon b}{n} = O_p \left(\sqrt{\frac{K_{2,n}}{n}} \right).$$

Since the above result holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we further deduce that

$$\begin{aligned} & \frac{\Upsilon' Z_2' P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon}{n} \\ = & \frac{1}{n} \sum_{(i,t)=1}^{m_n} \sum_{(j,s)=1}^{m_n} \sum_{(k,v)=1}^{m_n} P_{(i,t),(j,s)}^\perp P_{(i,t),(k,v)}^\perp \gamma_{(j,s)} \gamma'_{(k,v)} \left\{ \varepsilon_{(i,t)}^2 - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right\} \\ = & O_p \left(\sqrt{\frac{K_{2,n}}{n}} \right). \end{aligned}$$

Next, consider part (b). Using the fact that $A = P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)}$ and that $M^{(Z,Q)} Z_2 = 0$

and applying the result of part (a) above, we obtain

$$\begin{aligned}
& \frac{\Upsilon' Z'_2 A D(\varepsilon \circ \varepsilon) A Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} \\
= & \frac{\Upsilon' Z'_2 P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon}{n} \\
& - \frac{\Upsilon' Z'_2 P^\perp D(\varepsilon \circ \varepsilon) M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} Z_2 \Upsilon}{n} + \frac{\Upsilon' Z'_2 M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} D(\varepsilon \circ \varepsilon) M^{(Z, Q)} D_{\widehat{\vartheta}} M^{(Z, Q)} Z_2 \Upsilon}{n} \\
= & \frac{\Upsilon' Z'_2 P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} \\
& + \frac{\Upsilon' Z'_2 P^\perp D(\varepsilon \circ \varepsilon) P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon}{n} \\
= & \frac{\Upsilon' Z'_2 P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} + O_p \left(\sqrt{\frac{K_{2,n}}{n}} \right)
\end{aligned} \tag{58}$$

Next, note that

$$\begin{aligned}
& \frac{\Upsilon' Z'_2 P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} \\
= & \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} + \frac{\Upsilon' Z'_2 [P^\perp - M^{(Z_1, Q)}] D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} \\
& + \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} [P^\perp - M^{(Z_1, Q)}] Z_2 \Upsilon}{n} + \frac{\Upsilon' Z'_2 [P^\perp - M^{(Z_1, Q)}] D_{\sigma^2} [P^\perp - M^{(Z_1, Q)}] Z_2 \Upsilon}{n} \\
= & \frac{\Upsilon' Z'_2 [P^\perp - M^{(Z_1, Q)}] D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} + \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} D_{\sigma^2} [P^\perp - M^{(Z_1, Q)}] Z_2 \Upsilon}{n} \\
& + \frac{\Upsilon' Z'_2 [P^\perp - M^{(Z_1, Q)}] D_{\sigma^2} [P^\perp - M^{(Z_1, Q)}] Z_2 \Upsilon}{n} \\
= & 0.
\end{aligned}$$

where the last equality follows from the fact that

$$\begin{aligned}
[P^\perp - M^{(Z_1, Q)}] Z_2 &= [P^{(Z, Q)} - P^{(Z_1, Q)} - (I_{m_n} - P^{(Z_1, Q)})] Z_2 \\
&= -M^{(Z, Q)} Z_2 \\
&= 0.
\end{aligned}$$

It follows from expression (58) that

$$\begin{aligned}
& \frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A Z_2 \Upsilon}{n} - \frac{\Upsilon' Z_2' M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} \\
&= \frac{\Upsilon' Z_2' P^\perp D_{\sigma^2} P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z_2' M^{(Z_1, Q)} D_{\sigma^2} M^{(Z_1, Q)} Z_2 \Upsilon}{n} + O_p \left(\sqrt{\frac{K_{2,n}}{n}} \right) \\
&= O_p \left(\sqrt{\frac{K_{2,n}}{n}} \right) = o_p(1),
\end{aligned}$$

as required. \square

Lemma OA-9:

Suppose that Assumptions 1-6 are satisfied. Then,

(a)

$$\begin{aligned}
& D_\mu^{-1} U' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} \\
& - \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] D_\mu^{-1} E \left[U_{(j,s)} U'_{(j,s)} | \mathcal{F}_n^Z \right] D_\mu^{-1} \\
&= O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) = o_p(1)
\end{aligned}$$

(b)

$$\frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1}}{\sqrt{n}} = o_p(1)$$

Proof of Lemma OA-9:

To show part (a), let $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, and define $u_{a,(i,t)} = a' D_\mu^{-1} U_{(i,t)}$ and

$u_{b,(j,s)} = b'D_\mu^{-1}U_{(j,s)}$. Note that

$$\begin{aligned}
& a'D_\mu^{-1}U'AD(\varepsilon \circ \varepsilon)AUD_\mu^{-1}b \\
& - \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] a'D_\mu^{-1}E \left[U_{(j,s)}U'_{(j,s)} | \mathcal{F}_n^Z \right] D_\mu^{-1}b \\
= & \sum_{(j,s),(k,v)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \\
& - \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \\
= & \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left(\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right) \\
& + \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)}
\end{aligned} \tag{59}$$

Focusing on the first term in expression (59) above, we apply the CS inequality, parts (b) and

(c) of Lemma S2-1, and Assumptions 2(i) and 3(ii) to obtain

$$\begin{aligned}
& E \left[\left(\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left(\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right) \right)^2 | \mathcal{F}_n^Z \right] \\
&= \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 E \left\{ \left(\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right)^2 | \mathcal{F}_n^Z \right\} \\
&\quad + \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 \left\{ E \left[\varepsilon_{(j,s)}^2 u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(i,t)}^2 u_{a,(i,t)} u_{b,(i,t)} | \mathcal{F}_n^Z \right] \right. \\
&\quad \left. - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(j,s)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(i,t)} u_{b,(i,t)} | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right\} \\
&\quad + \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (k,v)}}^{m_n} \sum_{(j,s) \neq \{(i,t),(k,v)\}} A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 \left\{ E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(k,v)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)}^2 u_{b,(j,s)}^2 | \mathcal{F}_n^Z \right] \right. \\
&\quad \left. - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(k,v)}^2 | \mathcal{F}_n^Z \right] \left(E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right)^2 \right\} \\
&\quad + 2 \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)}^2 A_{(k,v),(i,t)}^2 \left\{ E \left[\varepsilon_{(i,t)}^2 u_{a,(i,t)} u_{b,(i,t)} | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(k,v)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right. \\
&\quad \left. - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(k,v)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(i,t)} u_{b,(i,t)} | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right\} \\
&\quad + \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (k,v)}}^{m_n} \sum_{(j,s) \neq \{(i,t),(k,v)\}} \left\{ A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 \right. \\
&\quad \times \left(E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] E \left[u_{a,(k,v)} u_{b,(k,v)} | \mathcal{F}_n^Z \right] \right. \\
&\quad \left. - \left(E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] E \left[u_{a,(k,v)} u_{b,(k,v)} | \mathcal{F}_n^Z \right] \right) \Big\} \\
&\leq 2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right) \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z \right] \right) \frac{1}{(\mu_n^{\min})^4} \left\{ 2 \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 \right. \\
&\quad + \sum_{\substack{(i,t),(k,v),(j,s)=1 \\ (i,t) \neq (k,v), \\ (j,s) \neq (i,t), (j,s) \neq (k,v)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(k,v),(j,s)}^2 + 2 \sum_{\substack{(i,t),(k,v),(j,s)=1 \\ (j,s) \neq (k,v), \\ (i,t) \neq (j,s), (i,t) \neq (k,v)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(k,v),(i,t)}^2 + \sum_{\substack{(i,t),(k,v),(j,s)=1 \\ (i,t) \neq (k,v), \\ (j,s) \neq (i,t), (j,s) \neq (k,v)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.} \left(\frac{K_{2,n}^3}{(\mu_n^{\min})^4 n^2} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right) \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$\begin{aligned}
& E \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 (\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] E[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z]) \right)^2 \right] \\
&= \frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \times \\
& \quad E_Z \left(E \left[\left(\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 (\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] E[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z]) \right)^2 | \mathcal{F}_n^Z \right] \right) \\
&\leq \bar{C}.
\end{aligned}$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\begin{aligned}
& \Pr \left(\left| \frac{(\mu_n^{\min})^2 \sqrt{n}}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 (\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] E[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z]) \right| \geq \sqrt{\frac{\bar{C}}{\epsilon}} \right) \\
&= \Pr \left\{ \frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 (\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] E[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z]) \right)^2 \geq \frac{\bar{C}}{\epsilon} \right\} \\
&\leq \epsilon \frac{E \left[\frac{(\mu_n^{\min})^4 n}{K_{2,n}^2} \left(\sum_{(i,t) \neq (j,s)} A_{(i,t),(j,s)}^2 \left\{ \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] E[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z] \right\} \right)^2 \right]}{\bar{C}} \\
&\leq \epsilon
\end{aligned}$$

for all n sufficiently large, which shows that

$$\begin{aligned}
& \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 (\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] E[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z]) \\
&= O_p \left(\frac{K_{2,n}}{(\mu_n^{\min})^2 \sqrt{n}} \right). \tag{60}
\end{aligned}$$

Turning our attention to the second term in expression (59), we first define $\varsigma_{(i,t)} = \varepsilon_{(i,t)}^2 - E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z]$. Note that by the decoupling inequality given in Lemma OA-7, there exist finite constants C_2 and C_3 , whose explicit forms are given in Lemma OA-7, and independent copies $\{(u_{a,(i,t)}^{(g)}, u_{b,(i,t)}^{(g)}, \varsigma_{(i,t)}^{(g)})\}_{(i,t)=1}^{m_n}$ (for $g = 1, 2, 3$) of the sequence $\{(u_{a,(i,t)}, u_{b,(i,t)}, \varsigma_{(i,t)})\}_{(i,t)=1}^{m_n}$ such

that

$$\begin{aligned}
& E \left[\left(\sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
\leq & 2E \left[\left(\sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] u_{a,(j,s)} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
& + 2E \left[\left(\sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} \left[\varepsilon_{(i,t)}^2 - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right] \right)^2 | \mathcal{F}_n^Z \right] \\
\leq & 2C_2 E \left[\left(\sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] u_{a,(j,s)}^{(1)} u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \\
& + 2C_3 E \left[\left(\sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} u_{a,(j,s)}^{(1)} u_{b,(k,v)}^{(2)} \varsigma_{(i,t)}^{(3)} \right)^2 | \mathcal{F}_n^Z \right] \\
= & 2C_2 \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v)}}^m \sum_{(i,t) \neq \{(j,s),(k,v)\}} \sum_{(\ell,h) \neq \{(i,t),(j,s),(k,v)\}} \left\{ A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(\ell,h),(j,s)} A_{(\ell,h),(k,v)} \right. \\
& \quad \times E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(\ell,h)}^2 | \mathcal{F}_n^Z \right] E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \} \\
& + 2C_2 \sum_{\substack{(j,s),(k,v),(i,t)=1 \\ (j,s) \neq (k,v) \\ (i,t) \neq \{(j,s),(k,v)\}}}^m A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 \left(E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \\
& + 2C_3 \sum_{\substack{(j,s),(k,v),(i,t)=1 \\ (j,s) \neq (k,v) \\ (i,t) \neq \{(j,s),(k,v)\}}}^m A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 \left\{ E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \right. \\
& \quad \times E \left[\left(\varsigma_{(i,t)}^{(3)} \right)^2 | \mathcal{F}_n^Z \right] \} \tag{61}
\end{aligned}$$

Define $D_{\sigma^2} = \text{diag} \left(\sigma_{(1,2)}^2, \dots, \sigma_{(n,T_n)}^2 \right)$ where $\sigma_{(i,t)}^2 = E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right]$ for $(i,t) = 1, \dots, m_n$. By applying Assumptions 2(i), 3(ii), and 5 as well as parts (a) and (b) of Lemma OA-1, we can estimate the

(almost sure) order of magnitude of the first term on the right-hand side of (61) as follows

$$\begin{aligned}
& \sum_{(j,s),(k,v)=1}^{m_n} \sum_{\substack{(i,t) \neq \{(j,s),(k,v)\} \\ (j,s) \neq (k,v)}} \sum_{(\ell,h) \neq \{(i,t),(j,s),(k,v)\}} \left\{ A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(\ell,h),(j,s)} A_{(\ell,h),(k,v)} \right. \\
& \quad \times E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(\ell,h)}^2 | \mathcal{F}_n^Z \right] E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \left. \right\} \\
= & \sum_{(j,s)=1}^{m_n} E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] \sum_{\substack{(k,v)=1 \\ (k,v) \neq (j,s)}}^{m_n} e'_{(j,s)} A \sum_{(i,t) \neq \{(j,s),(k,v)\}} e_{(i,t)} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] e'_{(i,t)} A e_{(k,v)} \\
& \quad \times E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] e'_{(k,v)} \sum_{(\ell,h) \neq \{(i,t),(j,s),(k,v)\}} A e_{(\ell,h)} E \left[\varepsilon_{(\ell,h)}^2 | \mathcal{F}_n^Z \right] e'_{(\ell,h)} A e_{(j,s)} \\
= & \sum_{(j,s)=1}^{m_n} E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] \sum_{\substack{(k,v)=1 \\ (k,v) \neq (j,s)}}^{m_n} e'_{(j,s)} A \sum_{(i,t) \neq \{(j,s),(k,v)\}} e_{(i,t)} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] e'_{(i,t)} A e_{(k,v)} \\
& \quad \times E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \left[e'_{(k,v)} A D_{\sigma^2} A e_{(j,s)} - e'_{(k,v)} A e_{(i,t)} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] e'_{(i,t)} A e_{(j,s)} \right] \\
= & \sum_{(j,s)=1}^{m_n} E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] \sum_{\substack{(k,v)=1 \\ (k,v) \neq (j,s)}}^{m_n} E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \left[e'_{(k,v)} A D_{\sigma^2} A e_{(j,s)} \right]^2 \\
& \quad - \sum_{(j,s)=1}^{m_n} E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] \\
& \quad \times \sum_{\substack{(k,v),(i,t)=1 \\ (k,v) \neq (j,s) \\ (i,t) \neq \{(j,s),(k,v)\}}}^{m_n} \left[e'_{(j,s)} A e_{(i,t)} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] e'_{(i,t)} A e_{(k,v)} \right]^2 E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \\
\leq & \sum_{(j,s)=1}^{m_n} E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] \sum_{(k,v)=1}^{m_n} E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] e'_{(j,s)} A D_{\sigma^2} A e_{(k,v)} e'_{(k,v)} A D_{\sigma^2} A e_{(j,s)} \\
\leq & \left(\max_{1 \leq (i,t) \leq m_n} E \left[\| U_{(i,t)} \|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \frac{1}{(\mu_n^{\min})^4} \sum_{(j,s)=1}^{m_n} e'_{(j,s)} A D_{\sigma^2} A^2 D_{\sigma^2} A e_{(j,s)}
\end{aligned}$$

$$\begin{aligned}
&\leq \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \left(1 + \max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \right) \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 \\
&\quad \times \frac{1}{(\mu_n^{\min})^4} \sum_{(j,s)=1}^{m_n} e'_{(j,s)} A^2 e_{(j,s)} \\
&\leq \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \left(1 + \max_{1 \leq (i,t) \leq m_n} |\widehat{\vartheta}_{(i,t)}|^2 \right) \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 \\
&\quad \times \frac{1}{(\mu_n^{\min})^4} \left[\text{tr} \left\{ P^\perp \right\} + \text{tr} \left\{ D_{\widehat{\vartheta}}^2 \right\} \right] \\
&= O_{a.s.} \left(\frac{K_{2,n}}{(\mu_n^{\min})^4} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right) = O_{a.s.} \left(\frac{K_{2,n}}{(\mu_n^{\min})^4} \right).
\end{aligned}$$

By applying Assumptions 2(i) and 3(ii) as well as part (c) of Lemma S2-1, we can also derive the (almost sure) order of magnitude for the second and the third terms of (61) as follows

$$\begin{aligned}
&\sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v), (i,t) \\ (k,v) \neq (i,t)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 \left(E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \\
&\leq \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \frac{1}{(\mu_n^{\min})^4} \\
&\quad \times \sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right) = o_{a.s.}(1)
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq (k,v), (i,t) \\ (k,v) \neq (i,t)}}^m \left\{ A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 E \left[\left(u_{a,(j,s)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] \right. \\
& \quad \times E \left[\left(u_{b,(k,v)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(\zeta_{(i,t)}^{(3)} \right)^2 | \mathcal{F}_n^Z \right] \left. \right\} \\
& \leq \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right) \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \frac{1}{(\mu_n^{\min})^4} \\
& \quad \times \sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}}^m A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 \\
& = O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^4 n} \right) = o_{a.s.}(1).
\end{aligned}$$

These results imply that

$$\begin{aligned}
& E \left[\left(\sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
& = O_{a.s.} \left(\frac{K_{2,n}}{(\mu_n^{\min})^4} \right)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$\begin{aligned}
& E \left[\frac{(\mu_n^{\min})^4}{K_{2,n}} \left(\sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \right)^2 \right] \\
& = E_Z \left(E \left[\frac{(\mu_n^{\min})^4}{K_{2,n}} \left(\sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}}^m A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \right) \\
& \leq \bar{C}.
\end{aligned}$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\begin{aligned}
& \Pr \left(\frac{(\mu_n^{\min})^2}{\sqrt{K_{2,n}}} \left| \sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \right| \geq \sqrt{\frac{C}{\epsilon}} \right) \\
&= \Pr \left\{ \frac{(\mu_n^{\min})^4}{K_{2,n}} \left(\sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \right)^2 \geq \frac{C}{\epsilon} \right\} \\
&\leq \frac{\epsilon}{\bar{C}} E \left[\frac{(\mu_n^{\min})^4}{K_{2,n}} \left(\sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \right)^2 \right] \leq \epsilon
\end{aligned}$$

for all n sufficiently large, which shows that

$$\sum_{(i,t)=1}^{m_n} \sum_{\substack{(j,s),(k,v)=1 \\ (j,s) \neq \{(k,v),(i,t)\}, (k,v) \neq (i,t)}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} = O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) \quad (62)$$

Now, (60) and (62) together imply that

$$\begin{aligned}
& a' D_\mu^{-1} U' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} b \\
& - \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] a' D_\mu^{-1} E \left[U_{(j,s)} U'_{(j,s)} | \mathcal{F}_n^Z \right] D_\mu^{-1} b \\
&= \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left(\varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(j,s)} - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right) \\
&+ \sum_{(j,s),(k,v)=1}^{m_n} \sum_{\substack{(i,t) \neq \{(j,s),(k,v)\} \\ (j,s) \neq (k,v)}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} \varepsilon_{(i,t)}^2 u_{a,(j,s)} u_{b,(k,v)} \\
&= O_p \left(\frac{K_{2,n}}{(\mu_n^{\min})^2 \sqrt{n}} \right) + O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right) \\
&= O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right).
\end{aligned}$$

Finally, since the above argument holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we obtain the

desired result

$$\begin{aligned}
& D_\mu^{-1} U' A D (\varepsilon \circ \varepsilon) A U D_\mu^{-1} - \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] D_\mu^{-1} E \left[U_{(j,s)} U'_{(j,s)} | \mathcal{F}_n^Z \right] D_\mu^{-1} \\
&= O_p \left(\frac{\sqrt{K_{2,n}}}{(\mu_n^{\min})^2} \right).
\end{aligned}$$

To show part (b), again let $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$ and define $u_{b,(k,v)} = b' D_\mu^{-1} U_{(k,v)}$ and $u_b = U D_\mu^{-1} b$. We can apply Loèeve's c_r inequality to obtain

$$\begin{aligned}
& E \left[\left(\frac{a' \Upsilon' Z'_2 A D (\varepsilon \circ \varepsilon) A U D_\mu^{-1} b}{\sqrt{n}} \right)^2 | \mathcal{F}_n^Z \right] \\
&= \frac{1}{n} E \left[\left(\sum_{(j,s),(k,v)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} a' \Upsilon' Z_{2,(j,s)} \varepsilon_{(i,t)}^2 u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
&\leq \frac{2}{n} E \left[\left(\sum_{(j,s),(k,v)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} a' \Upsilon' Z_{2,(j,s)} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
&\quad + \frac{2}{n} E \left[\left(\sum_{(j,s),(k,v)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} a' \Upsilon' Z_{2,(j,s)} \left\{ \varepsilon_{(i,t)}^2 - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right\} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
&= \frac{2}{n} E \left[(a' \Upsilon' Z'_2 A D_{\sigma^2} A u_b)^2 | \mathcal{F}_n^Z \right] \\
&\quad + \frac{2}{n} E \left[\left(\sum_{(j,s),(k,v)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} a' \Upsilon' Z_{2,(j,s)} \varsigma_{(i,t)} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \tag{63}
\end{aligned}$$

where $\varsigma_{(i,t)} = \varepsilon_{(i,t)}^2 - E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right]$. Focusing on the first term of expression (63) above, we can

apply Assumptions 2(i), 3(ii), and 3(iii) as well as part (b) of Lemma OA-1 to obtain

$$\begin{aligned}
& \frac{2}{n} E \left[(a' \Upsilon' Z_2' A D_{\sigma^2} A u_b)^2 | \mathcal{F}_n^Z \right] \\
= & \frac{2}{n} a' \Upsilon' Z_2' A D_{\sigma^2} A E \left[u_b u_b' | \mathcal{F}_n^Z \right] A D_{\sigma^2} A Z_2 \Upsilon a \\
\leq & \frac{1}{(\mu_n^{\min})^2} \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right) \frac{a' \Upsilon' Z_2' A D_{\sigma^2} A^2 D_{\sigma^2} A Z_2 \Upsilon a}{n} \\
\leq & \frac{1}{(\mu_n^{\min})^2} \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right) \left(1 + \max_{1 \leq (i,t) \leq m_n} |\hat{\vartheta}_{(i,t)}|^2 \right)^2 \\
& \times \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 \frac{a' \Upsilon' Z_2' \Upsilon a}{n} \\
= & O_{a.s.} \left(\frac{1}{(\mu_n^{\min})^2} \right) = o_{a.s.}(1).
\end{aligned}$$

Turning our attention now to the second term of (63), note that we can apply the inequality $|XY| \leq (1/2) X^2 + (1/2) Y^2$, Assumptions 1, 2(i), and 3(iii) as well as part (c) of Lemma S2-1 to

get

$$\begin{aligned}
& \frac{2}{n} E \left[\left(\sum_{(j,s),(k,v)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} a' \Upsilon' Z_{2,(j,s)} \varsigma_{(i,t)} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
& \leq \frac{2}{n} \sum_{(j,s),(k,v),(\ell,h)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v),(\ell,h)\}} \left| A_{(i,t),(j,s)} A_{(i,t),(\ell,h)} A_{(i,t),(k,v)}^2 \right. \\
& \quad \left. \times a' \Upsilon' Z_{2,(j,s)} a' \Upsilon' Z_{2,(\ell,h)} E \left[\varsigma_{(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{b,(k,v)}^2 | \mathcal{F}_n^Z \right] \right| \\
& \quad + \frac{2}{n} \sum_{(j,s),(k,v),(\ell,h)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v),(\ell,h)\}} \left| A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(k,v),(\ell,h)} A_{(k,v),(i,t)} \right. \\
& \quad \left. \times a' \Upsilon' Z_{2,(j,s)} a' \Upsilon' Z_{2,(\ell,h)} E \left[\varsigma_{(i,t)} u_{b,(i,t)} | \mathcal{F}_n^Z \right] E \left[\varsigma_{(k,v)} u_{b,(k,v)} | \mathcal{F}_n^Z \right] \right| \\
& \leq \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right)^2 \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n (\mu_n^{\min})^2} \sum_{(i,t)=1}^{m_n} \sum_{\substack{(k,v),(\ell,h)=1 \\ (k,v) \neq (i,t) \\ (\ell,h) \neq (i,t)}}^{m_n} A_{(i,t),(\ell,h)}^2 A_{(i,t),(k,v)}^2 \\
& \quad + \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n (\mu_n^{\min})^2} \sum_{(i,t)=1}^{m_n} \sum_{\substack{(k,v),(j,s)=1 \\ (k,v) \neq (i,t) \\ (j,s) \neq (i,t)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 \\
& \quad + \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right) \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right) \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n (\mu_n^{\min})^2} \\
& \quad \times \sum_{(k,v)=1}^{m_n} \sum_{\substack{(i,t),(\ell,h)=1 \\ (i,t) \neq (k,v), (\ell,h) \neq (k,v)}}^{m_n} A_{(k,v),(\ell,h)}^2 A_{(i,t),(k,v)}^2 \\
& \quad + \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right) \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right) \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n (\mu_n^{\min})^2} \\
& \quad \times \sum_{(i,t)=1}^{m_n} \sum_{\substack{(k,v),(j,s)=1 \\ (k,v) \neq (i,t), (j,s) \neq (i,t)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(k,v),(i,t)}^2 \\
& = O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^2 n} \right)
\end{aligned}$$

It follows from these results that

$$\begin{aligned}
& E \left[\left(\frac{a' \Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} b}{\sqrt{n}} \right)^2 | \mathcal{F}_n^Z \right] \\
& \leq \frac{2}{n} E \left[(a' \Upsilon' Z_2' A D_{\sigma^2} A u_b)^2 | \mathcal{F}_n^Z \right] \\
& \quad + \frac{2}{n} E \left[\left(\sum_{(j,s),(k,v)=1}^{m_n} \sum_{(i,t) \neq \{(j,s),(k,v)\}} A_{(i,t),(j,s)} A_{(i,t),(k,v)} a' \gamma_{(j,s)} \varsigma_{(i,t)} u_{b,(k,v)} \right)^2 | \mathcal{F}_n^Z \right] \\
& = O_{a.s.} \left(\frac{1}{(\mu_n^{\min})^2} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{(\mu_n^{\min})^2 n} \right) = o_{a.s.}(1)
\end{aligned}$$

Now, by the conditional version of the Markov's inequality, we deduce that, for any $\epsilon > 0$,

$$\Pr \left(\left| \frac{a' \Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \rightarrow 0 \text{ a.s.}$$

Since

$$\sup_n E \left[\left| \Pr \left(\left| \frac{a' \Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \right|^2 \right] < \infty,$$

it then follows by a version of the dominated convergence theorem, as given by Theorem 25.12 of Billingsley (1986), that as $n \rightarrow \infty$

$$\begin{aligned}
& \Pr \left(\left| \frac{a' \Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \right) \\
& = E \left[\Pr \left(\left| \frac{a' \Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \right] \rightarrow 0,
\end{aligned}$$

Finally, since the above result holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we further deduce that

$$\frac{\Upsilon' Z_2' A D(\varepsilon \circ \varepsilon) A U D_\mu^{-1}}{\sqrt{n}} \xrightarrow{p} 0, \text{ as } n \rightarrow \infty$$

as required for part (b). \square

Lemma OA-10:

Under Assumptions 5 and 6, the following statements are true.

(a)

$$\begin{aligned}
\text{tr} \{A^4\} &= \text{tr} \left\{ \left(P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right)^4 \right\} \\
&= O_{a.s.}(K_{2,n}).
\end{aligned}$$

(b) $|S_n| = O_{a.s.}(K_{2,n})$, where

$$\begin{aligned}
S_n &= \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} (A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(k,v),(\ell,h)} \\
&\quad + A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(j,s),(\ell,h)} A_{(k,v),(\ell,h)} \\
&\quad + A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)}) \\
(c) \quad &\left| \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)} \right| = O_{a.s.}(K_{2,n}).
\end{aligned}$$

Proof of Lemma OA-10:

To show part (a), note first that $P^\perp M^{(Z,Q)} = 0 = M^{(Z,Q)}P^\perp$. Now, write

$$\begin{aligned}
&\left(P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right)^4 \\
&= \left[\left(P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right) \left(P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right) \right]^2 \\
&= \left(P^\perp + M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right)^2 \\
&= \left(P^\perp + M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right) \left(P^\perp + M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right) \\
&= P^\perp + M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)}
\end{aligned}$$

Hence,

$$\begin{aligned}
&\text{tr} \left\{ \left(P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right)^4 \right\} \\
&= \text{tr} \left\{ P^\perp \right\} + \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right\} \\
&= K_{2,n} + \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right\}
\end{aligned}$$

In addition, note that

$$\begin{aligned}
0 &\leq \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right\} \\
&\leq \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}}^2 M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right\} \\
&\leq \left(\max_{(i,t) \in \Lambda_1} |\widehat{\vartheta}_{(i,t)}|^2 \right) \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right\} \\
&\leq \left(\max_{(i,t) \in \Lambda_1} |\widehat{\vartheta}_{(i,t)}|^2 \right) \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}}^2 M^{(Z,Q)} \right\} \\
&= \left(\max_{(i,t) \in \Lambda_1} |\widehat{\vartheta}_{(i,t)}|^2 \right) \text{tr} \left\{ D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} \right\} \\
&\leq \left(\max_{(i,t) \in \Lambda_1} |\widehat{\vartheta}_{(i,t)}|^2 \right) \text{tr} \left\{ D_{\widehat{\vartheta}}^2 \right\}
\end{aligned}$$

It follows from applying parts (a) and (b) of Lemma OA-1 that

$$\begin{aligned}
\text{tr} \{A^4\} &= \text{tr} \left\{ \left(P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right)^4 \right\} \\
&= K_{2,n} + \text{tr} \left\{ M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} \right\} \\
&= K_{2,n} + \left(\max_{(i,t) \in \Lambda_1} \left| \widehat{\vartheta}_{(i,t)} \right|^2 \right) \text{tr} \{D_{\widehat{\vartheta}}^2\} \\
&= O_{a.s.}(K_{2,n}).
\end{aligned}$$

Next, to show part (b), decompose $A = P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)}$ as follows

$$A = L + L',$$

where L be the lower triangular matrix such that $L_{(i,t),(j,s)} = A_{(i,t),(j,s)} \mathbb{I}\{(i,t) > (j,s)\}$, i.e., L is lower triangular matrix whose lower triangular elements correspond to the lower triangular elements of A . It follows that

$$\begin{aligned}
A^4 &= (L + L')^4 = \left(L^2 + LL' + L'L + (L')^2 \right)^2 \\
&= L^4 + L^3 L' + L^2 L'L + L^2 (L')^2 + LL'L^2 + LL'LL' + L (L')^2 L + L (L')^3 \\
&\quad + L'L^3 + L'L^2 L' + L'LL'L + L'L (L')^2 + (L')^2 L^2 + (L')^2 LL' + (L')^3 L + (L')^4
\end{aligned}$$

Using the fact that $\text{tr} \{AB\} = \text{tr} \{BA\}$ and $\text{tr} \{A\} = \text{tr} \{A'\}$, we obtain

$$\text{tr} \{A^4\} = 2\text{tr} \{L^4\} + 8\text{tr} \{L^3 L'\} + 4\text{tr} \{L^2 (L')^2\} + 2\text{tr} \{LL'LL'\}$$

We compute each of the terms on the right-hand side above as follows.

$$\begin{aligned}
&\text{tr} \{L^4\} \\
&= \sum_{(i,t)} \sum_{(j,s)} \sum_{(k,v)} \sum_{(\ell,h)} [A_{(i,t),(j,s)} \mathbb{I}\{(i,t) > (j,s)\} A_{(j,s),(k,v)} \mathbb{I}\{(j,s) > (k,v)\} \\
&\quad \times A_{(k,v),(\ell,h)} \mathbb{I}\{(k,v) > (\ell,h)\} A_{(\ell,h),(i,t)} \mathbb{I}\{(\ell,h) > (i,t)\}] \\
&= 0
\end{aligned}$$

$$\begin{aligned}
& \text{tr} \{ L^3 L' \} \\
&= \sum_{(i,t)} \sum_{(j,s)} \sum_{(k,v)} \sum_{(\ell,h)} [A_{(i,t),(j,s)} \mathbb{I}\{(i,t) > (j,s)\} A_{(j,s),(k,v)} \mathbb{I}\{(j,s) > (k,v)\} \\
&\quad \times A_{(k,v),(\ell,h)} \mathbb{I}\{(k,v) > (\ell,h)\} A_{(\ell,h),(i,t)} \mathbb{I}\{(i,t) > (\ell,h)\}] \\
&= \sum_{1 \leq (\ell,h) < (k,v) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(\ell,h),(i,t)} \\
&= \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(j,s)} A_{(k,v),(j,s)} A_{(j,s),(i,t)} A_{(i,t),(\ell,h)} \\
&= \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(\ell,h),(i,t)} A_{(k,v),(\ell,h)} \text{ (by symmetry of } A) \\
\\
& \text{tr} \{ L^2 (L')^2 \} \\
&= \sum_{(i,t)} \sum_{(j,s)} \sum_{(k,v)} \sum_{(\ell,h)} [A_{(i,t),(j,s)} \mathbb{I}\{(i,t) > (j,s)\} A_{(j,s),(k,v)} \mathbb{I}\{(j,s) > (k,v)\} \\
&\quad \times A_{(k,v),(\ell,h)} \mathbb{I}\{(\ell,h) > (k,v)\} A_{(\ell,h),(i,t)} \mathbb{I}\{(i,t) > (\ell,h)\}] \\
&= \sum_{1 \leq (k,v) < (\ell,h) = (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(\ell,h),(i,t)} \\
&\quad + \sum_{1 \leq (k,v) < (\ell,h) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(\ell,h),(i,t)} \\
&\quad + \sum_{1 \leq (k,v) < (j,s) < (\ell,h) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(\ell,h),(i,t)} \\
&= \sum_{1 \leq (k,v) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(j,s)} A_{(j,s),(i,t)} \\
&\quad + \sum_{1 \leq (k,v) < (\ell,h) < (j,s) < (i,t) \leq m_n} [A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(\ell,h),(i,t)} \\
&\quad \quad + A_{(i,t),(\ell,h)} A_{(\ell,h),(k,v)} A_{(k,v),(j,s)} A_{(j,s),(i,t)}] \\
&= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 \\
&\quad + \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} [A_{(\ell,h),(k,v)} A_{(k,v),(i,t)} A_{(i,t),(j,s)} A_{(j,s),(\ell,h)} \\
&\quad \quad + A_{(\ell,h),(j,s)} A_{(j,s),(i,t)} A_{(i,t),(k,v)} A_{(k,v),(\ell,h)}] \\
&= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 \\
&\quad + 2 \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(j,s),(\ell,h)} A_{(k,v),(\ell,h)}
\end{aligned}$$

$$\begin{aligned}
& \operatorname{tr} \{LL'LL'\} \\
= & \sum_{(i,t)} \sum_{(j,s)} \sum_{(k,v)} \sum_{(\ell,h)} [A_{(i,t),(j,s)} \mathbb{I}\{(i,t) > (j,s)\} A_{(j,s),(k,v)} \mathbb{I}\{(k,v) > (j,s)\} \\
& \quad \times A_{(k,v),(\ell,h)} \mathbb{I}\{(k,v) > (\ell,h)\} A_{(\ell,h),(i,t)} \mathbb{I}\{(i,t) > (\ell,h)\}] \\
= & \sum_{1 \leq (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(i,t)} A_{(i,t),(j,s)} A_{(j,s),(i,t)} \\
& + \sum_{1 \leq (j,s) < (k,v) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(j,s)} A_{(j,s),(i,t)} \\
& + \sum_{1 \leq (j,s) < (i,t) < (k,v) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(j,s)} A_{(j,s),(i,t)} \\
& + \sum_{1 \leq (j,s) < (\ell,h) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(i,t)} A_{(i,t),(\ell,h)} A_{(\ell,h),(i,t)} \\
& + \sum_{1 \leq (\ell,h) < (j,s) < (i,t) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(i,t)} A_{(i,t),(\ell,h)} A_{(\ell,h),(i,t)} \\
& + \left(\sum_{1 \leq (\ell,h) < (j,s) < (k,v) < (i,t) \leq m_n} + \sum_{1 \leq (j,s) < (\ell,h) < (k,v) < (i,t) \leq m_n} \right) A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(\ell,h),(i,t)} \\
& + \left(\sum_{1 \leq (\ell,h) < (j,s) < (i,t) < (k,v) \leq m_n} + \sum_{1 \leq (j,s) < (\ell,h) < (i,t) < (k,v) \leq m_n} \right) A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(\ell,h),(i,t)} \\
= & \sum_{1 \leq (i,t) < (j,s) \leq m_n} A_{(i,t),(j,s)}^4 + 2 \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} (A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2) \\
& + 4 \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)}
\end{aligned}$$

It follows that

$$\begin{aligned}
& \operatorname{tr} \{A^4\} \\
= & 8 \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} [A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(k,v),(\ell,h)} \\
& + A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(j,s),(\ell,h)} A_{(k,v),(\ell,h)} \\
& + A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)}] \\
& + 2 \sum_{1 \leq (i,t) < (j,s) \leq m_n} A_{(i,t),(j,s)}^4 \\
& + 4 \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} (A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2 + A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2) \\
= & 8S_n + 2 \sum_{1 \leq (i,t) < (j,s) \leq m_n} A_{(i,t),(j,s)}^4 \\
& + 4 \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} (A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2 + A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2)
\end{aligned}$$

It follows from the triangle inequality, the result given in part (a) above, parts (b) and (c) of Lemma S2-1, and the symmetry of A that

$$\begin{aligned}
|S_n| & \leq \frac{1}{8} \operatorname{tr} \{A^4\} + \frac{1}{4} \sum_{1 \leq (i,t) < (j,s) \leq m_n} A_{(i,t),(j,s)}^4 \\
& + \frac{1}{2} \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} (A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2 + A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2) \\
& \leq \frac{1}{8} \operatorname{tr} \{A^4\} + \frac{1}{4} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 + \frac{3}{2} \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 \\
& = O_{a.s.}(K_{2,n}) + O_{a.s.}\left(\frac{K_{2,n}^2}{n^2}\right) + O_{a.s.}\left(\frac{K_{2,n}^2}{n}\right) \\
& = O_{a.s.}(K_{2,n})
\end{aligned}$$

Finally, for part (c), we take $\{\eta_{(i,t)}\}$ to be a double-indexed sequence of *i.i.d.* random variables with mean 0 and variance 1 and where $\eta_{(i,t)}$ and Z are independent for all (i,t) and n . Define the

random quantities

$$\begin{aligned}
\Delta_1 &= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} \left[A_{(i,t),(j,s)} A_{(i,t),(k,v)} \eta_{(j,s)} \eta_{(k,v)} + A_{(i,t),(j,s)} A_{(j,s),(k,v)} \eta_{(i,t)} \eta_{(k,v)} \right. \\
&\quad \left. + A_{(i,t),(k,v)} A_{(j,s),(k,v)} \eta_{(i,t)} \eta_{(j,s)} \right] \\
\Delta_2 &= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} \left[A_{(i,t),(j,s)} A_{(i,t),(k,v)} \eta_{(j,s)} \eta_{(k,v)} + A_{(i,t),(j,s)} A_{(j,s),(k,v)} \eta_{(i,t)} \eta_{(k,v)} \right] \\
\Delta_3 &= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} \eta_{(i,t)} \eta_{(j,s)}
\end{aligned}$$

Next, note that using part (c) of Lemma S2-1 and the symmetry of A , we have

$$\begin{aligned}
E[\Delta_3^2 | \mathcal{F}_n^Z] &= E \left[\left(\sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} \eta_{(i,t)} \eta_{(j,s)} \right)^2 | \mathcal{F}_n^Z \right] \\
&= \sum_{1 \leq (i,t) < (j,s) < \{(k,v), (\ell,h)\} \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)} \\
&= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2 \\
&\quad + 2 \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)} \\
&\leq \sum_{(k,v)=1}^{m_n} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(k,v)}^2 A_{(k,v),(j,s)}^2 \\
&\quad + 2 \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)} \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) + 2 \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)}
\end{aligned}$$

$$\begin{aligned}
& E [\Delta_2 \Delta_3 | \mathcal{F}_n^Z] \\
= & E \left[\left(\sum_{1 \leq (i_1, t_1) < (j_1, s_1) < (k_1, g_1) \leq m_n} \left\{ A_{(i_1, t_1), (j_1, s_1)} A_{(i_1, t_1), (k_1, g_1)} \eta_{(j_1, s_1)} \eta_{(k_1, g_1)} \right. \right. \right. \\
& \quad \left. \left. \left. + A_{(i_1, t_1), (j_1, s_1)} A_{(j_1, s_1), (k_1, g_1)} \eta_{(i_1, t_1)} \eta_{(k_1, g_1)} \right\} \right) \right. \\
& \quad \times \left. \left(\sum_{1 \leq (i_2, t_2) < (j_2, s_2) < (k_2, g_2) \leq m_n} A_{(i_2, t_2), (k_2, g_2)} A_{(j_2, s_2), (k_2, g_2)} \eta_{(i_2, t_2)} \eta_{(j_2, s_2)} \right) \right| \mathcal{F}_n^Z \Bigg] \\
= & \sum_{1 \leq (i, t) < (j, s) < (k, v) < (\ell, h) \leq m_n} \left[A_{(i, t), (j, s)} A_{(i, t), (k, v)} A_{(j, s), (\ell, h)} A_{(k, v), (\ell, h)} \right. \\
& \quad \left. + A_{(i, t), (j, s)} A_{(j, s), (k, v)} A_{(i, t), (\ell, h)} A_{(k, v), (\ell, h)} \right]
\end{aligned}$$

and

$$\begin{aligned}
& E [\Delta_2^2 | \mathcal{F}_n^Z] \\
&= E \left[\left(\sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} \left\{ A_{(i,t),(j,s)} A_{(i,t),(k,v)} \zeta_{(j,s)} \zeta_{(k,v)} + A_{(i,t),(j,s)} A_{(j,s),(k,v)} \eta_{(i,t)} \eta_{(k,v)} \right\} \right)^2 | \mathcal{F}_n^Z \right] \\
&= \sum_{1 \leq \{(i,t),(\ell,h)\} < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(\ell,h),(j,s)} A_{(\ell,h),(k,v)} \\
&\quad + \sum_{1 \leq (i,t) < \{(j,s),(\ell,h)\} < (k,v) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(\ell,h),(k,v)} \\
&\quad + \sum_{1 \leq (i,t) < (j,s) < (\ell,h) < (k,v) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(j,s),(\ell,h)} A_{(\ell,h),(k,v)} \\
&\quad + \sum_{1 \leq (\ell,h) < (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(\ell,h),(i,t)} A_{(\ell,h),(k,v)} \\
&= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 \\
&\quad + 2 \sum_{1 \leq (i,t) < (\ell,h) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(k,v)} A_{(\ell,h),(j,s)} A_{(\ell,h),(k,v)} \\
&\quad + 2 \sum_{1 \leq (i,t) < (j,s) < (\ell,h) < (k,v) \leq m_n} A_{(i,t),(j,s)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(\ell,h),(k,v)} \\
&\quad + \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(j,s)} A_{(i,t),(\ell,h)} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} \\
&\quad + \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(j,s),(k,v)} A_{(k,v),(\ell,h)} A_{(i,t),(j,s)} A_{(i,t),(\ell,h)} \\
&= \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 + 2S_n \\
&\leq 2 \sum_{(j,s)=1}^{m_n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (j,s), (k,v) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 A_{(j,s),(k,v)}^2 + 2S_n \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) + O_{a.s.} (K_{2,n}) = O_{a.s.} (K_{2,n})
\end{aligned}$$

Since $\Delta_1 = \Delta_2 + \Delta_3$, it follows from part (b) of this lemma and the results given above that

$$\begin{aligned}
E [\Delta_1^2 | \mathcal{F}_n^Z] &= E [\Delta_2^2 | \mathcal{F}_n^Z] + E [\Delta_3^2 | \mathcal{F}_n^Z] + 2E [\Delta_2 \Delta_3 | \mathcal{F}_n^Z] \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) + 4S_n \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) + O_{a.s.} (K_{2,n}) = O_{a.s.} (K_{2,n}).
\end{aligned}$$

Moreover, note that

$$\begin{aligned}
& \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)} \\
&= \frac{1}{2} E [\Delta_3^2 | \mathcal{F}_n^Z] - \frac{1}{2} \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2 \\
&= \frac{1}{2} E [(\Delta_1 - \Delta_2)^2 | \mathcal{F}_n^Z] - \frac{1}{2} \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2
\end{aligned}$$

so that, by making use of the triangle inequality, Loèeve's c_r inequality, and the symmetry of A ; we obtain

$$\begin{aligned}
& \left| \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} A_{(i,t),(\ell,h)} A_{(j,s),(\ell,h)} \right| \\
&\leq E [\Delta_1^2 | \mathcal{F}_n^Z] + E [\Delta_2^2 | \mathcal{F}_n^Z] + \frac{1}{2} \sum_{1 \leq (i,t) < (j,s) < (k,v)} A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2 \\
&\leq E [\Delta_1^2 | \mathcal{F}_n^Z] + E [\Delta_2^2 | \mathcal{F}_n^Z] + \frac{1}{2} \sum_{(k,v)=1}^{m_n} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (k,v), (j,s) \neq (k,v)}}^{m_n} A_{(i,t),(k,v)}^2 A_{(k,v),(j,s)}^2 \\
&= O_{a.s.}(K_{2,n}). \quad \square
\end{aligned}$$

Lemma OA-11:

Let L be the lower triangular matrix such that $L_{(i,t),(j,s)} = A_{(i,t),(j,s)} \mathbb{I}\{(i,t) > (j,s)\}$. Then, under Assumptions 5-6,

$$\|LL'\|_F = O_{a.s.} \left(\sqrt{K_{2,n}} \right),$$

where $\|\cdot\|_F$ denotes the Frobenius norm, i.e., $\|A\|_F = [tr(A'A)]^{1/2}$.

Proof of Lemma OA-11:

By parts (b) and (c) of Lemma S2-1 and part (c) of Lemma OA-10, we have

$$\begin{aligned}
& \|LL'\|_F^2 \\
&= \text{tr} \{ LL' LL' \} \\
&= \sum_{1 \leq (i,t) < (j,s) \leq m_n} A_{(i,t),(j,s)}^4 + 2 \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} [A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2] \\
&\quad + 4 \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(k,v),(j,s)} A_{(j,s),(\ell,h)} A_{(\ell,h),(i,t)} \\
&\leq \sum_{1 \leq (i,t) < (j,s) \leq m_n} A_{(i,t),(j,s)}^4 + 2 \sum_{1 \leq (i,t) < (j,s) < (k,v) \leq m_n} [A_{(i,t),(j,s)}^2 A_{(i,t),(k,v)}^2 + A_{(i,t),(k,v)}^2 A_{(j,s),(k,v)}^2] \\
&\quad + 4 \left| \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(k,v),(j,s)} A_{(j,s),(\ell,h)} A_{(\ell,h),(i,t)} \right| \\
&\leq \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^4 + 4 \sum_{(k,v)=1}^{m_n} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (k,v), (j,s) \neq (k,v)}}^{m_n} A_{(i,t),(k,v)}^2 A_{(k,v),(j,s)}^2 \\
&\quad + 4 \left| \sum_{1 \leq (i,t) < (j,s) < (k,v) < (\ell,h) \leq m_n} A_{(i,t),(k,v)} A_{(k,v),(j,s)} A_{(j,s),(\ell,h)} A_{(\ell,h),(i,t)} \right| \\
&= O_{a.s.} \left(\frac{K_{2,n}^2}{n^2} \right) + O_{a.s.} \left(\frac{K_{2,n}^2}{n} \right) + O_{a.s.} (K_{2,n}) = O_{a.s.} (K_{2,n}),
\end{aligned}$$

from which the required result follows. \square

Lemma OA-12:

Under Assumptions 1-6, the following statements are true.

(a)

$$\frac{\Upsilon' Z'_2 A Z_2 \Upsilon}{n} = \frac{\Upsilon' Z'_2 M^{(Z_1, Q)} Z_2 \Upsilon}{n} = O_{a.s.} (1)$$

(b)

$$D_\mu^{-1} U' A U D_\mu^{-1} = o_p (1)$$

(c)

$$\frac{\Upsilon' Z'_2 A U D_\mu^{-1}}{\sqrt{n}} = o_p (1), \quad \frac{D_\mu^{-1} U' A Z_2 \Upsilon}{\sqrt{n}} = o_p (1).$$

Proof of Lemma OA-12:

To show part (a), note that, by making use of the fact that $M^{(Z,Q)}Z_2 = 0$, we have

$$\begin{aligned}
\frac{\Upsilon' Z'_2 A Z_2 \Upsilon}{n} &= \frac{\Upsilon' Z'_2 P^\perp Z_2 \Upsilon}{n} - \frac{\Upsilon' Z'_2 M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} Z_2 \Upsilon}{n} \\
&= \frac{\Upsilon' Z'_2 P^\perp Z_2 \Upsilon}{n} \\
&= \frac{\Upsilon' Z'_2 (P^{(Z,Q)} - P^{(Z_1,Q)}) Z_2 \Upsilon}{n} \\
&= \frac{\Upsilon' Z'_2 (M^{(Z_1,Q)} - M^{(Z,Q)}) Z_2 \Upsilon}{n} \\
&= \frac{\Upsilon' Z'_2 M^{(Z_1,Q)} Z_2 \Upsilon}{n}
\end{aligned}$$

Moreover, part (iii) of Assumption 3 implies that

$$\frac{\Upsilon' Z'_2 M^{(Z_1,Q)} Z_2 \Upsilon}{n} \leq \frac{\Upsilon' Z'_2 Z_2 \Upsilon}{n} = O_{a.s.}(1)$$

where we take $A \leq B$ for two square matrices A and B to mean that $A - B$ is negative semi-definite, or, alternatively, $B - A$ is positive semidefinite. It follows immediately that

$$\frac{\Upsilon' Z'_2 A Z_2 \Upsilon}{n} = \frac{\Upsilon' Z'_2 M^{(Z_1,Q)} Z_2 \Upsilon}{n} = O_{a.s.}(1),$$

as required.

To show part (b), note that, for $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we write

$$\begin{aligned}
a'D_\mu^{-1}U'AUD_\mu^{-1}b &= a'D_\mu^{-1} \left(\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)} U_{(i,t)} U'_{(j,s)} \right) D_\mu^{-1}b \\
&= \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)} u_{a,(i,t)} u_{b,(j,s)},
\end{aligned}$$

where $u_{a,(i,t)} = a'D_\mu^{-1}U_{(i,t)}$ and $u_{b,(j,s)} = b'D_\mu^{-1}U_{(j,s)}$. Next, making use of the CS inequality, part

(a) of Lemma S2-1, and Assumptions 1, 2(i), 3(ii), and 5; we get

$$\begin{aligned}
& E \left[(a' D_\mu^{-1} U' A U D_\mu^{-1} b)^2 | \mathcal{F}_n^Z \right] \\
= & E \left[\left(\sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)} u_{a,(i,t)} u_{b,(j,s)} \right)^2 | \mathcal{F}_n^Z \right] \\
= & \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} \sum_{\substack{(k,v),(\ell,h)=1 \\ (k,v) \neq (\ell,h)}}^{m_n} A_{(i,t),(j,s)} A_{(k,v),(\ell,h)} E \left[u_{a,(i,t)} u_{b,(j,s)} u_{a,(k,v)} u_{b,(\ell,h)} | \mathcal{F}_n^Z \right] \\
= & \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ E \left[u_{a,(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{b,(j,s)}^2 | \mathcal{F}_n^Z \right] + E \left[u_{a,(i,t)} u_{b,(i,t)} | \mathcal{F}_n^Z \right] E \left[u_{a,(j,s)} u_{b,(j,s)} | \mathcal{F}_n^Z \right] \right\} \\
\leq & \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ E \left[u_{a,(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{b,(j,s)}^2 | \mathcal{F}_n^Z \right] + E \left[|u_{a,(i,t)} u_{b,(i,t)}| | \mathcal{F}_n^Z \right] E \left[|u_{a,(j,s)} u_{b,(j,s)}| | \mathcal{F}_n^Z \right] \right\} \\
\leq & \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 E \left[u_{a,(i,t)}^2 | \mathcal{F}_n^Z \right] E \left[u_{b,(j,s)}^2 | \mathcal{F}_n^Z \right] \\
& + \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sqrt{E \left[u_{a,(i,t)}^2 | \mathcal{F}_n^Z \right]} \sqrt{E \left[u_{b,(i,t)}^2 | \mathcal{F}_n^Z \right]} \sqrt{E \left[u_{a,(j,s)}^2 | \mathcal{F}_n^Z \right]} \sqrt{E \left[u_{b,(j,s)}^2 | \mathcal{F}_n^Z \right]} \\
\leq & \frac{CK_{2,n}}{(\mu_n^{\min})^4} \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
= & O_{a.s.} \left(\frac{K_{2,n}}{(\mu_n^{\min})^4} \right) \\
= & o_{a.s.}(1),
\end{aligned}$$

It follows from the conditional version of the Markov's inequality that for any $\epsilon > 0$

$$\Pr(|a' D_\mu^{-1} U' A U D_\mu^{-1} b| \geq \epsilon | \mathcal{F}_n^Z) \rightarrow 0 \text{ a.s.}$$

Moreover, note that

$$\sup_n E \left[\left| \Pr(|a' D_\mu^{-1} U' A U D_\mu^{-1} b| \geq \epsilon | \mathcal{F}_n^Z) \right|^2 \right] < \infty$$

Hence, by a version of the dominated convergence theorem, as given by Theorem 25.12 of Billingsley

(1986), it follows that as $n \rightarrow \infty$

$$\begin{aligned} & \Pr(|a'D_\mu^{-1}U'AUD_\mu^{-1}b| \geq \epsilon) \\ &= E[\Pr(|a'D_\mu^{-1}U'AUD_\mu^{-1}b| \geq \epsilon | \mathcal{F}_n^Z)] \rightarrow 0 \end{aligned}$$

Since the above result holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we further deduce that

$$D_\mu^{-1}U'AUD_\mu^{-1} = o_p(1),$$

as required.

To show part (c), again, let $u_{b,(i,t)} = U_{(i,t)}D_\mu^{-1}b$. Note again that, by making use of the conditional serial independence assumption in Assumption 1, we have

$$\begin{aligned} & E\left(\left[\frac{a'\Upsilon'Z'_2AUD_\mu^{-1}b}{\sqrt{n}}\right]^2 | \mathcal{F}_n^Z\right) \\ &= E\left(\left[\frac{1}{\sqrt{n}} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (k,v)}}^{m_n} A_{(i,t),(k,v)} a'\Upsilon'Z_{2,(i,t)} u_{b,(k,v)}\right]^2 | \mathcal{F}_n^Z\right) \\ &= \frac{1}{n} \sum_{\substack{(i,t),(k,v)=1 \\ (i,t) \neq (k,v)}}^{m_n} \sum_{\substack{(j,s),(\ell,h)=1 \\ (j,s) \neq (\ell,h)}}^{m_n} A_{(i,t),(k,v)} A_{(j,s),(\ell,h)} a'\Upsilon'Z_{2,(i,t)} a'\Upsilon'Z_{2,(j,s)} E[u_{b,(k,v)} u_{b,(\ell,h)} | \mathcal{F}_n^Z] \\ &= \frac{1}{n} \sum_{(k,v)=1}^{m_n} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (k,v), (j,s) \neq (k,v)}}^{m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} a'\Upsilon'Z_{2,(i,t)} a'\Upsilon'Z_{2,(j,s)} b'D_\mu^{-1} E[U_{(k,v)} U'_{(k,v)} | \mathcal{F}_n^Z] D_\mu^{-1} b \\ &\leq \frac{\max_{1 \leq (k,v) \leq m_n} E[\|U_{(k,v)}\|_2^2 | \mathcal{F}_n^Z]}{(\mu_n^{\min})^2} \frac{1}{n} \sum_{(k,v)=1}^{m_n} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (k,v), (j,s) \neq (k,v)}}^{m_n} A_{(i,t),(k,v)} A_{(j,s),(k,v)} a'\Upsilon'Z_{2,(i,t)} a'\Upsilon'Z_{2,(j,s)} \\ &= \frac{\max_{1 \leq (k,v) \leq m_n} E[\|U_{(k,v)}\|_2^2 | \mathcal{F}_n^Z]}{(\mu_n^{\min})^2} \frac{a'\Upsilon'Z'_2 A A' Z_2 \Upsilon a}{n} \end{aligned}$$

Next, by note that

$$\begin{aligned}
& \frac{a' \Upsilon' Z_2' A A' Z_2 \Upsilon a}{n} \\
= & \frac{a' \Upsilon' Z_2' (P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)}) (P^\perp - M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)}) Z_2 \Upsilon a}{n} \\
= & \frac{a' \Upsilon' Z_2' (P^\perp + M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)} D_{\widehat{\vartheta}} M^{(Z,Q)}) Z_2 \Upsilon a}{n} \\
= & \frac{a' \Upsilon' Z_2' P^\perp Z_2 \Upsilon a}{n} \\
\leq & \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n}
\end{aligned}$$

from which it follows by Assumptions 2 and 3(iii) that

$$\begin{aligned}
& E \left(\left[\frac{a' \Upsilon' Z_2' A U D_\mu^{-1} b}{\sqrt{n}} \right]^2 \mid \mathcal{F}_n^Z \right) \\
\leq & \frac{\max_{1 \leq (k,v) \leq m_n} E \left[\|U_{(k,v)}\|_2^2 \mid \mathcal{F}_n^Z \right]}{(\mu_n^{\min})^2} \frac{a' \Upsilon' Z_2' Z_2 \Upsilon a}{n} \\
= & O_{a.s.} \left(\frac{1}{(\mu_n^{\min})^2} \right) = o_{a.s.}(1)
\end{aligned}$$

It further follows from the conditional version of the Markov's inequality that for any $\epsilon > 0$

$$\Pr \left(\left| \frac{a' \Upsilon' Z_2' A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \rightarrow 0 \text{ a.s.}$$

In addition, note that

$$\sup_n E \left[\left| \Pr \left(\left| \frac{a' \Upsilon' Z_2' A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \right|^2 \right] < \infty$$

Hence, by a version of the dominated convergence theorem, as given by Theorem 25.12 of Billingsley (1986), it follows that as $n \rightarrow \infty$

$$\Pr \left(\left| \frac{a' \Upsilon' Z_2' A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \right) = E \left[\Pr \left(\left| \frac{a' \Upsilon' Z_2' A U D_\mu^{-1} b}{\sqrt{n}} \right| \geq \epsilon \mid \mathcal{F}_n^Z \right) \right] \rightarrow 0$$

Since the above result holds for all $a, b \in \mathbb{R}^d$ such that $\|a\|_2 = \|b\|_2 = 1$, we further deduce that

$$\frac{\Upsilon' Z_2' A U D_\mu^{-1}}{\sqrt{n}} = o_p(1),$$

Moreover, it follows immediately that

$$\frac{D_\mu^{-1}U'AZ_2\Upsilon}{\sqrt{n}} = \left(\frac{\Upsilon'Z_2'AUD_\mu^{-1}}{\sqrt{n}} \right)' = o_p(1),$$

as required. \square

Lemma OA-13: Let $A = P^\perp - M^{(Z,Q)}D_{\widehat{\vartheta}}M^{(Z,Q)}$. Then, under Assumptions 2-6, the following result holds as $K_{2,n}$, $n \rightarrow \infty$.

$$\sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_1),(j,s_1)}^2 A_{(i,t_2),(j,s_2)}^2 = O_{a.s.} \left(\frac{K_{2,n}^3}{n^2} \right).$$

Proof of Lemma OA-13:

To proceed, we apply the inequality $|XY| \leq (1/2)X^2 + (1/2)Y^2$ as well as the result given part (b) of Lemma S2-1 to obtain

$$\begin{aligned} & \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_1),(j,s_1)}^2 A_{(i,t_2),(j,s_2)}^2 \\ & \leq \frac{1}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_1),(j,s_1)}^4 + \frac{1}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_2),(j,s_2)}^4 \\ & \leq \frac{\bar{T}^2}{2} \sum_{\substack{(i,t_1),(j,s_1)=1 \\ (i,t_1) \neq (j,s_1)}}^{m_n} A_{(i,t_1),(j,s_1)}^4 + \frac{\bar{T}^2}{2} \sum_{\substack{(i,t_2),(j,s_2)=1 \\ (i,t_2) \neq (j,s_2)}}^{m_n} A_{(i,t_2),(j,s_2)}^4 \quad (\text{by Assumption 6(ii)}) \\ & = O_{a.s.} \left(\frac{K_{2,n}^3}{n^2} \right) + O_{a.s.} \left(\frac{K_{2,n}^3}{n^2} \right) \\ & = O_{a.s.} \left(\frac{K_{2,n}^3}{n^2} \right). \quad \square \end{aligned}$$

Lemma OA-14: Suppose that Assumptions 1, 2, 5, and 6 are satisfied. Then, the following statements are true.

(a)

$$\mathcal{A}_{1,2} = O_p \left(\frac{1}{\sqrt{n}} \right),$$

where

$$\begin{aligned} \mathcal{A}_{1,2} = & \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} \left(M_{(i,h),(i,g)}^Q \right)^2 \varepsilon_{(i,g)}^2 \right. \\ & \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(j,q)} \varepsilon_{(j,c)} \right\} \end{aligned}$$

(b)

$$\mathcal{A}_{1,3} = O_p \left(\frac{1}{\sqrt{n}} \right),$$

where

$$\begin{aligned} \mathcal{A}_{1,3} = & \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{\substack{g=1 \\ r=1 \\ r \neq g}}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \varepsilon_{(i,g)} \varepsilon_{(i,r)} \right. \\ & \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \left(M_{(j,v),(j,q)}^Q \right)^2 \varepsilon_{(j,q)}^2 \right\} \end{aligned}$$

(c)

$$\mathcal{A}_{1,4} = O_p \left(\frac{K_{2,n}}{n} \right),$$

where

$$\begin{aligned} \mathcal{A}_{1,4} = & \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{\substack{g=1 \\ r=1 \\ r \neq g}}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \varepsilon_{(i,g)} \varepsilon_{(i,r)} \right. \\ & \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(j,q)} \varepsilon_{(j,c)} \right\} \end{aligned}$$

(d)

$$\mathcal{A}_1 = O_p \left(\frac{1}{\sqrt{n}} \right),$$

where

$$\begin{aligned}
& \mathcal{A}_1 \\
= & \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} M^Q \varepsilon \right)^2 \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
& - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2
\end{aligned}$$

Proof of Lemma OA-14:

To show part (a), write

$$\begin{aligned}
\mathcal{A}_{1,2} & = \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \varepsilon_{(i,g)}^2 \\
& \quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(j,q)} \varepsilon_{(j,c)} \\
& = \frac{1}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{t,s=1 \\ t \neq s}}^{T_i} A_{(i,t),(i,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \\
& \quad \times \sum_{v=1}^{T_j} J_{(i,s),(i,v)} \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_i} M_{(i,v),(i,q)}^Q M_{(i,v),(i,c)}^Q \varepsilon_{(i,q)}^2 \varepsilon_{(i,c)} \\
& \quad + \frac{1}{K_{2,n}} \sum_{i,j=1}^n \sum_{\substack{t=1 \\ i \neq j}}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \\
& \quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(j,q)}^2 \varepsilon_{(j,c)} \\
& = \mathcal{A}_{1,2,1} + \mathcal{A}_{1,2,2}
\end{aligned}$$

Now, applying part (f) of Lemma S2-1 as well as Assumptions 2(i), 5, and 6; we obtain

$$\begin{aligned}
& E[|\mathcal{A}_{1,2,1}| \mid \mathcal{F}_n^Z] \\
\leq & \frac{1}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{t,s=1 \\ t \neq s}}^{T_i} A_{(i,t),(i,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left(M_{(i,h),(i,g)}^Q \right)^2 \\
& \times \sum_{v=1}^{T_j} |J_{(i,s),(i,v)}| \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_i} \left| M_{(i,v),(i,q)}^Q \right| \left| M_{(i,v),(i,c)}^Q \right| E \left[\left| \varepsilon_{(i,g)}^2 \varepsilon_{(i,q)} \varepsilon_{(i,c)} \right| \mid \mathcal{F}_n^Z \right] \\
\leq & \bar{T}^3 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} \left| M_{(i,t),(j,s)}^Q \right| \right)^4 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 \mid \mathcal{F}_n^Z \right] \right) \frac{1}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{t,s=1 \\ t \neq s}}^{T_i} A_{(i,t),(i,s)}^2 \\
= & O_{a.s.} \left(\frac{K_{2,n}}{n} \right)
\end{aligned}$$

Moreover, let $\varsigma_{(i,g)} = \varepsilon_{(i,g)}^2 - E \left[\varepsilon_{(i,g)}^2 \mid \mathcal{F}_n^Z \right]$, and, making use of Loève's c_r inequality, we can write

$$\begin{aligned}
& E[\mathcal{A}_{1,2,2}^2 \mid \mathcal{F}_n^Z] \\
\leq & 2E \left[\left(\frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \right. \right. \\
& \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varsigma_{(i,g)} \varepsilon_{(j,q)} \varepsilon_{(j,c)} \left. \right)^2 \mid \mathcal{F}_n^Z \left. \right] \\
& + 2E \left[\left(\frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \right. \right. \\
& \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q E \left[\varepsilon_{(i,g)}^2 \mid \mathcal{F}_n^Z \right] \varepsilon_{(j,q)} \varepsilon_{(j,c)} \left. \right)^2 \mid \mathcal{F}_n^Z \left. \right].
\end{aligned}$$

Now, note that, by the decoupling inequality given in Lemma OA-7, there exists a finite constant C_3 , whose explicit form is given in Lemma OA-7, and independent copies $\left\{ \left(\varsigma_{(i,t)}^{(\ell)}, \varepsilon_{(i,t)}^{(\ell)} \right) \right\}_{(i,t)=1}^{m_n}$ (for

$\ell = 1, 2, 3$) of the sequence $\{(\zeta_{(i,t)}, \varepsilon_{(i,t)})\}_{(i,t)=1}^{m_n}$ such that

$$\begin{aligned}
& 2E \left[\left(\frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \right. \right. \\
& \quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \zeta_{(i,g)} \varepsilon_{(j,q)} \varepsilon_{(j,c)} \left. \left. \right) ^2 | \mathcal{F}_n^Z \right] \\
& \leq (2 \cdot C_3) E \left[\left(\frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \right. \right. \\
& \quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \zeta_{(i,g)}^{(1)} \varepsilon_{(j,q)}^{(2)} \varepsilon_{(j,c)}^{(3)} \left. \left. \right) ^2 | \mathcal{F}_n^Z \right] \\
& = (2 \cdot C_3) \frac{1}{K_{2,n}^2} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_1),(j,s_1)}^2 A_{(i,t_2),(j,s_2)}^2 \\
& \quad \times \sum_{g=1}^{T_i} \sum_{h_1,h_2=1}^{T_i} |J_{(i,t_1),(i,h_1)}| |J_{(i,t_2),(i,h_2)}| \left(M_{(i,h_1),(i,g)}^Q \right)^2 \left(M_{(i,h_2),(i,g)}^Q \right)^2 \\
& \quad \times \sum_{v_1,v_2=1}^{T_j} |J_{(j,s_1),(j,v_1)}| |J_{(j,s_2),(j,v_2)}| \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}} |M_{(j,v_1),(j,q)}^Q| |M_{(j,v_1),(j,c)}^Q| |M_{(j,v_2),(j,q)}^Q| |M_{(j,v_2),(j,c)}^Q| \\
& \quad \times E \left[\left(\zeta_{(i,g)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(\varepsilon_{(j,q)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(\varepsilon_{(j,c)}^{(3)} \right)^2 | \mathcal{F}_n^Z \right] \\
& \leq (2 \cdot C_3) \overline{T}^3 \|J\|_\infty^4 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^8 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^2 \\
& \quad \times \frac{1}{K_{2,n}^2} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_1),(j,s_1)}^2 A_{(i,t_2),(j,s_2)}^2 \\
& = O_{a.s.} \left(\frac{K_{2,n}}{n^2} \right) = o_{a.s.} \left(\frac{1}{n} \right),
\end{aligned}$$

where the (almost sure) order of magnitude above is calculated using Lemma OA-13 and Assumptions 2(i), 5, and 6. Moreover, a further application of the decoupling inequality given in Lemma OA-7 shows that there exists a finite constant C_2 , whose explicit form is given in Lemma OA-7,

and independent copies $\{\varepsilon_{(i,t)}^{(\ell)}\}_{(i,t)=1}^{m_n}$ (for $\ell = 1, 2$) of the sequence $\{\varepsilon_{(i,t)}\}_{(i,t)=1}^{m_n}$ such that

$$\begin{aligned}
& 2E \left[\left(\frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \right. \right. \\
& \quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q E \left[\varepsilon_{(i,g)}^2 | \mathcal{F}_n^Z \right] \varepsilon_{(j,q)} \varepsilon_{(j,c)} \left. \right)^2 | \mathcal{F}_n^Z \left. \right] \\
& \leq 2C_2 E \left[\left(\frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \right. \right. \\
& \quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q E \left[\varepsilon_{(i,g)}^2 | \mathcal{F}_n^Z \right] \varepsilon_{(j,q)}^{(1)} \varepsilon_{(j,c)}^{(2)} \left. \right)^2 | \mathcal{F}_n^Z \left. \right] \\
& \leq 2C_2 \frac{1}{K_{2,n}^2} \sum_{i_1,i_2=1}^n \sum_{\substack{j=1 \\ j \neq i_1,i_2}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{t_2=1}^{T_{i_2}} \sum_{\substack{s_1,s_2=1 \\ j \neq i_1,i_2}}^{T_j} A_{(i_1,t_1),(j,s_1)}^2 A_{(i_2,t_2),(j,s_2)}^2 \\
& \quad \times \sum_{g_1=1}^{T_{i_1}} \sum_{g_2=1}^{T_{i_2}} \sum_{h_1=1}^{T_{i_1}} \sum_{h_2=1}^{T_{i_2}} |J_{(i_1,t_1),(i_1,h_1)}| |J_{(i_2,t_2),(i_2,h_2)}| \left(M_{(i_1,h_1),(i_1,g_1)}^Q \right)^2 \left(M_{(i_2,h_2),(i_2,g_2)}^Q \right)^2 \\
& \quad \times \sum_{v_1,v_2=1}^{T_j} |J_{(j,s_1),(j,v_1)}| |J_{(j,s_2),(j,v_2)}| \sum_{\substack{q=1 \\ c=1 \\ c \neq q}}^{T_j} \left| M_{(j,v_1),(j,q)}^Q \right| \left| M_{(j,v_1),(j,c)}^Q \right| \left| M_{(j,v_2),(j,q)}^Q \right| \left| M_{(j,v_2),(j,c)}^Q \right| \\
& \quad \times E \left[\varepsilon_{(i_1,g_1)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(i_2,g_2)}^2 | \mathcal{F}_n^Z \right] E \left[\left(\varepsilon_{(j,q)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(\varepsilon_{(j,c)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] \\
& \leq 2C_2 \bar{T}^4 \|J\|_\infty^4 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} \left| M_{(i,t),(j,s)}^Q \right| \right)^8 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^4 \\
& \quad \times \frac{1}{K_{2,n}^2} \sum_{i_1,i_2=1}^n \sum_{\substack{j=1 \\ j \neq i_1,i_2}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{t_2=1}^{T_{i_2}} \sum_{\substack{s_1,s_2=1 \\ j \neq i_1,i_2}}^{T_j} A_{(i_1,t_1),(j,s_1)}^2 A_{(i_2,t_2),(j,s_2)}^2 \\
& = O_{a.s.} \left(\frac{1}{n} \right)
\end{aligned}$$

where the (almost sure) order of magnitude above is calculated using part (e) of Lemma S2-1 and Assumptions 2(i), 5, and 6. It follows that

$$E \left[\mathcal{A}_{1,2,2}^2 | \mathcal{F}_n^Z \right] = O_{a.s.} \left(\frac{K_{2,n}}{n^2} \right) + O_{a.s.} \left(\frac{1}{n} \right) = O_{a.s.} \left(\frac{1}{n} \right),$$

so that, by the triangle inequality, the conditional version of Liapunov's inequality, and Assumption 5(ii), we further obtain

$$\begin{aligned} E[|\mathcal{A}_{1,2}| | \mathcal{F}_n^Z] &\leq E[|\mathcal{A}_{1,2,1}| | \mathcal{F}_n^Z] + E[|\mathcal{A}_{1,2,2}| | \mathcal{F}_n^Z] \\ &\leq E[|\mathcal{A}_{1,2,1}| | \mathcal{F}_n^Z] + \sqrt{E[\mathcal{A}_{1,2,2}^2 | \mathcal{F}_n^Z]} \\ &= O_{a.s.}\left(\frac{K_{2,n}}{n}\right) + O_{a.s.}\left(\frac{1}{\sqrt{n}}\right) = O_{a.s.}\left(\frac{1}{\sqrt{n}}\right) \end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E[\sqrt{n}|\mathcal{A}_{1,2}|] = E_W(\sqrt{n}E[|\mathcal{A}_{1,2}| | \mathcal{F}_n^W]) \leq \bar{C}.$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\Pr\left(|\sqrt{n}\mathcal{A}_{1,2}| \geq \frac{\bar{C}}{\epsilon}\right) \leq \epsilon \frac{\sqrt{n}E[|\mathcal{A}_{1,2}|]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{1,2} = O_p\left(\frac{1}{\sqrt{n}}\right). \quad (64)$$

To show part (b), note that since A is symmetric, so that $A_{(i,t),(j,s)} = A_{(j,s),(i,t)}$, it follows that $\mathcal{A}_{1,2} = \mathcal{A}_{1,3}$. Hence, using the same argument as that given for part (a) above, we can show that

$$\mathcal{A}_{1,3} = O_p\left(\frac{1}{\sqrt{n}}\right). \quad (65)$$

Turning our attention to part (c), note that, in this case, we can write

$$\begin{aligned}
\mathcal{A}_{1,4} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} \sum_{\substack{r=1 \\ r \neq g}}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \varepsilon_{(i,g)} \varepsilon_{(i,r)} \\
&\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(j,q)} \varepsilon_{(j,c)} \\
&= \frac{1}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{s,t=1 \\ s \neq t}}^{T_i} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} \sum_{\substack{r=1 \\ r \neq g}}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \\
&\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_i} \sum_{\substack{c=1 \\ c \neq q}}^{T_i} M_{(i,v),(i,q)}^Q M_{(i,v),(i,c)}^Q \varepsilon_{(i,g)} \varepsilon_{(i,r)} \varepsilon_{(i,q)} \varepsilon_{(i,c)} \\
&\quad + \frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} \sum_{\substack{r=1 \\ r \neq g}}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \\
&\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(i,g)} \varepsilon_{(i,r)} \varepsilon_{(j,q)} \varepsilon_{(j,c)} \\
&= \mathcal{A}_{1,4,1} + \mathcal{A}_{1,4,2}.
\end{aligned}$$

Now, by the triangle inequality, part (f) of Lemma S2-1, and Assumptions 2(i), 5, and 6; we have

$$\begin{aligned}
&E[|\mathcal{A}_{1,4,1}| |\mathcal{F}_n^Z] \\
&\leq \frac{1}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{s,t=1 \\ s \neq t}}^{T_i} A_{(i,t),(i,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{g=1}^{T_i} \sum_{\substack{r=1 \\ r \neq g}}^{T_i} \left| M_{(i,h),(i,g)}^Q \right| \left| M_{(i,h),(i,r)}^Q \right| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{q=1}^{T_i} \sum_{\substack{c=1 \\ c \neq q}}^{T_i} \left| M_{(i,v),(i,q)}^Q \right| \left| M_{(i,v),(i,c)}^Q \right| E[|\varepsilon_{(i,g)} \varepsilon_{(i,r)} \varepsilon_{(i,q)} \varepsilon_{(i,c)}| |\mathcal{F}_n^Z] \\
&\leq \bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} \left| M_{(i,t),(j,s)}^Q \right| \right)^4 \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^4] \right) \frac{1}{K_{2,n}} \sum_{i=1}^n \sum_{s,t=1}^{T_i} A_{(i,t),(i,s)}^2 \\
&= O_{a.s.} \left(\frac{K_{2,n}}{n} \right) = o_{a.s.}(1).
\end{aligned}$$

In addition, by the decoupling inequality given in Lemma OA-7, there exists a finite constant C_4 , whose explicit form is given in Lemma OA-7, and independent copies $\{\varepsilon_{(i,t)}^{(\ell)}\}_{(i,t)=1}^{m_n}$ (for $\ell = 1, 2, 3, 4$)

of the sequence $\{\varepsilon_{(i,t)}\}_{(i,t)=1}^{m_n}$ such that

$$\begin{aligned}
& E[\mathcal{A}_{1,4.2}^2 | \mathcal{F}_n^Z] \\
= & E \left[\left(\frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} \sum_{\substack{r=1 \\ r \neq g}}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \right. \right. \\
& \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(i,g)} \varepsilon_{(i,r)} \varepsilon_{(j,q)} \varepsilon_{(j,c)} \left. \left. \right)^2 | \mathcal{F}_n^Z \right] \\
\leq & \frac{C_4}{K_{2,n}^2} E \left[\left(\sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} \sum_{\substack{r=1 \\ r \neq g}}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \right. \right. \\
& \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(i,g)}^{(1)} \varepsilon_{(i,r)}^{(2)} \varepsilon_{(j,q)}^{(3)} \varepsilon_{(j,c)}^{(4)} \left. \left. \right)^2 | \mathcal{F}_n^Z \right] \\
\leq & \frac{C_4}{K_{2,n}^2} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_1),(j,s_1)}^2 A_{(i,t_1),(j,s_1)}^2 \sum_{h_1,h_2=1}^{T_i} |J_{(i,t_1),(i,h_1)}| |J_{(i,t_2),(i,h_2)}| \\
& \times \sum_{g=1}^{T_i} \sum_{\substack{r=1 \\ r \neq g}}^{T_i} \left| M_{(i,h_1),(i,g)}^Q \right| \left| M_{(i,h_1),(i,r)}^Q \right| \left| M_{(i,h_2),(i,g)}^Q \right| \left| M_{(i,h_2),(i,r)}^Q \right| \sum_{v_1,v_2=1}^{T_j} |J_{(j,s_1),(j,v_1)}| |J_{(j,s_2),(j,v_2)}| \\
& \times \sum_{q=1}^{T_j} \sum_{\substack{c=1 \\ c \neq q}}^{T_j} \left| M_{(j,v_1),(j,q)}^Q \right| \left| M_{(j,v_1),(j,c)}^Q \right| \left| M_{(j,v_2),(j,q)}^Q \right| \left| M_{(j,v_2),(j,c)}^Q \right| E \left[\left(\varepsilon_{(i,g)}^{(1)} \right)^2 | \mathcal{F}_n^Z \right] \\
& \times E \left[\left(\varepsilon_{(i,r)}^{(2)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(\varepsilon_{(j,q)}^{(3)} \right)^2 | \mathcal{F}_n^Z \right] E \left[\left(\varepsilon_{(j,c)}^{(4)} \right)^2 | \mathcal{F}_n^Z \right] \\
\leq & C_4 \bar{T}^4 \|J\|_\infty^4 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} \left| M_{(i,t),(j,s)}^Q \right| \right)^8 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^4 \\
& \times \frac{1}{K_{2,n}^2} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t_1,t_2=1}^{T_i} \sum_{s_1,s_2=1}^{T_j} A_{(i,t_1),(j,s_1)}^2 A_{(i,t_1),(j,s_1)}^2 \\
= & O_{a.s.} \left(\frac{K_{2,n}}{n^2} \right)
\end{aligned}$$

where the (almost sure) order of magnitude above is calculated using Lemma OA-13 and Assumptions 2(i), 5, and 6. By the triangle inequality, the conditional version of Liapunov's inequality,

and Assumption 5(ii); we then obtain

$$\begin{aligned}
E[|\mathcal{A}_{1,4}| |\mathcal{F}_n^Z] &\leq E[|\mathcal{A}_{1,4,1}| |\mathcal{F}_n^Z] + E[|\mathcal{A}_{1,4,2}| |\mathcal{F}_n^Z] \\
&\leq E[|\mathcal{A}_{1,4,1}| |\mathcal{F}_n^Z] + \sqrt{E[\mathcal{A}_{1,4,2}^2 |\mathcal{F}_n^Z]} \\
&= O_{a.s.}\left(\frac{K_{2,n}}{n}\right) + O_{a.s.}\left(\frac{\sqrt{K_{2,n}}}{n}\right) = O_{a.s.}\left(\frac{K_{2,n}}{n}\right) = o_{a.s.}(1).
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E\left[\frac{n}{K_{2,n}} |\mathcal{A}_{1,4}|\right] = E_Z\left(\frac{n}{K_{2,n}} E[|\mathcal{A}_{1,4}| |\mathcal{F}_n^Z]\right) \leq \bar{C}.$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\Pr\left(\left|\frac{n}{K_{2,n}} \mathcal{A}_{1,4}\right| \geq \frac{\bar{C}}{\epsilon}\right) \leq \epsilon \frac{n}{K_{2,n}} \frac{E[|\mathcal{A}_{1,4}|]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{1,4} = O_p\left(\frac{K_{2,n}}{n}\right). \quad (66)$$

Finally to show part (d), note that we can decompose \mathcal{A}_1 as

$$\begin{aligned}
&\mathcal{A}_1 \\
&= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} M^{Q_\varepsilon}\right)^2 \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^{Q_\varepsilon}\right)^2 \\
&\quad - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 \\
&= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} \sum_{r=1}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \varepsilon_{(i,g)} \varepsilon_{(i,r)} \\
&\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{q=1}^{T_j} \sum_{c=1}^{T_j} M_{(j,v),(j,q)}^Q M_{(j,v),(j,c)}^Q \varepsilon_{(j,q)} \varepsilon_{(j,c)} - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 \\
&= \mathcal{A}_{1,1} + \mathcal{A}_{1,2} + \mathcal{A}_{1,3} + \mathcal{A}_{1,4}
\end{aligned}$$

where

$$\begin{aligned} \mathcal{A}_{1,1} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{g=1}^{T_i} \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \varepsilon_{(i,g)}^2 \sum_{q=1}^{T_j} \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(M_{(j,v),(j,q)}^Q \right)^2 \varepsilon_{(j,q)}^2 \\ &\quad - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 \end{aligned}$$

and where $\mathcal{A}_{1,2}$, $\mathcal{A}_{1,3}$, and $\mathcal{A}_{1,4}$ are as defined in parts (a)-(c) above. We will first show that $\mathcal{A}_{1,1} = 0$. To proceed, note first that it is easily seen that $M^Q \circ M^Q$ is a block diagonal matrix, which can be written as

$$M^Q \circ M^Q = \begin{pmatrix} (M^Q \circ M^Q)_1 & 0 & \cdots & 0 \\ 0 & (M^Q \circ M^Q)_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & (M^Q \circ M^Q)_n \end{pmatrix}$$

where the i^{th} diagonal matrix $(M^Q \circ M^Q)_i$ is $T_i \times T_i$. Moreover, by definition $J = (M^Q \circ M^Q)^{-1}$, where the inverse exists in light of Assumption 6. Hence, J can also be written in block diagonal form as

$$J = \begin{pmatrix} J_1 & 0 & \cdots & 0 \\ 0 & J_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & J_n \end{pmatrix} = \begin{pmatrix} (M^Q \circ M^Q)_1^{-1} & 0 & \cdots & 0 \\ 0 & (M^Q \circ M^Q)_2^{-1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & (M^Q \circ M^Q)_n^{-1} \end{pmatrix}$$

Let $e_{(i,t)}$ be an $m_n \times 1$ elementary vector whose $(i,t)^{th}$ component is one and all other components are zero, and let e_{h,T_i} be a $T_i \times 1$ elementary vector, whose h^{th} component equals one and all other components equal to zero. Using these notations, one can show by direct calculation that, for any

$t, g \in \{1, \dots, T_i\}$ and for any $i \in \{1, 2, \dots, n\}$,

$$\begin{aligned}
\sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 &= \sum_{h=1}^{T_i} e'_{(i,t)} J e_{(i,h)} e'_{(i,h)} [M^Q \circ M^Q] e_{(i,g)} \\
&= \sum_{h=1}^{T_i} e'_{t,T_i} J_i e_{h,T_i} e'_{h,T_i} (M^Q \circ M^Q)_i e_{g,T_i} \\
&= e'_{t,T_i} (M^Q \circ M^Q)_i^{-1} \sum_{h=1}^{T_i} e_{h,T_i} e'_{h,T_i} (M^Q \circ M^Q)_i e_{g,T_i} \\
&= e'_{t,T_i} (M^Q \circ M^Q)_i^{-1} (M^Q \circ M^Q)_i e_{g,T_i} \\
&= e'_{t,T_i} I_{T_i} e_{g,T_i} \\
&= \begin{cases} 1 & \text{if } t = g \\ 0 & \text{if } t \neq g \end{cases}
\end{aligned}$$

These calculations imply that

$$\begin{aligned}
&\mathcal{A}_{1,1} \\
&= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{g,h=1}^{T_i} J_{(i,t),(i,h)} \left(M_{(i,h),(i,g)}^Q \right)^2 \varepsilon_{(i,g)}^2 \sum_{q,v=1}^{T_j} J_{(j,s),(j,v)} \left(M_{(j,v),(j,q)}^Q \right)^2 \varepsilon_{(j,q)}^2 \\
&\quad - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 \\
&= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 - \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \varepsilon_{(i,t)}^2 \varepsilon_{(j,s)}^2 \\
&= 0. \tag{67}
\end{aligned}$$

Combining (67) with the results obtained in expressions (64)-(66) above and making use of Assumption 5(ii), we further obtain

$$\begin{aligned}
\mathcal{A}_1 &= \mathcal{A}_{1,1} + \mathcal{A}_{1,2} + \mathcal{A}_{1,3} + \mathcal{A}_{1,4} \\
&= 0 + O_p \left(\frac{1}{\sqrt{n}} \right) + O_p \left(\frac{1}{\sqrt{n}} \right) + O_p \left(\frac{K_{2,n}}{n} \right) = O_p \left(\frac{1}{\sqrt{n}} \right). \square
\end{aligned}$$

Lemma OA-15: Let Assumptions 1-6 be satisfied, and let $\{\hat{\delta}_n\}$ be a sequence of estimators such that

$$\left\| \hat{\delta}_n - \delta_0 \right\|_2 \xrightarrow{p} 0 \text{ as } n \rightarrow \infty,$$

as long as $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Then, the following statements are true.

(a)

$$\mathcal{A}_{2,1} = O_p \left(\frac{K_n}{n} \right) = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{2,1} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right\} \end{aligned}$$

(b)

$$\mathcal{A}_{2,2} = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{2,2} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^{(Z,Q)} X [\hat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right\} \end{aligned}$$

(c)

$$\mathcal{A}_{2,3} = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{2,3} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} M^{(Z,Q)} X [\hat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right\} \end{aligned}$$

(d)

$$\mathcal{A}_{2,4} = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{2,4} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right\} \end{aligned}$$

$$(e) \quad \mathcal{A}_2 = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_2 &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \\ &\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \end{aligned}$$

Proof of Lemma OA-15:

To show part (a), we first apply the triangle inequality and the inequality $|XY| \leq (1/2)X^2 + (1/2)Y^2$ to obtain

$$\begin{aligned} |\mathcal{A}_{2,1}| &\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right| \right\} \\ &\leq \mathcal{A}_{2,1,1} + \mathcal{A}_{2,1,2} \end{aligned}$$

where

$$\begin{aligned} \mathcal{A}_{2,1,1} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \left(e'_{(j,v)} M^Q \varepsilon \right)^2, \\ \mathcal{A}_{2,1,2} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^Q \varepsilon \right)^2 \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right)^2. \end{aligned}$$

Clearly $\mathcal{A}_{2,1,1} \geq 0$ and $\mathcal{A}_{2,1,2} \geq 0$. Next, note that we can apply Assumptions 1, 2(i), 5, and 6 as well as part (a) of Lemma S2-1 to obtain

$$\begin{aligned}
& E[\mathcal{A}_{2,1,1} | \mathcal{F}_n^Z] \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{q=1}^{T_j} M_{(j,v),(j,q)}^Q \sum_{g=1}^{T_j} M_{(j,v),(j,g)}^Q \\
&\quad \times \sum_{(l,r)=1}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \sum_{(k,c)=1}^{m_n} P_{(i,h),(k,c)}^{Z^\perp} E[\varepsilon_{(l,r)} \varepsilon_{(k,c)} \varepsilon_{(j,q)} \varepsilon_{(j,g)} | \mathcal{F}_n^Z] \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \\
&\quad \times \sum_{q=1}^{T_j} \left(M_{(j,v),(j,q)}^Q \right)^2 \left(P_{(i,h),(j,q)}^{Z^\perp} \right)^2 E[\varepsilon_{(j,q)}^4 | \mathcal{F}_n^Z] \\
&\quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{q=1}^{T_j} \left(M_{(j,v),(j,q)}^Q \right)^2 \\
&\quad \times \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp} \right)^2 E[\varepsilon_{(l,r)}^2 | \mathcal{F}_n^Z] E[\varepsilon_{(j,q)}^2 | \mathcal{F}_n^Z] \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{q=1}^{T_j} \left| M_{(j,v),(j,q)}^Q \right| \left| P_{(i,h),(j,q)}^{Z^\perp} \right| \\
&\quad \times \sum_{g=1}^{T_j} \left| M_{(j,v),(j,g)}^Q \right| \left| P_{(i,h),(j,g)}^{Z^\perp} \right| E[\varepsilon_{(j,q)}^2 | \mathcal{F}_n^Z] E[\varepsilon_{(j,g)}^2 | \mathcal{F}_n^Z]
\end{aligned}$$

$$\begin{aligned}
&\leq \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \\
&\quad \times \frac{2\bar{T} \|J\|_\infty^2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 2\bar{T} \|J\|_\infty^2 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right)^2 \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 4\bar{T}^2 \|J\|_\infty^2 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right)^2 \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.} \left(\frac{K_n^2}{n^2} \right) + O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) = O_{a.s.} \left(\frac{K_n}{n} \right). \tag{68}
\end{aligned}$$

In the same way, we can also show that $E [\mathcal{A}_{2,1,2} | \mathcal{F}_n^Z] = O_{a.s.} (K_n/n)$, from which it follows

$$\begin{aligned}
E [|\mathcal{A}_{2,1}| | \mathcal{F}_n^Z] &\leq E [\mathcal{A}_{2,1,1} | \mathcal{F}_n^Z] + E [\mathcal{A}_{2,1,2} | \mathcal{F}_n^Z] \\
&= O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\frac{K_n}{n} \right) = O_{a.s.} \left(\frac{K_n}{n} \right).
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left[\frac{n}{K_n} |\mathcal{A}_{2,1}| \right] = E_Z \left(\frac{n}{K_n} E [|\mathcal{A}_{2,1}| | \mathcal{F}_n^Z] \right) \leq \bar{C}.$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\Pr \left(\left| \frac{n}{K_n} \mathcal{A}_{2,1} \right| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{n}{K_n} \frac{E [|\mathcal{A}_{2,1}|]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{2,1} = O_p \left(\frac{K_n}{n} \right) = o_p(1). \tag{69}$$

To show part (b), we again apply the triangle inequality and the inequality $|XY| \leq (1/2) X^2 +$

$(1/2) Y^2$ to get

$$\begin{aligned}
& |\mathcal{A}_{2,2}| \\
& \leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
& \quad \times \left| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^{Q_\varepsilon} \right) \left(e'_{(j,v)} M^{(Z,Q)} X \left[\widehat{\delta}_n - \delta_0 \right] \right) \left(e'_{(j,v)} M^{Q_\varepsilon} \right) \right| \\
& = \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
& \quad \times \left| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^{Q_\varepsilon} \right) \left(e'_{(j,v)} M^{(Z,Q)} U \left[\widehat{\delta}_n - \delta_0 \right] \right) \left(e'_{(j,v)} M^{Q_\varepsilon} \right) \right| \\
& \leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \\
& \quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^{Q_\varepsilon} \right)^2 \\
& \quad \times e'_{(j,v)} M^{(Z,Q)} U \left[\widehat{\delta}_n - \delta_0 \right] \left[\widehat{\delta}_n - \delta_0 \right]' U' M^{(Z,Q)} e_{(j,v)} \\
& \leq \mathcal{A}_{2,2,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{2,2,2},
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{A}_{2,2,1} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2, \\
\mathcal{A}_{2,2,2} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^{Q_\varepsilon} \right)^2 \\
&\quad \times e'_{(j,v)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(j,v)}
\end{aligned}$$

Note that $\mathcal{A}_{2,2,1}$ is the same as $\mathcal{A}_{2,1,1}$ above, so from expression (68), we get

$$0 \leq E [\mathcal{A}_{2,2,1} | \mathcal{F}_n^Z] = O_{a.s.} \left(\frac{K_n}{n} \right).$$

Moreover, using a similar argument as that given above to show expression (69) allows us to deduce that

$$\mathcal{A}_{2,2,1} = O_p \left(\frac{K_n}{n} \right) = o_p(1). \quad (70)$$

Next, making use of the decomposition $M^{(Z,Q)} = M^Q - P^{Z^\perp}$, with $P^{Z^\perp} = M^Q Z (Z' M^Q Z)^{-1} Z' M^Q$, we can obtain the inequality

$$\begin{aligned}
& e'_{(j,v)} M^{(Z,Q)} UU' M^{(Z,Q)} e_{(j,v)} \\
&= e'_{(j,v)} \left[M^Q - P^{Z^\perp} \right] UU' \left[M^Q - P^{Z^\perp} \right] e_{(j,v)} \\
&\leq e'_{(j,v)} M^Q UU' M^Q e_{(j,v)} + 2 \left| e'_{(j,v)} M^Q UU' P^{Z^\perp} e_{(j,v)} \right| + e'_{(j,v)} P^{Z^\perp} UU' P^{Z^\perp} e_{(j,v)} \\
&\leq 2e'_{(j,v)} M^Q UU' M^Q e_{(j,v)} + 2e'_{(j,v)} P^{Z^\perp} UU' P^{Z^\perp} e_{(j,v)} \\
&\quad \left(\text{since } \left| e'_{(j,v)} M^Q UU' P^{Z^\perp} e_{(j,v)} \right| \leq \frac{e'_{(j,v)} M^Q UU' M^Q e_{(j,v)}}{2} + \frac{e'_{(j,v)} P^{Z^\perp} UU' P^{Z^\perp} e_{(j,v)}}{2} \right) \tag{71}
\end{aligned}$$

Applying the inequality given in expression (71) above, we further obtain

$$\begin{aligned}
0 &\leq \mathcal{A}_{2,2,2} \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(j,v)} M^{(Z,Q)} UU' M^{(Z,Q)} e_{(j,v)} \\
&\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^Q \varepsilon \right)^2 e'_{(j,v)} M^Q UU' M^Q e_{(j,v)} \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^Q \varepsilon \right)^2 e'_{(j,v)} P^{Z^\perp} UU' P^{Z^\perp} e_{(j,v)} \\
&= \mathcal{A}_{2,2,2,1} + \mathcal{A}_{2,2,2,2}, \quad (\text{say})
\end{aligned}$$

Applying part (a) of Lemma S2-1 and Assumptions 2(i), 5, and 6; we get

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{2,2,2,1}|\mathcal{F}_n^Z] \\
&= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad \times \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| E\left[\left(e'_{(i,h)} M^Q \varepsilon\right)^2 e'_{(j,v)} M^Q UU' M^Q e_{(j,v)} |\mathcal{F}_n^Z\right] \\
&\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{g=1}^{T_i} \sum_{r=1}^{T_i} |M_{(i,h),(i,g)}^Q| |M_{(i,h),(i,r)}^Q| \\
&\quad \times \sum_{q=1}^{T_j} \sum_{c=1}^{T_j} |M_{(j,v),(j,q)}^Q| |M_{(j,v),(j,c)}^Q| E\left[|\varepsilon_{(i,g)} \varepsilon_{(i,r)} U'_{(j,q)} U_{(j,c)}| |\mathcal{F}_n^Z\right] \\
&\leq 4\bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q|\right)^4 \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^4 |\mathcal{F}_n^Z]\right)^{1/2} \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E[\|U_{(i,t)}\|_2^4 |\mathcal{F}_n^Z]\right)^{1/2} \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.}(1).
\end{aligned}$$

In addition, for $E[\mathcal{A}_{2,2,2,2}|\mathcal{F}_n^Z]$, we have, by straightforward calculations using Assumption 1 and

the triangle inequality,

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{2,2,2,2}|\mathcal{F}_n^Z] \\
&= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| E\left[\left(e'_{(i,h)} M^Q \varepsilon\right)^2 e'_{(j,v)} P^{Z^\perp} UU' P^{Z^\perp} e_{(j,v)} |\mathcal{F}_n^Z\right] \\
&= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{g=1}^{T_i} \sum_{r=1}^{T_i} M_{(i,h),(i,g)}^Q M_{(i,h),(i,r)}^Q \\
&\quad \times \sum_{(k,q)=1}^{m_n} P_{(j,v),(k,q)}^{Z^\perp} \sum_{(l,c)=1}^{m_n} P_{(j,v),(l,c)}^{Z^\perp} E\left[\varepsilon_{(i,g)} \varepsilon_{(i,r)} U'_{(k,q)} U_{(l,c)} |\mathcal{F}_n^Z\right] \\
&\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \right. \\
&\quad \times \left. \sum_{g=1}^{T_i} \left(M_{(i,h),(i,g)}^Q\right)^2 \left(P_{(j,v),(i,g)}^{Z^\perp}\right)^2 E\left[\varepsilon_{(i,g)}^2 U'_{(i,g)} U_{(i,g)} |\mathcal{F}_n^Z\right] \right\} \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{g=1}^{T_i} \left(M_{(i,h),(i,g)}^Q\right)^2 \right. \\
&\quad \times \left. \sum_{(k,q)=1}^{m_n} \left(P_{(j,v),(k,q)}^{Z^\perp}\right)^2 E\left[\varepsilon_{(i,g)}^2 |\mathcal{F}_n^Z\right] E\left[U'_{(k,q)} U_{(k,q)} |\mathcal{F}_n^Z\right] \right\} \\
&\quad + \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{g=1}^{T_i} \sum_{r=1}^{T_i} \left|M_{(i,h),(i,g)}^Q\right| \left|M_{(i,h),(i,r)}^Q\right| \right. \\
&\quad \times \left. \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left|P_{(j,v),(i,g)}^{Z^\perp}\right| \left|P_{(j,v),(i,r)}^{Z^\perp}\right| E\left[\left|\varepsilon_{(i,g)} \varepsilon_{(i,r)} U'_{(i,g)} U_{(i,r)}\right| |\mathcal{F}_n^Z\right] \right\} \tag{72}
\end{aligned}$$

Applying the CS inequality to (72) and making use of part (a) of Lemma S2-1 as well as Assumptions

2(i), 5, and 6; we then obtain

$$\begin{aligned}
& E [\mathcal{A}_{2,2,2,2} | \mathcal{F}_n^Z] \\
& \leq 4\bar{T} \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^{1/2} \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
& \quad + 4\bar{T} \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
& \quad + 8\bar{T}^2 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^{1/2} \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
& = O_{a.s.} \left(\frac{K_n^2}{n^2} \right) + O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) = O_{a.s.} \left(\frac{K_n}{n} \right) = o_{a.s.}(1).
\end{aligned}$$

It follows from these calculations that

$$\begin{aligned}
0 & \leq E [\mathcal{A}_{2,2,2}] \\
& = E [\mathcal{A}_{2,2,2,1}] + E [\mathcal{A}_{2,2,2,2}] \\
& = O_{a.s.}(1) + O_{a.s.} \left(\frac{K_n}{n} \right) \\
& = O_{a.s.}(1)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E [|\mathcal{A}_{2,2,2}|] = E [\mathcal{A}_{2,2,2}] = E_Z (E [\mathcal{A}_{2,2,2} | \mathcal{F}_n^Z]) \leq \bar{C}.$$

Application of the Markov's inequality then implies that, for any $\epsilon > 0$,

$$\Pr \left(|\mathcal{A}_{2,2,2}| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{E [\mathcal{A}_{2,2,2}]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{2,2,2} = O_p(1). \tag{73}$$

Combining the results on $\mathcal{A}_{2,2,1}$ and $\mathcal{A}_{2,2,2}$, we have

$$\begin{aligned} |\mathcal{A}_{2,2}| &\leq \mathcal{A}_{2,2,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{2,2,2} \\ &= O_p \left(\frac{K_n}{n} \right) + o_p(1) O_p(1) = o_p(1). \end{aligned} \quad (74)$$

Next, to show part (c), note that since A is symmetric, so that $A_{(i,t),(j,s)} = A_{(j,s),(i,t)}$, it follows that $\mathcal{A}_{2,2} = \mathcal{A}_{2,3}$. Hence, using the same argument as that given for part (b) above, we can show that

$$\mathcal{A}_{2,3} = o_p(1). \quad (75)$$

Turning our attention to part (d), note that, applying the triangle inequality and the inequality that $|XY| \leq (1/2)X^2 + (1/2)Y^2$, we obtain

$$\begin{aligned} &|\mathcal{A}_{2,4}| \\ &\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\ &\quad \times \left| \left(e'_{(i,h)} M^{(Z,Q)} X \left[\widehat{\delta}_n - \delta_0 \right] \right) \left(e'_{(i,h)} M^{Q,\varepsilon} \right) \left(e'_{(j,v)} M^{(Z,Q)} X \left[\widehat{\delta}_n - \delta_0 \right] \right) \left(e'_{(j,v)} M^{Q,\varepsilon} \right) \right| \\ &\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^{Q,\varepsilon} \right)^2 \\ &\quad \times e'_{(j,v)} M^{(Z,Q)} X \left[\widehat{\delta}_n - \delta_0 \right] \left[\widehat{\delta}_n - \delta_0 \right]' X' M^{(Z,Q)} e_{(j,v)} \\ &\quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{Q,\varepsilon} \right)^2 \\ &\quad \times e'_{(i,h)} M^{(Z,Q)} X \left[\widehat{\delta}_n - \delta_0 \right] \left[\widehat{\delta}_n - \delta_0 \right]' X' M^{(Z,Q)} e_{(i,h)} \\ &\leq \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 (\mathcal{A}_{2,4,1} + \mathcal{A}_{2,4,2}), \end{aligned}$$

where

$$\begin{aligned}
\mathcal{A}_{2,4,1} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(j,v)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(j,v)} \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(j,v)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(j,v)} \\
\mathcal{A}_{2,4,2} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(i,h)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(i,h)}
\end{aligned}$$

Note that both $\mathcal{A}_{2,4,1}$ and $\mathcal{A}_{2,4,2}$ are of the same form as $\mathcal{A}_{2,2,2}$, so that the result given in expression (73) allows us to deduce that

$$\mathcal{A}_{2,4,1} = O_p(1) \text{ and } \mathcal{A}_{2,4,2} = O_p(1),$$

from which it further follows that

$$|\mathcal{A}_{2,4}| \leq \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 (\mathcal{A}_{2,4,1} + \mathcal{A}_{2,4,2}) = o_p(1) [O_p(1) + O_p(1)] = o_p(1). \quad (76)$$

Finally, for part (e), note that we can decompose \mathcal{A}_2 as

$$\begin{aligned}
\mathcal{A}_2 &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right. \\
&\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right\} \\
&= \mathcal{A}_{2,1} + \mathcal{A}_{2,2} + \mathcal{A}_{2,3} + \mathcal{A}_{2,4},
\end{aligned}$$

where $\mathcal{A}_{2,1}$, $\mathcal{A}_{2,2}$, $\mathcal{A}_{2,3}$, and $\mathcal{A}_{2,4}$ are as defined in parts (a)-(d) above. It follows from the results

given in expressions (69), (74), (75), and (76) that

$$\begin{aligned}\mathcal{A}_2 &= \mathcal{A}_{2,1} + \mathcal{A}_{2,2} + \mathcal{A}_{2,3} + \mathcal{A}_{2,4} \\ &= O_p\left(\frac{K_n}{n}\right) + o_p(1) + o_p(1) + o_p(1) \\ &= o_p(1). \quad \square\end{aligned}$$

Lemma OA-16: Let Assumptions 1-6 be satisfied, and let $\{\hat{\delta}_n\}$ be a sequence of estimators such that

$$\|\hat{\delta}_n - \delta_0\|_2 \xrightarrow{p} 0 \text{ as } n \rightarrow \infty,$$

as long as $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Then, the following statements are true.

(a)

$$\mathcal{A}_{3,1} = O_p\left(\frac{K_n^2}{n^2}\right) = o_p(1),$$

where

$$\begin{aligned}\mathcal{A}_{3,1} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon\right)^2 \left(e'_{(j,v)} P^{Z^\perp} \varepsilon\right)^2\end{aligned}$$

(b)

$$\mathcal{A}_{3,2} = O_p\left(\frac{K_n}{n}\right) = o_p(1),$$

where

$$\begin{aligned}\mathcal{A}_{3,2} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon\right)^2 \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| e'_{(j,v)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(j,v)} \right\}\end{aligned}$$

(c)

$$\mathcal{A}_{3,3} = O_p\left(\frac{K_n}{n}\right) = o_p(1).$$

where

$$\begin{aligned} \mathcal{A}_{3,3} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \right. \\ &\quad \times \left. \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right)^2 \right\} \end{aligned}$$

(d)

$$\mathcal{A}_{3,4} = O_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{3,4} &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \right. \\ &\quad \times \left. \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| e'_{(j,v)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(j,v)} \right\} \end{aligned}$$

(e)

$$\mathcal{A}_3 = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_3 &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\hat{\delta}_n - \delta_0] \right]^2 \right. \\ &\quad \times \left. \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left[e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\hat{\delta}_n - \delta_0] \right]^2 \right\} \end{aligned}$$

Proof of Lemma OA-16:

To show part (a), note that, by applying part (a) of Lemma S2-1 and Assumptions 1, 2(i), 5,

and 6; we have

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{3,1}|\mathcal{F}_n^Z] \\
&= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| E\left[\left(e'_{(i,h)} P^{Z^\perp} \varepsilon\right)^2 \left(e'_{(j,v)} P^{Z^\perp} \varepsilon\right)^2 |\mathcal{F}_n^Z\right] \\
&= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(l,r)=1}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \sum_{(k,c)=1}^{m_n} P_{(i,h),(k,c)}^{Z^\perp} \\
&\quad \times \sum_{(p,q)=1}^{m_n} P_{(j,v),(p,q)}^{Z^\perp} \sum_{(g,w)=1}^{m_n} P_{(j,v),(g,w)}^{Z^\perp} E[\varepsilon_{(l,r)} \varepsilon_{(k,c)} \varepsilon_{(p,q)} \varepsilon_{(g,w)} |\mathcal{F}_n^Z] \\
&\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
&\quad \times \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp}\right)^2 \left(P_{(j,v),(l,r)}^{Z^\perp}\right)^2 E[\varepsilon_{(l,r)}^4 |\mathcal{F}_n^Z] \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp}\right)^2 \\
&\quad \times \sum_{(p,q)=1}^{m_n} \left(P_{(j,v),(p,q)}^{Z^\perp}\right)^2 E[\varepsilon_{(l,r)}^2 |\mathcal{F}_n^Z] E[\varepsilon_{(p,q)}^2 |\mathcal{F}_n^Z] \\
&\quad + \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(l,r)=1}^{m_n} \left|P_{(i,h),(l,r)}^{Z^\perp}\right| \left|P_{(j,v),(l,r)}^{Z^\perp}\right| \\
&\quad \times \sum_{(k,c)=1}^{m_n} \left|P_{(i,h),(k,c)}^{Z^\perp}\right| \left|P_{(j,v),(k,c)}^{Z^\perp}\right| E[\varepsilon_{(l,r)}^2 |\mathcal{F}_n^Z] E[\varepsilon_{(p,q)}^2 |\mathcal{F}_n^Z]
\end{aligned}$$

$$\begin{aligned}
&\leq 4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^3 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 8 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.} \left(\frac{K_n^3}{n^3} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) \\
&= O_{a.s.} \left(\frac{K_n^2}{n^2} \right).
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left[\frac{n^2}{K_n^2} |\mathcal{A}_{3,1}| \right] = E \left[\frac{n^2}{K_n^2} \mathcal{A}_{3,1} \right] = E_Z \left(E \left[\frac{n^2}{K_n^2} \mathcal{A}_{3,1} | \mathcal{F}_n^Z \right] \right) \leq \bar{C}.$$

Application of the Markov's inequality then implies that, for any $\epsilon > 0$,

$$\Pr \left(\frac{n^2}{K_n^2} |\mathcal{A}_{3,1}| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{n^2}{K_n^2} \frac{E[\mathcal{A}_{3,1}]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{3,1} = O_p \left(\frac{K_n^2}{n^2} \right). \tag{77}$$

To show part (b), first let $e_{b,d}$ denote a $d \times 1$ elementary vector whose b^{th} component equals one for $b \in \{1, \dots, d\}$ and whose other components all equal to zero. Using the fact that

$$M^{(Z,Q)} X = M^Q U - P^{Z^\perp} U,$$

we can apply the inequality $\left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r$ to obtain

$$\begin{aligned}
0 &\leq \mathcal{A}_{3,2} \\
&= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 e'_{(j,v)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(j,v)} \\
&= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \\
&\quad \times \sum_{b=1}^d \left(e'_{(j,v)} M^Q U e_{b,d} - e'_{(j,v)} P^{Z^\perp} U e_{b,d} \right)^2 \\
&\leq \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \sum_{b=1}^d \left(e'_{(j,v)} M^Q U e_{b,d} \right)^2 \\
&\quad + \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \sum_{b=1}^d \left(e'_{(j,v)} P^{Z^\perp} U e_{b,d} \right)^2 \\
&= \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 e'_{(j,v)} M^Q U U' M^Q e_{(j,v)} \\
&\quad + \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 e'_{(j,v)} P^{Z^\perp} U U' P^{Z^\perp} e_{(j,v)} \\
&= \mathcal{A}_{3,2,1} + \mathcal{A}_{3,2,2}
\end{aligned}$$

Making use of part (a) of Lemma S2-1 and Assumptions 1, 2(i), 5, and 6; we obtain

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{3,2,1}|\mathcal{F}_n^Z] \\
&= \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| E[e'_{(j,v)} M^Q U U' M^Q e_{(j,v)} e'_{(i,h)} P^{Z^\perp} \varepsilon \varepsilon' P^{Z^\perp} e_{(i,h)} |\mathcal{F}_n^Z] \\
&= \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{q=1}^{T_j} M_{(j,v),(j,q)}^Q \sum_{g=1}^{T_j} M_{(j,v),(j,g)}^Q \\
&\quad \times \sum_{(l,r)=1}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \sum_{(k,c)=1}^{m_n} P_{(i,h),(k,c)}^{Z^\perp} E[U'_{(j,q)} U_{(j,g)} \varepsilon_{(l,r)} \varepsilon_{(k,c)} |\mathcal{F}_n^Z] \\
&\leq \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
&\quad \times \sum_{q=1}^{T_j} \left(M_{(j,v),(j,q)}^Q \right)^2 \left(P_{(i,h),(j,q)}^{Z^\perp} \right)^2 E[U'_{(j,q)} U_{(j,q)} \varepsilon_{(j,q)}^2 |\mathcal{F}_n^Z] \\
&\quad + \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{q=1}^{T_j} \left(M_{(j,v),(j,q)}^Q \right)^2 \\
&\quad \times \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp} \right)^2 E[U'_{(j,q)} U_{(j,q)} |\mathcal{F}_n^Z] E[\varepsilon_{(l,r)}^2 |\mathcal{F}_n^Z] \\
&\quad + \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{q=1}^{T_j} \left| M_{(j,v),(j,q)}^Q \right| \left| P_{(i,h),(j,q)}^{Z^\perp} \right| \\
&\quad \times \sum_{g=1}^{T_j} \left| M_{(j,v),(j,g)}^Q \right| \left| P_{(i,h),(j,g)}^{Z^\perp} \right| E[\left| U'_{(j,q)} U_{(j,g)} \varepsilon_{(j,q)} \varepsilon_{(j,g)} \right| |\mathcal{F}_n^Z]
\end{aligned}$$

$$\begin{aligned}
&\leq 8\bar{T}\|J\|_\infty^2 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^{1/2} \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E[\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^W] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 8\bar{T}\|J\|_\infty^2 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^W] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 16\bar{T}^2\|J\|_\infty^2 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^{1/2} \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E[\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^W] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.}\left(\frac{K_n^2}{n^2}\right) + O_{a.s.}\left(\frac{K_n}{n}\right) + O_{a.s.}\left(\frac{K_n^2}{n^2}\right) = O_{a.s.}\left(\frac{K_n}{n}\right)
\end{aligned}$$

Finally, applying part (a) of Lemma S2-1 and Assumptions 1, 2(i), 5, and 6; we get

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{3,2,3}|\mathcal{F}_n^Z] \\
&= \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| E[e'_{(j,v)} P^{Z^\perp} UU' P^{Z^\perp} e_{(j,v)} e'_{(i,h)} P^{Z^\perp} \varepsilon \varepsilon' P^{Z^\perp} e_{(i,h)} |\mathcal{F}_n^Z] \\
&= \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(p,q)=1}^{m_n} P_{(j,v),(p,q)}^{Z^\perp} \sum_{(f,g)=1}^{m_n} P_{(j,v),(f,g)}^{Z^\perp} \\
&\quad \times \sum_{(l,r)=1}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \sum_{(k,c)=1}^{m_n} P_{(i,h),(k,c)}^{Z^\perp} E[U'_{(p,q)} U_{(f,g)} \varepsilon_{(l,r)} \varepsilon_{(k,c)} |\mathcal{F}_n^Z] \\
&\leq \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
&\quad \times \sum_{(l,r)=1}^{m_n} \left(P_{(j,v),(l,r)}^{Z^\perp} \right)^2 \left(P_{(i,h),(l,r)}^{Z^\perp} \right)^2 E[U'_{(l,r)} U_{(l,r)} \varepsilon_{(l,r)}^2 |\mathcal{F}_n^Z] \\
&\quad + \frac{8}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(p,q)=1}^{m_n} \left(P_{(j,v),(p,q)}^{Z^\perp} \right)^2 \\
&\quad \times \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp} \right)^2 E[U'_{(p,q)} U_{(p,q)} |\mathcal{F}_n^W] E[\varepsilon_{(l,r)}^2 |\mathcal{F}_n^Z] \\
&\quad + \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(p,q)=1}^{m_n} \left| P_{(j,v),(l,r)}^{Z^\perp} P_{(i,h),(l,r)}^{Z^\perp} \right| \\
&\quad \times \sum_{(k,c)=1}^{m_n} \left| P_{(j,v),(k,c)}^{Z^\perp} P_{(i,h),(k,c)}^{Z^\perp} \right| E[\left| U'_{(l,r)} U_{(k,c)} \varepsilon_{(l,r)} \varepsilon_{(k,c)} \right| |\mathcal{F}_n^Z]
\end{aligned}$$

$$\begin{aligned}
&\leq 8 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z \right] \right)^{1/2} \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^3 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 8 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z \right] \right) \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right) \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 16 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z \right] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z \right] \right)^{1/2} \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.} \left(\frac{K_n^3}{n^3} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) = O_{a.s.} \left(\frac{K_n^2}{n^2} \right)
\end{aligned}$$

It follows from these calculations that

$$\begin{aligned}
0 &\leq E [\mathcal{A}_{3,2} | \mathcal{F}_n^Z] \\
&= E [\mathcal{A}_{3,2,1} | \mathcal{F}_n^Z] + E [\mathcal{A}_{3,2,2} | \mathcal{F}_n^Z] \\
&= O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) \\
&= O_{a.s.} \left(\frac{K_n}{n} \right)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left[\frac{n}{K_n} |\mathcal{A}_{3,2}| \right] = E \left[\frac{n}{K_n} \mathcal{A}_{3,2} \right] = E_Z \left(E \left[\frac{n}{K_n} \mathcal{A}_{3,2} | \mathcal{F}_n^Z \right] \right) \leq \bar{C}.$$

Application of the Markov's inequality then implies that, for any $\epsilon > 0$,

$$\Pr \left(\frac{n}{K_n} |\mathcal{A}_{3,2}| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{n}{K_n} \frac{E[\mathcal{A}_{3,2}]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{3,2} = O_p \left(\frac{K_n}{n} \right). \tag{78}$$

Next, to show part (c), note that since A is symmetric, so that $A_{(i,t),(j,s)} = A_{(j,s),(i,t)}$, it follows that $\mathcal{A}_{3,2} = \mathcal{A}_{3,3}$. Hence, using the same argument as that given for part (b) above, we can show

that

$$\mathcal{A}_{3,3} = O_p \left(\frac{K_n}{n} \right). \quad (79)$$

Turning our attention to part (d), we apply the inequality $|XY| \leq (1/2)X^2 + (1/2)Y^2$ to obtain

$$\begin{aligned} 0 &\leq \mathcal{A}_{3,4} \\ &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \right. \\ &\quad \times \left. \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| e'_{(j,v)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(j,v)} \right\} \\ &= \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(i,h)} \right. \\ &\quad \times \left. \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| e'_{(j,v)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(j,v)} \right\} \\ &\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(i,h)} \right)^2 \\ &\quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(j,v)} \right)^2 \\ &= \mathcal{A}_{3,4,1} + \mathcal{A}_{3,4,2} \end{aligned}$$

Focusing first on $\mathcal{A}_{3,4,1}$, note that by applying the inequality $\left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r$ and

following an argument similar to that given for $\mathcal{A}_{3,2}$ above, we have

$$\begin{aligned}
0 &\leq \mathcal{A}_{3,4,1} \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^{(Z,Q)} UU' M^{(Z,Q)} e_{(i,h)} \right)^2 \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left[\sum_{b=1}^d \left(e'_{(i,h)} M^Q U e_{b,d} - e'_{(i,h)} P^{Z^\perp} U e_{b,d} \right)^2 \right]^2 \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
&\quad \times \left[2e'_{(i,h)} M^Q UU' M^Q e_{(i,h)} + 2e'_{(i,h)} P^{Z^\perp} UU' P^{Z^\perp} e_{(i,h)} \right]^2 \\
&\leq \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} M^Q UU' M^Q e_{(i,h)} \right)^2 \\
&\quad + \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} UU' P^{Z^\perp} e_{(i,h)} \right)^2 \\
&= \mathcal{A}_{3,4,1,1} + \mathcal{A}_{3,4,1,2}
\end{aligned}$$

Applying the triangle inequality, part (a) of Lemma S2-1, as well as Assumptions 2(i), 5, and 6; we

obtain

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{3,4,1,1}|\mathcal{F}_n^Z] \\
&= \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| E\left[\left(e'_{(i,h)} M^Q U U' M^Q e_{(i,h)}\right)^2 |\mathcal{F}_n^Z\right] \\
&\leq \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{g=1}^{T_i} |M_{(i,h),(i,g)}^Q| \sum_{r=1}^{T_i} |M_{(i,h),(i,r)}^Q| \\
&\quad \times \sum_{q=1}^{T_i} |M_{(i,h),(i,q)}^Q| \sum_{r=1}^{T_i} |M_{(i,h),(i,r)}^Q| E\left[\left|U'_{(i,g)} U_{(i,r)} U'_{(i,q)} U_{(i,c)}\right| |\mathcal{F}_n^Z\right] \\
&\leq 16\bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q|\right)^4 \left(\max_{1 \leq (i,t) \leq m_n} E\left[\|U_{(i,t)}\|_2^4 |\mathcal{F}_n^Z\right]\right) \\
&\quad \times \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.}(1)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E[|\mathcal{A}_{3,4,1,1}|] = E[\mathcal{A}_{3,4,1,1}] = E_{W_n}(E[\mathcal{A}_{3,4,1,1}|\mathcal{F}_n^W]) \leq \bar{C}.$$

Application of the Markov's inequality then implies that, for any $\epsilon > 0$,

$$\Pr\left(|\mathcal{A}_{3,4,1,1}| \geq \frac{\bar{C}}{\epsilon}\right) \leq \epsilon \frac{E[\mathcal{A}_{3,4,1,1}]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{3,4,1,1} = O_p(1).$$

Moreover, by applying part (a) of Lemma S2-1 as well as Assumptions 1, 2(i), 5, and 6; we have

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{3,4,1,2}|\mathcal{F}_n^Z] \\
&= \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} UU' P^{Z^\perp} e_{(i,h)} \right)^2 \\
&= \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(l,r)=1}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \sum_{(k,c)=1}^{m_n} P_{(i,h),(k,c)}^{Z^\perp} \\
&\quad \times \sum_{(p,q)=1}^{m_n} P_{(i,h),(p,q)}^{Z^\perp} \sum_{(k,c)=1}^{m_n} P_{(i,h),(f,g)}^{Z^\perp} E[U'_{(l,r)} U_{(k,c)} U'_{(p,q)} U_{(f,g)} |\mathcal{F}_n^Z] \\
&= \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp} \right)^4 E \left[\left(U'_{(l,r)} U_{(l,r)} \right)^2 |\mathcal{F}_n^Z \right] \\
&\quad + \frac{16}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp} \right)^2 \\
&\quad \times \sum_{\substack{(p,q)=1 \\ (p,q)\neq(l,r)}}^{m_n} \left(P_{(i,h),(p,q)}^{Z^\perp} \right)^2 E \left[\left(U'_{(l,r)} U_{(l,r)} \right) |\mathcal{F}_n^Z \right] E \left[\left(U'_{(p,q)} U_{(p,q)} \right) |\mathcal{F}_n^Z \right] \\
&\quad + \frac{32}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{(l,r)=1}^{m_n} \left(P_{(i,h),(l,r)}^{Z^\perp} \right)^2 \\
&\quad \times \sum_{\substack{(p,q)=1 \\ (p,q)\neq(l,r)}}^{m_n} \left(P_{(i,h),(p,q)}^{Z^\perp} \right)^2 E \left[\left(U'_{(l,r)} U_{(p,q)} \right) |\mathcal{F}_n^Z \right] E \left[\left(U'_{(p,q)} U_{(l,r)} \right) |\mathcal{F}_n^Z \right]
\end{aligned}$$

$$\begin{aligned}
&\leq 16 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z \right] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^3 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 16 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \\
&\quad \times \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad + 32 \|J\|_\infty^2 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z \right] \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \\
&\quad \times \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.} \left(\frac{K_n^3}{n^3} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) = O_{a.s.} \left(\frac{K_n^2}{n^2} \right).
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left[\frac{n^2}{K_n^2} |\mathcal{A}_{3,4,1,2}| \right] = E \left[\frac{n^2}{K_n^2} \mathcal{A}_{3,4,1,2} \right] = E_Z \left(\frac{n^2}{K_n^2} E [\mathcal{A}_{3,4,1,2} | \mathcal{F}_n^Z] \right) \leq \bar{C}.$$

Application of the Markov's inequality then implies that, for any $\epsilon > 0$,

$$\Pr \left(\frac{n^2}{K_n^2} |\mathcal{A}_{3,4,1,2}| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{n^2}{K_n^2} \frac{E [\mathcal{A}_{3,4,1,2}]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that $\mathcal{A}_{3,4,1,2} = O_p(K_n^2/n^2)$. It follows that

$$\begin{aligned}
\mathcal{A}_{3,4,1} &\leq \mathcal{A}_{3,4,1,1} + \mathcal{A}_{3,4,1,2} \\
&= O_p(1) + O_p \left(\frac{K_n^2}{n^2} \right) = O_p(1)
\end{aligned}$$

In the same way, we can also show that $\mathcal{A}_{3,4,2} = O_p(1)$, from which it further follows that

$$0 \leq \mathcal{A}_{3,4} \leq \mathcal{A}_{3,4,1} + \mathcal{A}_{3,4,2} = O_p(1) + O_p(1) = O_p(1). \quad (80)$$

Finally, consider part (e). Making use of the triangle inequality and the inequality $\left| \sum_{i=1}^m a_i \right|^r \leq$

$m^{r-1} \sum_{i=1}^m |a_i|^r$, we can write

$$\begin{aligned}
|\mathcal{A}_3| &\leq \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right]^2 \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left[e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right]^2 \\
&\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right)^2 \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \times \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \\
&\quad \times e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] [\widehat{\delta}_n - \delta_0]' X' M^{(Z,Q)} e_{(j,v)} \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
&\quad \times \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right)^2 e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] [\widehat{\delta}_n - \delta_0]' X' M^{(Z,Q)} e_{(i,h)} \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \\
&\quad \times [\widehat{\delta}_n - \delta_0]' X' M^{(Z,Q)} e_{(i,h)} e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] [\widehat{\delta}_n - \delta_0]' X' M^{(Z,Q)} e_{(j,v)}
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right)^2 \\
&+ \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
&\quad \times \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 e'_{(j,v)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(j,v)} \\
&+ \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
&\quad \times \left(e'_{(j,v)} P^{Z^\perp} \varepsilon \right)^2 e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \\
&+ \left\| \widehat{\delta}_n - \delta_0 \right\|_2^4 \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \\
&\quad \times e'_{(j,v)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(j,v)} \\
&= \mathcal{A}_{3,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{3,2} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{3,3} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^4 \mathcal{A}_{3,4}, \tag{81}
\end{aligned}$$

where $\mathcal{A}_{3,1}$, $\mathcal{A}_{3,2}$, $\mathcal{A}_{3,3}$, and $\mathcal{A}_{3,4}$ are as defined in parts (a)-(d) above. It follows from expressions (77)-(80) that

$$\begin{aligned}
|\mathcal{A}_3| &\leq \mathcal{A}_{3,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{3,2} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{3,3} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^4 \mathcal{A}_{3,4} \\
&= O_p \left(\frac{K_n^2}{n^2} \right) + o_p(1) O_p \left(\frac{K_n}{n} \right) + o_p(1) O_p \left(\frac{K_n}{n} \right) + o_p(1) O_p(1) = o_p(1). \tag{82}
\end{aligned}$$

Lemma OA-17: Let Assumptions 1-6 be satisfied, and let $\{\widehat{\delta}_n\}$ be a sequence of estimators such that

$$\left\| \widehat{\delta}_n - \delta_0 \right\|_2 \xrightarrow{p} 0 \text{ as } n \rightarrow \infty,$$

as long as $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Then, the following statements are true.

(a)

$$\mathcal{A}_{4,1} = O_p \left(\sqrt{\frac{K_n}{n}} \right) = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{4,1} &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \right\} \end{aligned}$$

(b)

$$\mathcal{A}_{4,2} = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{4,2} &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} M^{(Z,Q)} X [\hat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \right\} \end{aligned}$$

(c)

$$\mathcal{A}_4 = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_4 &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \right. \\ &\quad \left. \times \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\hat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \right\} \end{aligned}$$

Proof of Lemma OA-17:

To show part (a), note first that we can decompose $\mathcal{A}_{4,1}$ as

$$\begin{aligned}
\mathcal{A}_{4,1} &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \\
&\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&= -\frac{2}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{s,t=1 \\ s \neq t}}^{T_i} A_{(i,t),(i,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{r=1}^{T_i} P_{(i,h),(i,r)}^{Z^\perp} \sum_{g=1}^{T_i} M_{(i,h),(i,g)}^Q \sum_{v=1}^{T_i} J_{(i,s),(i,v)} \\
&\quad \times \sum_{q=1}^{T_i} M_{(i,v),(i,q)}^Q \sum_{c=1}^{T_i} M_{(i,v),(i,c)}^Q \varepsilon(i,r) \varepsilon(i,g) \varepsilon(i,q) \varepsilon(i,c) \\
&\quad -\frac{2}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{s,t=1 \\ s \neq t}}^{T_i} A_{(i,t),(i,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{\substack{(l,r)=1 \\ l \neq i}}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \sum_{g=1}^{T_i} M_{(i,h),(i,g)}^Q \sum_{v=1}^{T_i} J_{(i,s),(i,v)} \\
&\quad \times \sum_{q=1}^{T_i} M_{(i,v),(i,q)}^Q \sum_{c=1}^{T_i} M_{(i,v),(i,c)}^Q \varepsilon(l,r) \varepsilon(i,g) \varepsilon(i,q) \varepsilon(i,c) \\
&\quad -\frac{2}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{r=1}^{T_i} P_{(i,h),(i,r)}^{Z^\perp} \sum_{g=1}^{T_i} M_{(i,h),(i,g)}^Q \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \\
&\quad \times \sum_{q=1}^{T_j} M_{(j,v),(j,q)}^Q \sum_{c=1}^{T_j} M_{(j,v),(j,c)}^Q \varepsilon(i,r) \varepsilon(i,g) \varepsilon(j,q) \varepsilon(j,c) \\
&\quad -\frac{2}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} M_{(i,h),(i,g)}^Q \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{r=1}^{T_i} P_{(i,h),(j,r)}^{Z^\perp} \\
&\quad \times \sum_{q=1}^{T_j} M_{(j,v),(j,q)}^Q \sum_{c=1}^{T_j} M_{(j,v),(j,c)}^Q \varepsilon(i,g) \varepsilon(j,r) \varepsilon(j,q) \varepsilon(j,c) \\
&\quad -\frac{2}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} M_{(i,h),(i,g)}^Q \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{(l,r)=1 \\ l \neq i,j}}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \\
&\quad \times \sum_{q=1}^{T_j} M_{(j,v),(j,q)}^Q \sum_{c=1}^{T_j} M_{(j,v),(j,c)}^Q \varepsilon(l,r) \varepsilon(i,g) \varepsilon(j,q) \varepsilon(j,c) \\
&= \mathcal{A}_{4,1,1} + \mathcal{A}_{4,1,2} + \mathcal{A}_{4,1,3} + \mathcal{A}_{4,1,4} + \mathcal{A}_{4,1,5}
\end{aligned}$$

Consider first $\mathcal{A}_{4,1,1}$. Applying the triangle inequality, part (f) of Lemma S2-1, and Assumptions

2(i), 5, and 6; we obtain

$$\begin{aligned}
& E \left[|\mathcal{A}_{4,1,1}| \mid \mathcal{F}_n^Z \right] \\
& \leq \frac{2}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{s,t=1 \\ s \neq t}}^{T_i} A_{(i,t),(i,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{r=1}^{T_i} \left| P_{(i,h),(i,r)}^{Z^\perp} \right| \sum_{g=1}^{T_i} \left| M_{(i,h),(i,g)}^Q \right| \sum_{v=1}^{T_i} |J_{(i,s),(i,v)}| \\
& \quad \times \sum_{q=1}^{T_i} \left| M_{(i,v),(i,q)}^Q \right| \sum_{c=1}^{T_i} \left| M_{(i,v),(i,c)}^Q \right| E \left[|\varepsilon_{(i,r)} \varepsilon_{(i,g)} \varepsilon_{(i,q)} \varepsilon_{(i,c)}| \mid \mathcal{F}_n^Z \right] \\
& \leq 2\bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} \left| M_{(i,t),(j,s)}^Q \right| \right)^3 \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 \mid \mathcal{F}_n^Z \right] \right) \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \frac{1}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{s,t=1 \\ s \neq t}}^{T_i} A_{(i,t),(i,s)}^2 \\
& = O_{a.s.} \left(\frac{K_n K_{2,n}}{n^2} \right)
\end{aligned}$$

Next, consider $\mathcal{A}_{4,1,1}$. Here, by applying the triangle inequality, parts (f) and (g) of Lemma S2-1,

and Assumptions 1, 2(i), 5, and 6; we get

$$\begin{aligned}
& E [\mathcal{A}_{4,1,2}^2 | \mathcal{F}_n^Z] \\
= & E \left[\left(\frac{2}{K_{2,n}} \sum_{i=1}^n \sum_{\substack{s,t=1 \\ s \neq t}}^{T_i} A_{(i,t),(i,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{\substack{(l,r)=1 \\ l \neq i}}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \sum_{g=1}^{T_i} M_{(i,h),(i,g)}^Q \sum_{v=1}^{T_i} J_{(i,s),(i,v)} \right. \right. \\
& \times \sum_{q=1}^{T_i} M_{(i,v),(i,q)}^Q \sum_{c=1}^{T_i} M_{(i,v),(i,c)}^Q \varepsilon_{(l,r)} \varepsilon_{(i,g)} \varepsilon_{(i,q)} \varepsilon_{(i,c)} \left. \left. \right)^2 | \mathcal{F}_n^Z \right] \\
\leq & \left\{ \frac{4}{K_{2,n}^2} \sum_{i=1}^n \sum_{\substack{s_1,t_1=1 \\ s_1 \neq t_1}}^{T_i} \sum_{\substack{s_2,t_2=1 \\ s_2 \neq t_2}}^{T_i} A_{(i,t_1),(i,s_1)}^2 A_{(i,t_2),(i,s_2)}^2 \sum_{h_1=1}^{T_i} |J_{(i,t_1),(i,h_1)}| \sum_{h_2=1}^{T_i} |J_{(i,t_2),(i,h_2)}| \right. \\
& \times \sum_{\substack{(l,r)=1 \\ l \neq i}}^{m_n} \left| P_{(i,h_1),(l,r)}^{Z^\perp} P_{(i,h_2),(l,r)}^{Z^\perp} \right| \sum_{g_1=1}^{T_i} \left| M_{(i,h_1),(i,g_1)}^Q \right| \sum_{g_2=1}^{T_i} \left| M_{(i,h_2),(i,g_2)}^Q \right| \sum_{v_1=1}^{T_i} |J_{(i,s_1),(i,v_1)}| \\
& \times \sum_{v_2=1}^{T_i} |J_{(i,s_2),(i,v_2)}| \sum_{q_1=1}^{T_i} \left| M_{(i,v_1),(i,q_1)}^Q \right| \sum_{q_2=1}^{T_i} \left| M_{(i,v_2),(i,q_2)}^Q \right| \sum_{c_1=1}^{T_i} \left| M_{(i,v_1),(i,c_1)}^Q \right| \sum_{c_2=1}^{T_i} \left| M_{(i,v_2),(i,c_2)}^Q \right| \\
& \times E \left[\varepsilon_{(l,r)}^2 | \mathcal{F}_n^Z \right] E \left[|\varepsilon_{(i,g_1)} \varepsilon_{(i,q_1)} \varepsilon_{(i,c_1)} \varepsilon_{(i,g_2)} \varepsilon_{(i,q_2)} \varepsilon_{(i,c_2)}| | \mathcal{F}_n^Z \right] \} \\
& + \left\{ \frac{4}{K_{2,n}^2} \sum_{i_1=1}^n \sum_{\substack{s_1,t_1=1 \\ s_1 \neq t_1}}^{T_{i_1}} A_{(i_1,t_1),(i_1,s_1)}^2 \sum_{i_2=1}^n \sum_{\substack{s_2,t_2=1 \\ i_2 \neq i_1 \\ s_2 \neq t_2}}^{T_{i_2}} A_{(i_2,t_2),(i_2,s_2)}^2 \sum_{h_1=1}^{T_{i_1}} |J_{(i_1,t_1),(i_1,h_1)}| \sum_{h_2=1}^{T_{i_2}} |J_{(i_2,t_2),(i_2,h_2)}| \right. \\
& \times \sum_{r_1=1}^{T_{i_1}} \left| P_{(i_1,h_1),(i_2,r_1)}^{Z^\perp} \right| \sum_{r_2=1}^{T_{i_2}} \left| P_{(i_2,h_2),(i_2,r_2)}^{Z^\perp} \right| \sum_{g_1=1}^{T_{i_1}} \left| M_{(i_1,h_1),(i_1,g_1)}^Q \right| \sum_{g_2=1}^{T_{i_2}} \left| M_{(i_2,h_2),(i_2,g_2)}^Q \right| \\
& \times \sum_{v_1=1}^{T_{i_1}} |J_{(i_1,s_1),(i_1,v_1)}| \sum_{v_2=1}^{T_{i_2}} |J_{(i_2,s_2),(i_2,v_2)}| \sum_{q_1=1}^{T_{i_1}} \left| M_{(i_1,v_1),(i_1,q_1)}^Q \right| \\
& \times \sum_{q_2=1}^{T_{i_2}} \left| M_{(i_2,v_2),(i_2,q_2)}^Q \right| \sum_{c_1=1}^{T_{i_1}} \left| M_{(i_1,v_1),(i_1,c_1)}^Q \right| \sum_{c_2=1}^{T_{i_2}} \left| M_{(i_2,v_2),(i_2,c_2)}^Q \right| \\
& \times E \left[|\varepsilon_{(i_1,r_1)} \varepsilon_{(i_1,g_1)} \varepsilon_{(i_1,q_1)} \varepsilon_{(i_1,c_1)}| | \mathcal{F}_n^Z \right] E \left[|\varepsilon_{(i_2,r_2)} \varepsilon_{(i_2,g_2)} \varepsilon_{(i_2,q_2)} \varepsilon_{(i_2,c_2)}| | \mathcal{F}_n^Z \right] \}
\end{aligned}$$

$$\begin{aligned}
&\leq \left\{ 4\bar{T}^6 \|J\|_\infty^4 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^6 \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^2 | \mathcal{F}_n^W] \right) \right. \\
&\quad \times \sqrt{\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z]} \sqrt{\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^8 | \mathcal{F}_n^Z]} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \\
&\quad \times \frac{1}{K_{2,n}^2} \sum_{i=1}^n \sum_{\substack{s_1, t_1=1 \\ s_1 \neq t_1}}^{T_i} \sum_{\substack{s_2, t_2=1 \\ s_2 \neq t_2}}^{T_i} A_{(i,t_1),(i,s_1)}^2 A_{(i,t_2),(i,s_2)}^2 \Bigg\} \\
&\quad + \left\{ 4\bar{T}^8 \|J\|_\infty^4 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^6 \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^W] \right)^2 \right. \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{i_1=1}^n \sum_{\substack{s_1, t_1=1 \\ s_1 \neq t_1}}^{T_{i_1}} A_{(i_1,t_1),(i_1,s_1)}^2 \frac{1}{K_{2,n}} \sum_{i_2=1}^n \sum_{\substack{s_2, t_2=1 \\ i_2 \neq i_1 \\ s_2 \neq t_2}}^{T_{i_2}} A_{(i_2,t_2),(i_2,s_2)}^2 \Bigg\} \\
&= O_{a.s.} \left(\frac{K_n}{n} \frac{K_{2,n}^2}{n^3} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \frac{K_{2,n}^2}{n^2} \right) = O_{a.s.} \left(\frac{K_n^2}{n^2} \frac{K_{2,n}^2}{n^2} \right)
\end{aligned}$$

Now, consider $\mathcal{A}_{4,1,3}$. In this case, we can apply the triangle inequality, part (a) of Lemma S2-1, and Assumptions 2(i), 5, and 6 to get

$$\begin{aligned}
&E[|\mathcal{A}_{4,1,3}| | \mathcal{F}_n^Z] \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{r=1}^{T_i} |P_{(i,h),(i,r)}^{Z^\perp}| \sum_{g=1}^{T_i} |M_{(i,h),(i,g)}^Q| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{q=1}^{T_j} |M_{(j,v),(j,q)}^Q| \sum_{c=1}^{T_j} |M_{(j,v),(j,c)}^Q| E[|\varepsilon_{(i,r)} \varepsilon_{(i,g)} \varepsilon_{(j,q)} \varepsilon_{(j,c)}| | \mathcal{F}_n^Z] \\
&\leq 2\bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^3 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E[\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right) \frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.} \left(\frac{K_n}{n} \right)
\end{aligned}$$

Turning our attention to $\mathcal{A}_{4,1,4}$, note that, by applying the triangle inequality, part (a) of Lemma

S2-1, and Assumptions 2(i), 5, and 6; we obtain

$$\begin{aligned}
& E \left[|\mathcal{A}_{4,1,4}| \mid \mathcal{F}_n^Z \right] \\
& \leq \frac{2}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{g=1}^{T_i} \left| M_{(i,h),(i,g)}^Q \right| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \\
& \quad \times \sum_{r=1}^{T_j} \left| P_{(i,h),(j,r)}^{Z^\perp} \right| \sum_{q=1}^{T_j} \left| M_{(j,v),(j,q)}^Q \right| \sum_{c=1}^{T_j} \left| M_{(j,v),(j,c)}^Q \right| E \left[|\varepsilon_{(i,g)} \varepsilon_{(j,r)} \varepsilon_{(j,q)} \varepsilon_{(j,c)}| \mid \mathcal{F}_n^Z \right] \\
& \leq 2\bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} \left| M_{(i,t),(j,s)}^Q \right| \right)^3 \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} E \left[\varepsilon_{(i,t)}^4 \mid \mathcal{F}_n^Z \right] \right) \frac{1}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \\
& = O_{a.s.} \left(\frac{K_n}{n} \right).
\end{aligned}$$

Finally, consider

$$\begin{aligned}
& E \left[\mathcal{A}_{4,1,5}^2 \mid \mathcal{F}_n^Z \right] \\
& = E \left[\left(\frac{2}{K_{2,n}} \sum_{\substack{i,j=1 \\ i \neq j}}^n \sum_{t=1}^{T_i} \sum_{s=1}^{T_j} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \sum_{g=1}^{T_i} M_{(i,h),(i,g)}^Q \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \sum_{\substack{(l,r)=1 \\ l \neq i,j}}^{m_n} P_{(i,h),(l,r)}^{Z^\perp} \right. \right. \\
& \quad \times \left. \left. \sum_{q=1}^{T_j} M_{(j,v),(j,q)}^Q \sum_{c=1}^{T_j} M_{(j,v),(j,c)}^Q \varepsilon_{(l,r)} \varepsilon_{(i,g)} \varepsilon_{(j,q)} \varepsilon_{(j,c)} \right)^2 \mid \mathcal{F}_n^Z \right]
\end{aligned}$$

In this case, note that, by applying the triangle and Jensen's inequalities, parts (a) and (e) of Lemma

S2-1, and Assumptions 1, 2(i), 5, and 6; we obtain

$$\begin{aligned}
& E [\mathcal{A}_{4,1,5}^2 | \mathcal{F}_n^Z] \\
\leq & \frac{4}{K_{2,n}^2} \sum_{\substack{i_1, j_1=1 \\ i_1 \neq j_1}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_{j_1}} A_{(i_1, t_1), (j_1, s_1)}^2 \sum_{\substack{i_2, j_2=1 \\ i_2 \neq j_2}}^n \sum_{t_2=1}^{T_{i_2}} \sum_{s_2=1}^{T_{j_2}} A_{(i_2, t_2), (j_2, s_2)}^2 \sum_{h_1=1}^{T_{i_1}} |J_{(i_1, t_1), (i_1, h_1)}| \sum_{h_2=1}^{T_i} |J_{(i_2, t_2), (i_2, h_2)}| \\
& \times \sum_{v_1=1}^{T_{j_1}} |J_{(j_1, s_1), (j_1, v_1)}| \sum_{v_2=1}^{T_{j_2}} J_{(j_2, s_2), (j_2, v_2)} \sum_{\substack{(l,r)=1 \\ l \neq i_1, j_1, i_2, j_2}}^{m_n} \left| P_{(i_1, h_1), (l, r)}^{Z^\perp} P_{(i_2, h_2), (l, r)}^{Z^\perp} \right| \sum_{g_1=1}^{T_{i_1}} \left| M_{(i_1, h_1), (i_1, g_1)}^Q \right| \\
& \times \sum_{g_2=1}^{T_{i_2}} \left| M_{(i_2, h_2), (i_2, g_2)}^Q \right| \sum_{q_1=1}^{T_{j_1}} \left| M_{(j_1, v_1), (j_1, q_1)}^Q \right| \sum_{q_2=1}^{T_{j_2}} \left| M_{(j_2, v_2), (j_2, q_2)}^Q \right| \sum_{c_1=1}^{T_{j_1}} \left| M_{(j_1, v_1), (j_1, c_1)}^Q \right| \\
& \times \sum_{c_2=1}^{T_{j_2}} \left| M_{(j_2, v_2), (j_2, c_2)}^Q \right| E \left[\varepsilon_{(l, r)}^2 | \mathcal{F}_n^Z \right] E \left[|\varepsilon_{(i_1, g_1)} \varepsilon_{(j_1, q_1)} \varepsilon_{(j_1, c_1)} \varepsilon_{(i_2, g_2)} \varepsilon_{(j_2, q_2)} \varepsilon_{(j_2, c_2)}| | \mathcal{F}_n^Z \right] \\
& + \frac{4}{K_{2,n}^2} \sum_{\substack{i_1, j_1=1 \\ i_1 \neq j_1}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_{j_1}} A_{(i_1, t_1), (j_1, s_1)}^2 \sum_{\substack{i_2, j_2=1 \\ i_2 \neq j_2, i_1}}^n \sum_{t_2=1}^{T_{i_2}} \sum_{s_2=1}^{T_{j_2}} A_{(i_2, t_2), (j_2, s_2)}^2 \sum_{h_1=1}^{T_{i_1}} |J_{(i_1, t_1), (i_1, h_1)}| \sum_{h_2=1}^{T_{i_2}} |J_{(i_2, t_2), (i_2, h_2)}| \\
& \times \sum_{v_1=1}^{T_{j_1}} |J_{(j_1, s_1), (j_1, v_1)}| \sum_{v_2=1}^{T_{j_2}} |J_{(j_2, s_2), (j_2, v_2)}| \sum_{g_1=1}^{T_{i_1}} \left| M_{(i_1, h_1), (i_1, g_1)}^Q \right| \sum_{g_2=1}^{T_{i_2}} \left| M_{(i_2, h_2), (i_2, g_2)}^Q \right| \sum_{q_1=1}^{T_{j_1}} \left| M_{(j_1, v_1), (j_1, q_1)}^Q \right| \\
& \times \sum_{q_2=1}^{T_{j_2}} \left| M_{(j_2, v_2), (j_2, q_2)}^Q \right| \sum_{c_1=1}^{T_{j_1}} \left| M_{(j_1, v_1), (j_1, c_1)}^Q \right| \sum_{c_2=1}^{T_{j_2}} \left| M_{(j_2, v_2), (j_2, c_2)}^Q \right| \sum_{r_1=1}^{T_{i_2}} \left| P_{(i_1, h_1), (i_2, g_2)}^{Z^\perp} \right| \\
& \times \sum_{r_2=1}^{T_{i_1}} \left| P_{(i_2, h_2), (i_1, g_1)}^{Z^\perp} \right| E \left[\varepsilon_{(i_1, g_1)}^2 | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(i_2, g_2)}^2 | \mathcal{F}_n^Z \right] E \left[|\varepsilon_{(j_1, q_1)} \varepsilon_{(j_1, c_1)} \varepsilon_{(j_2, q_2)} \varepsilon_{(j_2, c_2)}| | \mathcal{F}_n^Z \right] \\
& + \frac{4}{K_{2,n}^2} \sum_{i=1}^n \sum_{\substack{j_1=1 \\ j_1 \neq i}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_{j_1}} A_{(i, t_1), (j_1, s_1)}^2 \sum_{\substack{j_2=1 \\ j_2 \neq i, j_1}}^n \sum_{t_2=1}^{T_{i_2}} \sum_{s_2=1}^{T_{j_2}} A_{(i, t_2), (j_2, s_2)}^2 \sum_{h_1=1}^{T_i} |J_{(i, t_1), (i, h_1)}| \sum_{h_2=1}^{T_i} |J_{(i, t_2), (i, h_2)}| \\
& \times \sum_{v_1=1}^{T_{j_1}} |J_{(j_1, s_1), (j_1, v_1)}| \sum_{v_2=1}^{T_{j_2}} |J_{(j_2, s_2), (j_2, v_2)}| \sum_{g=1}^{T_i} \left| M_{(i, h_1), (i, g)}^Q \right| \left| M_{(i, h_2), (i, g)}^Q \right| \sum_{q_1=1}^{T_{j_1}} \left| M_{(j_1, v_1), (j_1, q_1)}^Q \right| \\
& \times \sum_{q_2=1}^{T_{j_2}} \left| M_{(j_2, v_2), (j_2, q_2)}^Q \right| \sum_{c_1=1}^{T_{j_1}} \left| M_{(j_1, v_1), (j_1, c_1)}^Q \right| \sum_{c_2=1}^{T_{j_2}} \left| M_{(j_2, v_2), (j_2, c_2)}^Q \right| \sum_{r_1=1}^{T_i} \left| P_{(i, h_1), (j_2, r_1)}^{Z^\perp} \right| \\
& \times \sum_{r_2=1}^{T_i} \left| P_{(i, h_2), (j_1, r_2)}^{Z^\perp} \right| E \left[|\varepsilon_{(j_1, r_2)} \varepsilon_{(j_1, q_1)} \varepsilon_{(j_1, c_1)}| | \mathcal{F}_n^Z \right] E \left[|\varepsilon_{(j_2, r_1)} \varepsilon_{(j_2, q_2)} \varepsilon_{(j_2, c_2)}| | \mathcal{F}_n^Z \right] E \left[\varepsilon_{(i, g)}^2 | \mathcal{F}_n^Z \right]
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{4\bar{T}^6 \|J\|_\infty^4}{K_{2,n}^2} \sum_{\substack{i_1, j_1=1 \\ i_1 \neq j_1}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_{j_1}} A_{(i_1, t_1), (j_1, s_1)}^2 \sum_{\substack{i_2, j_2=1 \\ i_2 \neq j_2}}^n \sum_{t_2=1}^{T_{i_2}} \sum_{s_2=1}^{T_{j_2}} A_{(i_2, t_2), (j_2, s_2)}^2 \\
&\quad \times \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^6 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \sqrt{\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z]} \\
&\quad \times \sqrt{\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^8 | \mathcal{F}_n^Z]} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t), (i,t)}^{Z^\perp} \right) \\
&\quad + \frac{4\bar{T}^8 \|J\|_\infty^4}{K_{2,n}^2} \sum_{\substack{i_1, j_1=1 \\ i_1 \neq j_1}}^n \sum_{t_1=1}^{T_{i_1}} \sum_{s_1=1}^{T_{j_1}} A_{(i_1, t_1), (j_1, s_1)}^2 \sum_{\substack{i_2, j_2=1 \\ i_2 \neq j_2}}^n \sum_{t_2=1}^{T_{i_2}} \sum_{s_2=1}^{T_{j_2}} A_{(i_2, t_2), (j_2, s_2)}^2 \\
&\quad \times \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^6 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right)^2 \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t), (i,t)}^{Z^\perp} \right)^2 \\
&\quad + \frac{4\bar{T}^7 \|J\|_\infty^4}{K_{2,n}^2} \sum_{i=1}^n \sum_{\substack{j_1=1 \\ j_1 \neq i}}^{T_i} \sum_{t_1=1}^{T_{j_1}} \sum_{s_1=1}^{T_{j_1}} A_{(i, t_1), (j_1, s_1)}^2 \sum_{\substack{j_2=1 \\ j_2 \neq i}}^n \sum_{t_2=1}^{T_{j_2}} \sum_{s_2=1}^{T_{j_2}} A_{(i, t_2), (j_2, s_2)}^2 \left(\max_{1 \leq (i,t), (j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^6 \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t), (i,t)}^{Z^\perp} \right)^2 \\
&= O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) + O_{a.s.} \left(\frac{K_n^2}{n^3} \right) = O_{a.s.} \left(\frac{K_n}{n} \right).
\end{aligned}$$

It follows from these calculations and from Liapunov's inequality that

$$\begin{aligned}
&E [|\mathcal{A}_{4,1}| |\mathcal{F}_n^Z] \\
&\leq E [|\mathcal{A}_{4,1,1}| |\mathcal{F}_n^Z] + E [|\mathcal{A}_{4,1,2}| |\mathcal{F}_n^Z] + E [|\mathcal{A}_{4,1,3}| |\mathcal{F}_n^Z] + E [|\mathcal{A}_{4,1,4}| |\mathcal{F}_n^Z] \\
&\quad + E [|\mathcal{A}_{4,1,5}| |\mathcal{F}_n^Z] \\
&\leq E [|\mathcal{A}_{4,1,1}| |\mathcal{F}_n^Z] + \sqrt{E [\mathcal{A}_{4,1,2}^2 |\mathcal{F}_n^Z]} + E [|\mathcal{A}_{4,1,3}| |\mathcal{F}_n^Z] + E [|\mathcal{A}_{4,1,4}| |\mathcal{F}_n^Z] \\
&\quad + \sqrt{E [\mathcal{A}_{4,1,5}^2 |\mathcal{F}_n^Z]} \\
&= O_{a.s.} \left(\frac{K_n K_{2,n}}{n^2} \right) + O_{a.s.} \left(\frac{K_n K_{2,n}}{n^2} \right) + O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\sqrt{\frac{K_n}{n}} \right) \\
&= O_{a.s.} \left(\sqrt{\frac{K_n}{n}} \right)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left[\sqrt{\frac{n}{K_n}} |\mathcal{A}_{4,1}| \right] = E_W \left(\sqrt{\frac{n}{K_n}} E [|\mathcal{A}_{4,1}| | \mathcal{F}_n^Z] \right) \leq \bar{C}.$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\Pr \left(\left| \sqrt{\frac{n}{K_n}} \mathcal{A}_{4,1} \right| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \sqrt{\frac{n}{K_n}} \frac{E [|\mathcal{A}_{4,1}|]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{4,1} = O_p \left(\sqrt{\frac{K_n}{n}} \right) \tag{83}$$

Turning our attention to part (b), note that, here, we can apply the triangle and CS inequalities

and the inequality $|XY| \leq (1/2)X^2 + (1/2)Y^2$ to obtain

$$\begin{aligned}
|\mathcal{A}_{4,2}| &\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left| \left(e'_{(i,h)} M^{(Z,Q)} X \left[\widehat{\delta}_n - \delta_0 \right] \right) \left(e'_{(i,h)} M^{Q_\varepsilon} \right) \right| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad \times \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sqrt{e'_{(i,h)} M^{(Z,Q)} X \left[\widehat{\delta}_n - \delta_0 \right] \left[\widehat{\delta}_n - \delta_0 \right]' X' M^{(Z,Q)} e_{(i,h)}} \\
&\quad \times \sqrt{e'_{(i,h)} M^{Q_\varepsilon} \varepsilon' M^Q e_{(i,h)}} \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \\
&\leq \left\| \widehat{\delta}_n - \delta_0 \right\|_2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \\
&\quad + \left\| \widehat{\delta}_n - \delta_0 \right\|_2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^Q \varepsilon' M^Q e_{(i,h)} \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2 \\
&= \left\| \widehat{\delta}_n - \delta_0 \right\|_2 (\mathcal{A}_{4,2,1} + \mathcal{A}_{4,2,2}),
\end{aligned} \tag{84}$$

where

$$\begin{aligned}
\mathcal{A}_{4,2,1} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2, \\
\mathcal{A}_{4,2,2} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^Q \varepsilon' M^Q e_{(i,h)} \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^{Q_\varepsilon} \right)^2
\end{aligned}$$

Next, observe that

$$\begin{aligned}
0 &\leq \mathcal{A}_{4,2,1} \\
&= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(i,h)} M^{(Z,Q)} UU' M^{(Z,Q)} e_{(i,h)} \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 e'_{(i,h)} M^Q UU' M^Q e_{(i,h)} \\
&\quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 e'_{(i,h)} P^{Z^\perp} UU' P^{Z^\perp} e_{(i,h)} \\
&= \mathcal{A}_{4,2,1,1} + \mathcal{A}_{4,2,1,2}, \quad (\text{say}).
\end{aligned}$$

Considering first $\mathcal{A}_{4,2,1,1}$. Applying part (a) of Lemma S2-1 and Assumptions 2(i), 5, and 6; we get

$$\begin{aligned}
0 &\leq E [\mathcal{A}_{4,2,1,1} | \mathcal{F}_n^Z] \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&\quad \times \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| E \left[\left(e'_{(j,v)} M^Q \varepsilon \right)^2 e'_{(i,h)} M^Q UU' M^Q e_{(i,h)} | \mathcal{F}_n^Z \right] \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{g=1}^{T_i} \sum_{r=1}^{T_i} |M_{(i,h),(i,g)}^Q| |M_{(i,h),(i,r)}^Q| \\
&\quad \times \sum_{q=1}^{T_j} \sum_{c=1}^{T_j} |M_{(j,v),(j,q)}^Q| |M_{(j,v),(j,c)}^Q| E \left[|\varepsilon_{(j,q)} \varepsilon_{(j,c)} U'_{(i,g)} U_{(i,r)}| | \mathcal{F}_n^Z \right] \\
&\leq 2\bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^4 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^{1/2} \\
&\quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z] \right)^{1/2} \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.}(1).
\end{aligned}$$

In addition, for $E [\mathcal{A}_{4,2,1,2} | \mathcal{F}_n^Z]$, we have, by straightforward calculations using Assumption 1 and

the triangle inequality,

$$\begin{aligned}
0 &\leq E[\mathcal{A}_{4,2,1,2}|\mathcal{F}_n^Z] \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| E\left[\left(e'_{(j,v)} M^Q \varepsilon\right)^2 e'_{(i,h)} P^{Z^\perp} UU' P^{Z^\perp} e_{(i,h)} |\mathcal{F}_n^Z\right] \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{g=1}^{T_j} \sum_{r=1}^{T_j} M_{(j,v),(j,g)}^Q M_{(j,v),(j,r)}^Q \\
&\quad \times \sum_{(k,q)=1}^{m_n} P_{(i,h),(k,q)}^{Z^\perp} \sum_{(l,c)=1}^{m_n} P_{(i,h),(l,c)}^{Z^\perp} E\left[\varepsilon_{(j,g)} \varepsilon_{(j,r)} U'_{(k,q)} U_{(l,c)} |\mathcal{F}_n^Z\right] \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \right. \\
&\quad \times \left. \sum_{g=1}^{T_j} \left(M_{(j,v),(j,g)}^Q\right)^2 \left(P_{(i,h),(j,g)}^{Z^\perp}\right)^2 E\left[\varepsilon_{(j,g)}^2 U'_{(j,g)} U_{(j,g)} |\mathcal{F}_n^Z\right]\right\} \\
&\quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} \sum_{v=1}^{T_j} |J_{(i,t),(i,h)}| |J_{(j,s),(j,v)}| \sum_{g=1}^{T_j} \left(M_{(j,v),(j,g)}^Q\right)^2 \right. \\
&\quad \times \left. \sum_{(k,q)=1}^{m_n} \left(P_{(i,h),(k,q)}^{Z^\perp}\right)^2 E\left[\varepsilon_{(j,g)}^2 |\mathcal{F}_n^Z\right] E\left[U'_{(k,q)} U_{(k,q)} |\mathcal{F}_n^Z\right]\right\} \\
&\quad + \frac{4}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{g=1}^{T_i} \sum_{r=1}^{T_i} \left|M_{(j,v),(j,g)}^Q\right| \left|M_{(j,v),(j,r)}^Q\right| \right. \\
&\quad \times \left. \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left|P_{(i,h),(j,g)}^{Z^\perp}\right| \left|P_{(i,h),(j,r)}^{Z^\perp}\right| E\left[\left|\varepsilon_{(j,g)} \varepsilon_{(j,r)} U'_{(j,g)} U_{(j,r)}\right| |\mathcal{F}_n^Z\right]\right\} \tag{85}
\end{aligned}$$

Applying the CS inequality to (85) and making use of part (a) of Lemma S2-1 as well as Assumptions

2(i), 5, and 6; we then obtain

$$\begin{aligned}
& E [\mathcal{A}_{4,2,1,2} | \mathcal{F}_n^Z] \\
& \leq 2\bar{T} \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^{1/2} \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
& \quad + 2\bar{T} \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^2 | \mathcal{F}_n^Z] \right) \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\|U_{(i,t)}\|_2^2 | \mathcal{F}_n^Z] \right) \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right) \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
& \quad + 4\bar{T}^2 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^2 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right)^{1/2} \\
& \quad \times \left(\max_{1 \leq (i,t) \leq m_n} E [\|U_{(i,t)}\|_2^4 | \mathcal{F}_n^Z] \right)^{1/2} \left(\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z^\perp} \right)^2 \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
& = O_{a.s.} \left(\frac{K_n^2}{n^2} \right) + O_{a.s.} \left(\frac{K_n}{n} \right) + O_{a.s.} \left(\frac{K_n^2}{n^2} \right) = O_{a.s.} \left(\frac{K_n}{n} \right) = o_{a.s.}(1).
\end{aligned}$$

It follows from these calculations that

$$\begin{aligned}
0 & \leq E [\mathcal{A}_{4,2,1} | \mathcal{F}_n^Z] \\
& = E [\mathcal{A}_{4,2,1,1} | \mathcal{F}_n^W] + E [\mathcal{A}_{4,2,1,2} | \mathcal{F}_n^Z] \\
& = O_{a.s.}(1) + O_{a.s.} \left(\frac{K_n}{n} \right) \\
& = O_{a.s.}(1)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E [|\mathcal{A}_{4,2,1}|] = E [\mathcal{A}_{4,2,1}] = E_Z (E [\mathcal{A}_{4,2,1} | \mathcal{F}_n^Z]) \leq \bar{C}.$$

Application of the Markov's inequality then implies that, for any $\epsilon > 0$,

$$\Pr \left(|\mathcal{A}_{4,2,1}| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{E [\mathcal{A}_{4,2,2}]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{4,2,1} = O_p(1). \tag{86}$$

In addition, applying the triangle inequality, part (a) of Lemma S2-1, as well as Assumptions

2(i), 5, and 6; we obtain

$$\begin{aligned}
0 &\leq E [\mathcal{A}_{4,2,2} | \mathcal{F}_n^Z] \\
&\leq \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \sum_{r=1}^{T_i} |M_{(i,h),(i,r)}^Q| \sum_{c=1}^{T_i} |M_{(i,h),(i,c)}^Q| \\
&\quad \times \sum_{q=1}^{T_j} |M_{(j,v),(j,q)}^Q| \sum_{g=1}^{T_j} |M_{(j,v),(j,g)}^Q| E [|\varepsilon_{(i,r)} \varepsilon_{(i,c)} \varepsilon_{(j,q)} \varepsilon_{(j,g)}| | \mathcal{F}_n^Z] \\
&\leq \bar{T}^4 \|J\|_\infty^2 \left(\max_{1 \leq (i,t),(j,s) \leq m_n} |M_{(i,t),(j,s)}^Q| \right)^4 \left(\max_{1 \leq (i,t) \leq m_n} E [\varepsilon_{(i,t)}^4 | \mathcal{F}_n^Z] \right) \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \\
&= O_{a.s.}(1)
\end{aligned}$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E [|\mathcal{A}_{4,2,2}|] = E [\mathcal{A}_{4,2,2}] = E_Z (E [\mathcal{A}_{4,2,2} | \mathcal{F}_n^Z]) \leq \bar{C}.$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\Pr \left(|\mathcal{A}_{4,2,2}| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{E [|\mathcal{A}_{4,2,2}|]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{4,2,2} = O_p(1). \tag{87}$$

It follows from expressions (84)-(87) that

$$|\mathcal{A}_{4,2}| \leq \left\| \widehat{\delta}_n - \delta_0 \right\|_2 [\mathcal{A}_{4,2,1} + \mathcal{A}_{4,2,2}] = o_p(1) [O_p(1) + O_p(1)] = o_p(1). \tag{88}$$

Finally, for part (c), note that we can write

$$\begin{aligned}
\mathcal{A}_4 &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad \times \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \\
&= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \\
&\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad - \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(i,h)} M^Q \varepsilon \right) \\
&\quad \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&= \mathcal{A}_{4,1} + \mathcal{A}_{4,2},
\end{aligned}$$

where $\mathcal{A}_{4,1}$ and $\mathcal{A}_{4,2}$ are as defined in parts (a) and (b) above. It follows from expressions (83) and (88) that

$$\mathcal{A}_4 = \mathcal{A}_{4,1} + \mathcal{A}_{4,2} = O_p \left(\sqrt{\frac{K_n}{n}} \right) + o_p(1) = o_p(1). \quad \square \quad (89)$$

Lemma OA-18: Let Assumptions 1-6 be satisfied, and let $\{\widehat{\delta}_n\}$ be a sequence of estimators such that

$$\left\| \widehat{\delta}_n - \delta_0 \right\|_2 \xrightarrow{p} 0 \text{ as } n \rightarrow \infty,$$

as long as $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Then, the following statements are true.

(a)

$$\mathcal{A}_{5,1} = O_p \left(\frac{K_n}{n} \right) = o_p(1),$$

where

$$\mathcal{A}_{5,1} = \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \left(e'_{(j,v)} M^Q \varepsilon \right)^2.$$

(b)

$$\mathcal{A}_{5,2} = O_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{5,2} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \right\} \end{aligned}$$

(c)

$$\mathcal{A}_5 = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_5 &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\hat{\delta}_n - \delta_0] \right)^2 \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \right\} \end{aligned}$$

Proof of Lemma OA-18:

To show part (a), note that $\mathcal{A}_{5,1}$ is of the same form as $\mathcal{A}_{2,1,1}$, so from expression (68), we have that

$$E[\mathcal{A}_{5,1} | \mathcal{F}_n^Z] = O_{a.s.} \left(\frac{K_n}{n} \right).$$

Hence, there exists a constant $\bar{C} < \infty$ such that for all n sufficiently large

$$E \left[\frac{n}{K_n} |\mathcal{A}_{5,1}| \right] = E_Z \left(\frac{n}{K_n} E[|\mathcal{A}_{5,1}| | \mathcal{F}_n^Z] \right) \leq \bar{C}.$$

It follows from the Markov's inequality that for any $\epsilon > 0$,

$$\Pr \left(\left| \frac{n}{K_n} \mathcal{A}_{5,1} \right| \geq \frac{\bar{C}}{\epsilon} \right) \leq \epsilon \frac{n}{K_n} \frac{E[|\mathcal{A}_{5,1}|]}{\bar{C}} \leq \epsilon$$

for all n sufficiently large, which shows that

$$\mathcal{A}_{5,1} = O_p \left(\frac{K_n}{n} \right). \tag{90}$$

For part (b), note that $\mathcal{A}_{5,2}$ is of the same form as $\mathcal{A}_{2,2,2}$, so that by expression (73), we have

that

$$\mathcal{A}_{5,2} = O_p(1). \quad (91)$$

Finally, consider part (c). Applying the triangle inequality and the inequality $\left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r$, we obtain

$$\begin{aligned} |\mathcal{A}_5| &\leq \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right]^2 \\ &\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\ &\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\ &\quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\ &\quad \times \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] [\widehat{\delta}_n - \delta_0]' X' M^{(Z,Q)} e_{(i,h)} \\ &\leq \mathcal{A}_{5,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{5,2}, \end{aligned} \quad (92)$$

where $\mathcal{A}_{5,1}$ and $\mathcal{A}_{5,2}$ are as defined in parts (a) and (b) above. In light of the upper bound given by (92), it then follows from expressions (90) and (91) that

$$|\mathcal{A}_5| \leq \mathcal{A}_{5,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{5,2} = O_p \left(\frac{K_n}{n} \right) + o_p(1) O_p(1) = o_p(1). \quad \square \quad (93)$$

Lemma OA-19: Let Assumptions 1-6 be satisfied, and let $\{\widehat{\delta}_n\}$ be a sequence of estimators such that

$$\left\| \widehat{\delta}_n - \delta_0 \right\|_2 \xrightarrow{p} 0 \text{ as } n \rightarrow \infty,$$

as long as $\sqrt{K_{2,n}} / (\mu_n^{\min})^2 \rightarrow 0$. Then, the following statements are true.

(a)

$$\mathcal{A}_{6,1} = O_p \left(\frac{K_n}{n} \right) = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{6,1} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left[e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right]^2 \right\} \end{aligned}$$

(b)

$$\mathcal{A}_{6,2} = O_p(1),$$

where

$$\begin{aligned} \mathcal{A}_{6,2} &= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \right\} \end{aligned}$$

(c)

$$\mathcal{A}_6 = o_p(1),$$

where

$$\begin{aligned} \mathcal{A}_6 &= -\frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \left\{ \sum_{h=1}^{T_i} J_{(i,t),(i,h)} \left(e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right)^2 \right. \\ &\quad \left. \times \sum_{v=1}^{T_j} J_{(j,s),(j,v)} \left(e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right\} \end{aligned}$$

Proof of Lemma OA-19:

To show part (a), note that $\mathcal{A}_{6,1}$ has the same form as the term which we used to get an initial bound on \mathcal{A}_3 . See, in particular, the first inequality in expression (81). Hence, it follows from

expression (82) that

$$\begin{aligned}
& \mathcal{A}_{6,1} \\
&= \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left[e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right]^2 \\
&\leq \mathcal{A}_{3,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{3,2} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{3,3} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^4 \mathcal{A}_{3,4} \\
&= o_p(1). \tag{94}
\end{aligned}$$

Next, for part (b), note that we can apply the inequality $\left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r$ to

obtain

$$\begin{aligned}
0 &\leq \mathcal{A}_{6,2} \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad + \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(i,h)} M^{(Z,Q)} X \left(\widehat{\delta}_n - \delta_0 \right) \left(\widehat{\delta}_n - \delta_0 \right)' X' M^{(Z,Q)} e_{(i,h)} \\
&\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(i,h)} M^{(Z,Q)} X X' M^{(Z,Q)} e_{(i,h)} \\
&= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\
&\quad \times e'_{(i,h)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(i,h)} \\
&= \mathcal{A}_{6,2,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{6,2,2},
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{A}_{6,2,1} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left(e'_{(i,h)} P^{Z^\perp} \varepsilon \right)^2 \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2, \\
\mathcal{A}_{6,2,2} &= \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t)\neq(j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \\
&\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 e'_{(i,h)} M^{(Z,Q)} U U' M^{(Z,Q)} e_{(i,h)}.
\end{aligned}$$

Note that $\mathcal{A}_{6,2,1}$ is of the same form as $\mathcal{A}_{2,2,1}$ and $\mathcal{A}_{6,2,2}$ is of the same form as $\mathcal{A}_{2,2,2}$, so that from expressions (70) and (73), we get

$$\mathcal{A}_{6,2,1} = O_p\left(\frac{K_n}{n}\right) = o_p(1), \quad \mathcal{A}_{6,2,2} = O_p(1),$$

from which it follows that

$$0 \leq \mathcal{A}_{6,2} \leq \mathcal{A}_{6,2,1} + \left\| \widehat{\delta}_n - \delta_0 \right\|_2^2 \mathcal{A}_{6,2,2} = O_p\left(\frac{K_n}{n}\right) + o_p(1) O_p(1) = o_p(1). \quad (95)$$

Finally, to show part (c), note that, by applying the triangle inequality and the inequality $|XY| \leq (1/2)X^2 + (1/2)Y^2$ in this case, we obtain

$$\begin{aligned} |\mathcal{A}_6| &\leq \frac{2}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \\ &\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left| \left(e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right) \left(e'_{(j,v)} M^Q \varepsilon \right) \right| \\ &\leq \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \\ &\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left[e'_{(j,v)} P^{Z^\perp} \varepsilon + e'_{(j,v)} M^{(Z,Q)} X [\widehat{\delta}_n - \delta_0] \right]^2 \\ &\quad + \frac{1}{K_{2,n}} \sum_{\substack{(i,t),(j,s)=1 \\ (i,t) \neq (j,s)}}^{m_n} A_{(i,t),(j,s)}^2 \sum_{h=1}^{T_i} |J_{(i,t),(i,h)}| \left[e'_{(i,h)} P^{Z^\perp} \varepsilon + e'_{(i,h)} M^{(Z,Q)} X (\widehat{\delta}_n - \delta_0) \right]^2 \\ &\quad \times \sum_{v=1}^{T_j} |J_{(j,s),(j,v)}| \left(e'_{(j,v)} M^Q \varepsilon \right)^2 \\ &= \mathcal{A}_{6,1} + \mathcal{A}_{6,2}, \end{aligned} \quad (96)$$

where $\mathcal{A}_{6,1}$ and $\mathcal{A}_{6,2}$ are as defined in parts (a) and (b) above. In light of the upper bound for $|\mathcal{A}_6|$ given in (96), it then follows from expressions (94) and (95) that

$$|\mathcal{A}_6| \leq \mathcal{A}_{6,1} + \mathcal{A}_{6,2} = o_p(1) + o_p(1) = o_p(1). \quad \square \quad (97)$$

The next lemma provides more primitive sufficient conditions for part (iv) of Assumption 5 in the main paper. To facilitate the statement and proof of this lemma, we introduce a number of additional notations. Let $Z_{1,\cdot,k}$ and $Z_{1,(i,t),k}$ denote, respectively, the k^{th} column and the $((i,t), k)^{th}$ element of Z_1 for $(i,t) = 1, \dots, m_n$ and $k = 1, \dots, K_{1,n}$. Similarly, let $Z_{2,\cdot,k}$ and $Z_{2,(i,t),k}$ denote, respectively, the k^{th} column and the $((i,t), k)^{th}$ element of Z_2 for $(i,t) = 1, \dots, m_n$ and $k =$

$1, \dots, K_{2,n}$. Finally, let $\tilde{Z}_{2,(i,t),k} = e'_{(i,t)} P^{Z_1^\perp} Z_{2,.,k}$, where $P^{Z_1^\perp} = M^Q Z_1 (Z'_1 M^Q Z_1)^{-1} Z'_1 M^Q$.

Lemma OA-20:

(a) Assume the following conditions hold:

(a1) There exists a positive constant C such that

$$\lambda_{\min} \left(\frac{Z'_1 M^Q Z_1}{m_n} \right) \geq C > 0 \text{ a.s. for all } n \text{ sufficiently large.}$$

(a2)

$$\max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{1,n}} \sum_{k=1}^{K_{1,n}} Z_{1,(i,t),k}^2 \right) = O_{a.s.}(1)$$

Under conditions (a1) and (a2),

$$\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z_1^\perp} = O_{a.s.} \left(\frac{K_{1,n}}{n} \right),$$

where $P_{(i,t),(i,t)}^{Z_1^\perp}$ is the $((i,t), (i,t))^{th}$ of the projection matrix $P^{Z_1^\perp}$.

(b) Assume the following conditions hold:

(b1) There exists a positive constant C such that

$$\lambda_{\min} \left(\frac{Z'_2 M^{(Z_1,Q)} Z_2}{m_n} \right) \geq C > 0 \text{ a.s. for all } n \text{ sufficiently large.}$$

(b2)

$$\max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} Z_{2,(i,t),k}^2 \right) = O_{a.s.}(1).$$

(b3)

$$\max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} \tilde{Z}_{2,(i,t),k}^2 \right) = O_{a.s.}(1).$$

Under conditions (b1)-(b3),

$$\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp = O_{a.s.} \left(\frac{K_{2,n}}{n} \right),$$

where $P_{(i,t),(i,t)}^\perp$ is the $((i,t), (i,t))^{th}$ of the projection matrix $P^\perp = M^{(Z_1,Q)} Z_2 (Z'_2 M^{(Z_1,Q)} Z_2)^{-1} Z'_2 M^{(Z_1,Q)}$.

Proof of Lemma OA-20:

To show part (a), . Note that

$$\begin{aligned}
0 &\leq \max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z_1^\perp} \\
&= \max_{1 \leq (i,t) \leq m_n} e'_{(i,t)} M^Q Z_1 (Z'_1 M^Q Z_1)^{-1} Z'_1 M^Q e_{(i,t)} \\
&= \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} e'_{(i,t)} M^Q Z_1 \left(\frac{Z'_1 M^Q Z_1}{m_n} \right)^{-1} Z'_1 M^Q e_{(i,t)} \\
&\leq \frac{1}{\lambda_{\min}(Z'_1 M^Q Z_1 / m_n)} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} e'_{(i,t)} M^Q Z_1 Z'_1 M^Q e_{(i,t)} \\
&\leq \frac{1}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left\{ \sum_{k=1}^{K_{1,n}} e'_{(i,t)} M^Q Z_{1,\cdot,k} Z'_{1,\cdot,k} M^Q e_{(i,t)} \right\} \\
&= \frac{1}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \sum_{k=1}^{K_{1,n}} \left(e'_{(i,t)} M^Q Z_{1,\cdot,k} \right)^2 \\
&= \frac{1}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \sum_{k=1}^{K_{1,n}} \left(Z_{1,(i,t),k} - T_i^{-1} \sum_{s=1}^{T_i} Z_{1,(i,s),k} \right)^2 \\
&\leq \frac{2}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left[\sum_{k=1}^{K_{1,n}} Z_{1,(i,t),k}^2 + \sum_{k=1}^{K_{1,n}} \left(\frac{1}{T_i} \sum_{s=1}^{T_i} Z_{1,(i,s),k} \right)^2 \right] \\
&\quad \left(\text{by applying the inequality } \left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r \text{ for } r \geq 1 \right) \\
&\leq \frac{2}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left[\sum_{k=1}^{K_{1,n}} Z_{1,(i,t),k}^2 + \sum_{k=1}^{K_{1,n}} \frac{1}{T_i} \sum_{s=1}^{T_i} Z_{1,(i,s),k}^2 \right] \text{ (by Jensen's inequality)}
\end{aligned}$$

Hence, by multiplying and dividing the majorant side of the above expression by $K_{1,n}$ and making use of the assumption that

$$\max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{1,n}} \sum_{k=1}^{K_{1,n}} Z_{1,(i,t),k}^2 \right) = O_{a.s.}(1);$$

we obtain

$$\begin{aligned}
\max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z_1^\perp} &\leq \frac{2}{C} \frac{K_{1,n}}{m_n} \max_{1 \leq (i,t) \leq m_n} \left[\frac{1}{K_{1,n}} \sum_{k=1}^{K_{1,n}} Z_{1,(i,t),k}^2 + \frac{1}{T_i} \sum_{s=1}^{T_i} \left(\frac{1}{K_{1,n}} \sum_{k=1}^{K_{1,n}} Z_{1,(i,s),k}^2 \right) \right] \\
&\leq \frac{2}{C} \frac{K_{1,n}}{m_n} \left[\max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{1,n}} \sum_{k=1}^{K_{1,n}} Z_{1,(i,t),k}^2 \right) + \frac{1}{T_i} \sum_{s=1}^{T_i} \max_{1 \leq (i,s) \leq m_n} \left(\frac{1}{K_{1,n}} \sum_{k=1}^{K_{1,n}} Z_{1,(i,s),k}^2 \right) \right] \\
&= O_{a.s.} \left(\frac{K_{1,n}}{m_n} \right) = O_{a.s.} \left(\frac{K_{1,n}}{n} \right)
\end{aligned}$$

given that $m_n \sim n$.

To show part (b), note that

$$\begin{aligned}
0 &\leq \max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \\
&= \max_{1 \leq (i,t) \leq m_n} e'_{(i,t)} M^{(Z_1, Q)} Z_2 \left(Z'_2 M^{(Z_1, Q)} Z_2 \right)^{-1} Z'_2 M^{(Z_1, Q)} e_{(i,t)} \\
&= \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} e'_{(i,t)} M^{(Z_1, Q)} Z_2 \left(\frac{Z'_2 M^{(Z_1, Q)} Z_2}{m_n} \right)^{-1} Z'_2 M^{(Z_1, Q)} e_{(i,t)} \\
&\leq \frac{1}{\lambda_{\min}(Z'_2 M^{(Z_1, Q)} Z_2 / m_n)} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} e'_{(i,t)} M^{(Z_1, Q)} Z_2 Z'_2 M^{(Z_1, Q)} e_{(i,t)} \\
&\leq \frac{1}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} e'_{(i,t)} M^{(Z_1, Q)} Z_2 Z'_2 M^{(Z_1, Q)} e_{(i,t)}
\end{aligned}$$

Next, observe that

$$\begin{aligned}
&e'_{(i,t)} M^{(Z_1, Q)} Z_2 Z'_2 M^{(Z_1, Q)} e_{(i,t)} \\
&= e'_{(i,t)} \left[M^Q - P^{Z_1^\perp} \right] Z_2 Z'_2 \left[M^Q - P^{Z_1^\perp} \right] e_{(i,t)} \\
&= e'_{(i,t)} M^Q Z_2 Z'_2 M^Q e_{(i,t)} - 2e'_{(i,t)} P^{Z_1^\perp} Z_2 Z'_2 M^Q e_{(i,t)} + e'_{(i,t)} P^{Z_1^\perp} Z_2 Z'_2 P^{Z_1^\perp} e_{(i,t)} \\
&\leq 2e'_{(i,t)} M^Q Z_2 Z'_2 M^Q e_{(i,t)} + 2e'_{(i,t)} P^{Z_1^\perp} Z_2 Z'_2 P^{Z_1^\perp} e_{(i,t)} \\
&\quad \left(\text{by applying the inequality } \left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r \text{ for } r \geq 1 \right)
\end{aligned}$$

where $P^{Z_1^\perp} = M^Q Z_1 (Z_1' M^Q Z_1)^{-1} Z_1' M^Q$. It follows that we can write

$$\begin{aligned}
0 &\leq \max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^\perp \\
&\leq \frac{2}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left\{ e'_{(i,t)} M^Q Z_2 Z_2' M^Q e_{(i,t)} + e'_{(i,t)} P^{Z_1^\perp} Z_2 Z_2' P^{Z_1^\perp} e_{(i,t)} \right\} \\
&\leq \frac{2}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left\{ \sum_{k=1}^{K_{2,n}} e'_{(i,t)} M^Q Z_{2,\cdot,k} Z_{2,\cdot,k}' M^Q e_{(i,t)} + \sum_{k=1}^{K_{2,n}} e'_{(i,t)} P^{Z_1^\perp} Z_{2,\cdot,k} Z_{2,\cdot,k}' P^{Z_1^\perp} e_{(i,t)} \right\} \\
&= \frac{1}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left\{ \sum_{k=1}^{K_{2,n}} \left(e'_{(i,t)} M^Q Z_{2,\cdot,k} \right)^2 + \sum_{k=1}^{K_{2,n}} \left(e'_{(i,t)} P^{Z_1^\perp} Z_{2,\cdot,k} \right)^2 \right\} \\
&= \frac{1}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \sum_{k=1}^{K_{2,n}} \left(Z_{2,(i,t),k} - T_i^{-1} \sum_{s=1}^{T_i} Z_{2,(i,s),k} \right)^2 + \sum_{k=1}^{K_{2,n}} \tilde{Z}_{2,(i,t),k}^2 \\
&\leq \frac{2}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left[\sum_{k=1}^{K_{1,n}} Z_{2,(i,t),k}^2 + \sum_{k=1}^{K_{2,n}} \left(\frac{1}{T_i} \sum_{s=1}^{T_i} Z_{2,(i,s),k} \right)^2 + \sum_{k=1}^{K_{2,n}} \tilde{Z}_{2,(i,t),k}^2 \right] \\
&\quad \left(\text{by applying the inequality } \left| \sum_{i=1}^m a_i \right|^r \leq m^{r-1} \sum_{i=1}^m |a_i|^r \text{ for } r \geq 1 \right) \\
&\leq \frac{2}{C} \frac{1}{m_n} \max_{1 \leq (i,t) \leq m_n} \left[\sum_{k=1}^{K_{1,n}} Z_{1,(i,t),k}^2 + \sum_{k=1}^{K_{1,n}} \frac{1}{T_i} \sum_{s=1}^{T_i} Z_{1,(i,s),k}^2 + \sum_{k=1}^{K_{2,n}} \tilde{Z}_{2,(i,t),k}^2 \right] \quad (\text{by Jensen's inequality})
\end{aligned}$$

Hence, by multiplying and dividing the majorant side of the above expression by $K_{2,n}$ and making use of the assumptions that

$$\max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} Z_{2,(i,t),k}^2 \right) = O_{a.s.}(1) \quad \text{and} \quad \max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} \tilde{Z}_{2,(i,t),k}^2 \right) = O_{a.s.}(1);$$

we obtain

$$\begin{aligned}
& \max_{1 \leq (i,t) \leq m_n} P_{(i,t),(i,t)}^{Z_1^\perp} \\
& \leq \frac{2}{C} \frac{K_{2,n}}{m_n} \max_{1 \leq (i,t) \leq m_n} \left[\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} Z_{2,(i,t),k}^2 + \frac{1}{T_i} \sum_{s=1}^{T_i} \left(\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} Z_{2,(i,s),k}^2 \right) + \frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} \tilde{Z}_{2,(i,t),k}^2 \right] \\
& \leq \frac{2}{C} \frac{K_{2,n}}{m_n} \left[\max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} Z_{2,(i,t),k}^2 \right) + \frac{1}{T_i} \sum_{s=1}^{T_i} \max_{1 \leq (i,s) \leq m_n} \left(\frac{1}{K_{1,n}} \sum_{k=1}^{K_{2,n}} Z_{2,(i,s),k}^2 \right) \right. \\
& \quad \left. + \max_{1 \leq (i,t) \leq m_n} \left(\frac{1}{K_{2,n}} \sum_{k=1}^{K_{2,n}} \tilde{Z}_{2,(i,t),k}^2 \right) \right] \\
& = O_{a.s.} \left(\frac{K_{2,n}}{m_n} \right) = O_{a.s.} \left(\frac{K_{2,n}}{n} \right)
\end{aligned}$$

given that $m_n \sim n$. \square

Section 4: Additional Monte Carlo Results

This section reports some additional Monte Carlo results which complement those presented in the main paper. Whereas the class of DGPs which generated the Monte Carlo results in the main paper concerns weakly identified situations where the instruments are all relevant but have uniformly small coefficients; the Monte Carlo design presented here considers a more heterogeneous case where there is one weak but relevant instrument while all other instruments used in estimation are completely irrelevant. More precisely, the Monte Carlo results reported here are based on the following data generating process:

DGP*:

$$\begin{aligned}
y_{(i,t)} &= \underset{1 \times 1}{\delta} \underset{1 \times 1}{x_{(i,t)}} + \underset{1 \times 10}{\varphi'} \underset{10 \times 1}{Z_{1,(i,t)}} + \alpha_i + \varepsilon_{(i,t)}, \\
x_{(i,t)} &= \underset{1 \times 10}{\Phi'} \underset{10 \times 1}{Z_{1,(i,t)}} + \underset{1 \times 1}{\pi} \underset{1 \times 1}{z_{21,(i,t)}} + \xi_i + u_{(i,t)}.
\end{aligned}$$

Here, we specify $\varphi = \iota_{10}$ and $\Phi = \iota_{10}$, with ι_{10} being a 10×1 vector of ones; and we choose the scalar parameter π so that the concentration parameter $\mu^2 = 25, 35, 45$, and 55 . The $(i,t)^{th}$ observation of the vector of instruments used in estimation is taken to be the $K_2 \times 1$ vector $Z_{2,(i,t)} = (z_{21,(i,t)} \ z_{22,(i,t)} \ \cdots \ z_{2K_2,(i,t)})'$, where $\{Z_{2,(i,t)}\}_{(i,t)=1}^{600} \equiv i.i.d.N(0, I_{K_2})$; and we consider two choices of K_2 : $K_2 = 10, 30$. On the other hand, the $(i,t)^{th}$ observation of the vector of included exogenous regressors, or covariates, is specified to be

$Z_{1,(i,t)} = (z_{1,(i,t)} \ z_{1,(i,t)}^2 \ z_{1,(i,t)}^3 \ z_{1,(i,t)}^4 \ z_{1,(i,t)} D_{(i,t),1} \ \cdots \ z_{1,(i,t)} D_{(i,t),6})'$, where $\{z_{1,(i,t)}\}_{(i,t)=1}^{600} \equiv i.i.d.N(0, 1)$ and $D_{(i,t),k} \in \{0, 1\}$ for $k \in \{1, 2, \dots, 6\}$ is a binary variable such that $\Pr(D_{(i,t),k} = 1) = 1/2$, with $\{D_{(i,t),k}\}$ taken to be independent across both (i,t) and k . Moreover, in our experiments, we set $n = 200$ and $T_i = 3$, for each $i \in \{1, 2, \dots, 200\}$, so that $m_n = 600$. We also take

$\{u_{(i,t)}\}_{(i,t)=1}^{600} \equiv i.i.d.N(0, 1)$, $\{\alpha_i\}_{i=1}^{200} i.i.d.N(0, 1)$, and $\{\xi_i\}_{i=1}^{200} i.i.d.N(0, 1)$; and $z_{1,(i,t)}$, $D_{(i,t),k}$, $Z_{2,(i,t)}$, $u_{(i,t)}$, α_i and ξ_i are all specified to be independent of each other. We allow the structural disturbance, $\varepsilon_{(i,t)}$, to exhibit conditional heteroskedasticity in a manner similar to the design given in Hausman et al (2012) and in our main paper. In particular, we let

$$\varepsilon_{(i,t)} = \rho u_{(i,t)} + \sqrt{\frac{1 - \rho^2}{\phi^2 + (0.86)^2}} (\phi v_{1,(i,t)} + 0.86 v_{2,(i,t)}) , \quad (98)$$

where $v_{1,(i,t)}|Z_{1,(i,t)}, z_{21,(i,t)} \sim N\left(0, \kappa \left[1 + (l'_{10} Z_{1,(i,t)} + z_{21,(i,t)})^2\right]\right)$ and $v_{2,(i,t)} \sim N(0, 1)$. Both of these distributions are specified to be independent across the index (i, t) , and κ is a normalization constant chosen so that the unconditional variance, $Var(v_{1,(i,t)})$, is equal to 1. For all experiments reported below, we set $\rho = 0.3$ and we choose the parameter ϕ , so that the R-squared for the regression of ε^2 on the instruments and the included exogenous variables take the values 0, 0.1, and 0.2.

The results of our Monte Carlo study based on the above data generating process are reported in Tables A1- A6 below.

Table A1: Median Bias, $K_2 = 10$

μ^2	$\mathcal{R}_{\varepsilon^2 z_1^2}^2$	2SLS	IJIVE1	IJIVE2	UJIVE	FEJIV	FELIM	FEFUL
25	0	0.1085	0.0429	0.0427	0.3845	-0.0116	-0.0002	0.0127
	0.1	0.1123	0.0466	0.0461	0.3679	-0.0055	0.0048	0.0168
	0.2	0.1100	0.0469	0.0467	0.3737	-0.0036	0.0042	0.0161
35	0	0.0875	0.0313	0.0312	0.2474	-0.0122	0.0007	0.0090
	0.1	0.0876	0.0303	0.0304	0.2611	-0.0135	0.0004	0.0094
	0.2	0.0875	0.0333	0.0331	0.2565	-0.0081	0.0021	0.0107
45	0	0.0748	0.0286	0.0289	0.2007	-0.0065	0.0023	0.0091
	0.1	0.0709	0.0247	0.0247	0.1851	-0.0121	-0.0022	0.0047
	0.2	0.0731	0.0258	0.0257	0.1916	-0.0099	0.0037	0.0108
55	0	0.0624	0.0213	0.0214	0.1430	-0.0079	0.0003	0.0056
	0.1	0.0591	0.0188	0.0191	0.1453	-0.0103	-0.0006	0.0049
	0.2	0.0603	0.0180	0.0180	0.1408	-0.0105	-0.0034	0.0022

Results based on 10,000 simulations

Table A2: Nine Decile Range 0.05 to 0.95, $K_2 = 10$

μ^2	$\mathcal{R}_{\varepsilon^2 z_1^2}^2$	2SLS	IJIVE1	IJIVE2	UJIVE	FEJIV	FELIM	FEFUL
25	0	0.6583	1.0481	1.0475	6.4416	1.6217	1.2922	1.1595
	0.1	0.6561	1.0440	1.0445	6.0634	1.6434	1.3226	1.1749
	0.2	0.6438	1.0219	1.0188	6.0248	1.6240	1.2587	1.1100
35	0	0.5803	0.8138	0.8140	5.6388	1.0813	0.9453	0.8868
	0.1	0.5827	0.8073	0.8076	6.0580	1.0644	0.9292	0.8757
	0.2	0.5858	0.8182	0.8175	5.8156	1.1009	0.9418	0.8884
45	0	0.5372	0.6936	0.6940	5.3935	0.8511	0.7762	0.7432
	0.1	0.5303	0.6845	0.6839	5.1687	0.8344	0.7528	0.7248
	0.2	0.5216	0.6806	0.6810	5.2909	0.8326	0.7476	0.7161
55	0	0.5030	0.6168	0.6169	4.5398	0.7172	0.6683	0.6505
	0.1	0.4896	0.6073	0.6079	4.8072	0.7050	0.6535	0.6376
	0.2	0.4812	0.5919	0.5931	4.8070	0.6957	0.6439	0.6269

Results based on 10,000 simulations

 Table A3: 0.05 Rejection Frequencies, $K_2 = 10$

μ^2	$\mathcal{R}_{\varepsilon^2 z_1^2}^2$	2SLS	IJIVE1	IJIVE2	UJIVE	FEJIV	FELIM	FEFUL
25	0	0.1701	0.0926	0.0863	0.5229	0.0265	0.0506	0.0511
	0.1	0.1822	0.0981	0.0925	0.5198	0.0296	0.0553	0.0570
	0.2	0.1855	0.0933	0.0879	0.5279	0.0270	0.0515	0.0529
35	0	0.1606	0.1003	0.0936	0.5307	0.0311	0.0455	0.0476
	0.1	0.1663	0.1013	0.0958	0.5387	0.0330	0.0501	0.0530
	0.2	0.1705	0.1073	0.1006	0.5390	0.0355	0.0515	0.0537
45	0	0.1573	0.1099	0.1026	0.5608	0.0369	0.0506	0.0538
	0.1	0.1611	0.1108	0.1040	0.5573	0.0391	0.0499	0.0533
	0.2	0.1645	0.1082	0.1018	0.5611	0.0356	0.0484	0.0513
55	0	0.1488	0.1106	0.1040	0.5800	0.0395	0.0496	0.0522
	0.1	0.1595	0.1221	0.1146	0.5830	0.0404	0.0507	0.0539
	0.2	0.1533	0.1118	0.1047	0.5786	0.0386	0.0506	0.0529

Results based on 10,000 simulations

Table A4: Median Bias, $K_2 = 30$

μ^2	$\mathcal{R}_{\varepsilon^2 z_1^2}^2$	2SLS	IJIVE1	IJIVE2	UJIVE	FEJIV	FELIM	FEFUL
25	0	0.1923	0.1141	0.1141	0.5684	0.0282	0.0150	0.0248
	0.1	0.1903	0.1126	0.1121	0.5663	0.0225	0.0075	0.0188
	0.2	0.1911	0.1102	0.1101	0.5747	0.0171	0.0060	0.0175
35	0	0.1688	0.0901	0.0905	0.4131	0.0012	0.0019	0.0087
	0.1	0.1703	0.0930	0.0924	0.3973	0.0004	0.0004	0.0093
	0.2	0.1718	0.0964	0.0968	0.4085	0.0031	0.0070	0.0150
45	0	0.1498	0.0775	0.0780	0.2916	-0.0084	-0.0006	0.0060
	0.1	0.1511	0.0785	0.0781	0.2920	-0.0070	-0.0004	0.0069
	0.2	0.1512	0.0764	0.0768	0.3007	-0.0071	0.0003	0.0065
55	0	0.1368	0.0668	0.0669	0.2253	-0.0054	-0.0017	0.0039
	0.1	0.1319	0.0604	0.0606	0.2157	-0.0162	-0.0063	-0.0006
	0.2	0.1382	0.0692	0.0692	0.2302	-0.0046	0.0033	0.0086

Results based on 10,000 simulations

 Table A5: Nine Decile Range 0.05 to 0.95, $K_2 = 30$

μ^2	$\mathcal{R}_{\varepsilon^2 z_1^2}^2$	2SLS	IJIVE1	IJIVE2	UJIVE	FEJIV	FELIM	FEFUL
25	0	0.4818	0.9798	0.9786	6.2121	2.9598	2.2421	1.8123
	0.1	0.4809	0.9776	0.9795	5.8280	3.0383	2.1486	1.7496
	0.2	0.4694	0.9585	0.9611	6.2576	3.1007	2.1524	1.7858
35	0	0.4481	0.8032	0.8051	5.9240	1.7994	1.4269	1.2970
	0.1	0.4386	0.7834	0.7823	6.2369	1.7482	1.4107	1.2814
	0.2	0.4359	0.7880	0.7891	6.1653	1.7639	1.3784	1.2474
45	0	0.4196	0.6860	0.6854	5.6961	1.2340	1.0510	0.9908
	0.1	0.4165	0.6847	0.6853	5.7884	1.2340	1.0395	0.9831
	0.2	0.4138	0.6819	0.6832	5.9617	1.2156	1.0342	0.9763
55	0	0.4013	0.6176	0.6174	5.6182	0.9816	0.8813	0.8497
	0.1	0.3979	0.6105	0.6129	5.6836	0.9781	0.8523	0.8220
	0.2	0.3913	0.6063	0.6069	5.3067	0.9628	0.8368	0.8057

Results based on 10,000 simulations

Table A6: 0.05 Rejection Frequencies, $K_2 = 30$

μ^2	$\mathcal{R}_{\varepsilon^2 z_1^2}^2$	2SLS	IJIVE1	IJIVE2	UJIVE	FEJIV	FELIM	FEFUL
25	0	0.4182	0.1490	0.1272	0.5484	0.0237	0.0572	0.0590
	0.1	0.4217	0.1461	0.1275	0.5466	0.0240	0.0553	0.0574
	0.2	0.4325	0.1495	0.1301	0.5517	0.0240	0.0556	0.0577
35	0	0.3844	0.1519	0.1326	0.5462	0.0306	0.0536	0.0565
	0.1	0.3943	0.1521	0.1299	0.5413	0.0324	0.0579	0.0600
	0.2	0.4091	0.1587	0.1368	0.5500	0.0306	0.0562	0.0585
45	0	0.3559	0.1502	0.1295	0.5491	0.0314	0.0524	0.0541
	0.1	0.3650	0.1539	0.1329	0.5434	0.0344	0.0546	0.0565
	0.2	0.3792	0.1613	0.1401	0.5504	0.0380	0.0576	0.0602
55	0	0.3437	0.1511	0.1306	0.5633	0.0370	0.0530	0.0550
	0.1	0.3331	0.1508	0.1309	0.5581	0.0396	0.0552	0.0571
	0.2	0.3624	0.1601	0.1388	0.5679	0.0421	0.0588	0.0605

Results based on 10,000 simulations

Qualitatively, the results presented in Tables A1-A6 mirror the results given in Tables 1-6 of the main paper. In particular, note that, in terms of median bias, the performance of FEJIV, FELIM, and FEFUL are uniformly better, across all our experiments, when compared to 2SLS, IJIVE1, IJIVE2, and UJIVE; but our experiments do show 2SLS, IJIVE1, and IJIVE2 to be less dispersed than the three estimators proposed in this paper. Again, perhaps the most notable difference in performance is that t-statistics based on FELIM and FEFUL have much less size distortion than t-statistics constructed from any of the other five estimators. Overall, the results here seem to indicate that allowing for some instruments to be completely irrelevant does not seem to yield results that are substantially different from the case where all instruments are taken to be relevant but weak, as long as the magnitude of the concentration parameter is kept the same in both cases.

References

- [1] Billingsley, P. (1995). *Probability and Measure*. New York: John Wiley & Sons.
- [2] Gänssler, P. and W. Stute (1977). *Wahrscheinlichkeitstheorie*. New York: Springer-Verlag.
- [3] Hall, P. and C. Heyde (1980). *Martingale Limit Theory and Its Applications*. New York: Academic Press.
- [4] Horn, R.A. and C. R. Johnson (1985). *Matrix Analysis*. Cambridge: Cambridge University Press.